Formally Assessing Cryptographic Entropy

Daniel R. L. Brown*

October 20, 2011

Abstract

Cryptography relies on the secrecy of keys. Measures of information, and thus secrecy, are called entropy. Previous work does not adequately assess the cryptographically appropriate entropy of secret keys.

This report defines several new forms of entropy appropriate for cryptographic situations. This report defines statistical inference methods appropriate for assessing cryptographic entropy.

Contents

1 Introduction **5**
- 1.1 Further Motivation . . . 5
  - 1.1.1 Roles of Entropy in Cryptography . . . 5
  - 1.1.2 Entropy Source Examples . . . 8
- 1.2 Previous Work . . . 10
  - 1.2.1 Hypothesis Testing . . . 11
  - 1.2.2 Randomness Extraction . . . 11
  - 1.2.3 Entropy Assessment . . . 11
- 1.3 Overview of this Report . . . 12
  - 1.3.1 Contributions . . . 12
  - 1.3.2 Organization . . . 12

2 Probability Models **13**
- 2.1 Formal Definition of Probability Models . . . 13
- 2.2 Equivalence, Isomorphism and Restriction . . . 15
- 2.3 Examples of Models . . . 15
  - 2.3.1 Singular, Uniform, and Deterministic . . . 15
  - 2.3.2 Independent (Identically Distributed) . . . 17
  - 2.3.3 Markov . . . 18
  - 2.3.4 Hidden Markov . . . 19
  - 2.3.5 Unrestricted . . . 20
- 2.4 Combining and Transforming Models . . . 20
  - 2.4.1 Applied Models . . . 20
  - 2.4.2 Unions of Models . . . 21
  - 2.4.3 Vacuous Extensions . . . 21
  - 2.4.4 Hulls and Composite Models . . . 21
  - 2.4.5 Products of Models . . . 22
- 2.5 Models with Extra Structure . . . 23
  - 2.5.1 Measurable and Bayesian Models . . . 23
  - 2.5.2 Metric Models . . . 24
  - 2.5.3 Non-Categorical and Poisson Models . . . 25

*Certicom Research

3 Entropy Parameters **27**
3.1 Entropy . . . 27
3.1.1 Min-Entropy . . . 27
3.1.2 Shannon Entropy . . . 28
3.1.3 Renyi Entropy . . . 29
3.1.4 Generating Series of a Distribution . . . 29
3.1.5 Working Entropy . . . 31
3.2 Modifications of Entropy . . . 32
3.2.1 Applied Entropy . . . 32
3.2.2 Contingent Entropy . . . 33
3.2.3 Contingent Applied Min-Entropy . . . 33
3.2.4 Filtered Entropy . . . 34
3.3 Sample-Dependent Entropy Parameters . . . 35
3.3.1 Sample Entropy . . . 35
3.3.2 Eventuated Min-Entropy . . . 36
3.3.3 Applied Eventuated Min-Entropy . . . 36
3.3.4 Contingent Eventuated Min-Entropy . . . 37

4 Statistical Inference **38**
4.1 Inference functions . . . 38
4.1.1 Point-valued inferences . . . 38
4.1.2 Set-valued inferences . . . 38
4.1.3 Grading-valued inferences . . . 39
4.2 Inference Methods . . . 40
4.3 Set-Valued Inference From Grading-Valued Inferences . . . 41
4.3.1 Maximally Graded . . . 41
4.3.2 Threshold Graded and Confidence Levels . . . 41
4.4 Example Gradings . . . 42
4.4.1 Likelihood . . . 42
4.4.2 Typicality . . . 42
4.4.3 Generalized Typicality and Adjusted Likelihood . . . 43
4.4.4 Calibrated Typicality . . . 45
4.4.5 Agreeability Gradings . . . 46
4.4.6 Bayesian Grading and Posterior Probabilities . . . 46
4.5 Parameter Inference . . . 46
4.5.1 Distributions to Parameter . . . 47
4.5.2 Infima Inferences . . . 47

5 Sample Statistics **48**
5.1 Induced Model . . . 48
5.2 Induced Inference . . . 48
5.3 Model-Neutral Statistics . . . 49
5.4 Sample Statistics for the Independent Probability Model . . . 50
5.4.1 Identity . . . 50
5.4.2 Frequency . . . 50
5.4.3 Partition . . . 51
5.4.4 Mode . . . 51
5.5 Statistics for the Markov Model . . . 51
5.5.1 Markov Frequency Statistic . . . 51
5.5.2 Maximum Likelihood Markov Statistic . . . 53
5.5.3 Runs Test . . . 54
5.5.4 Maximal Likelihood Min-Entropy Statistic . . . 54

6 Examples **55**
6.1 Toy Example in Independent Model . . . 55
6.1.1 Simplified Description of the Model . . . 55
6.1.2 Maximal Likelihood . . . 56
6.1.3 Threshold Inclusive Typicality . . . 56
6.1.4 Threshold Balanced Typicality . . . 58
6.1.5 Maximal Adjusted Likelihood . . . 59
6.1.6 Threshold Adjusted Likelihood . . . 60
6.1.7 Frequency Statistic Induced Inferences . . . 60
6.1.8 Partition Statistic Induced Inferences . . . 61
6.1.9 Bayesian Inference . . . 61
6.1.10 Working Entropy . . . 62
6.1.11 Applied Min-Entropy . . . 62
6.1.12 Contingent Min-Entropy . . . 63
6.1.13 Filtered Min-Entropy . . . 63
6.1.14 Sample Entropy . . . 64
6.1.15 Eventuated Min-Entropy . . . 65
6.1.16 Applied Eventuated Min-Entropy . . . 66
6.1.17 Contingent Eventuated Min-Entropy . . . 66
6.2 Polling Inference . . . 67
6.2.1 Maximum Likelihood . . . 67
6.2.2 Inclusive Typicality . . . 68
6.2.3 Balanced Typicality . . . 68
6.2.4 Adjusted Likelihood . . . 68
6.2.5 Frequency Statistic Induced Inference . . . 68
6.3 Low Sample Sizes in the Independent Model . . . 69
6.3.1 Maximal Likelihood Estimate . . . 70
6.3.2 Maximal Inclusive Typicality . . . 71
6.3.3 Maximal Balanced Typicality . . . 72
6.3.4 Frequency Statistic Induced Inference . . . 72
6.3.5 Partition Statistic Induced Inference . . . 73
6.4 Toy Examples in the Markov Model . . . 73
6.4.1 Maximum Likelihood Estimate . . . 74
6.4.2 Inclusive Typicality . . . 74
6.5 Dice . . . 75
6.5.1 The Uniform Model . . . 75
6.5.2 The Independent Model . . . 76
6.5.3 The Markov model . . . 78
6.6 Toy Model for a Ring Oscillator . . . 79
6.7 Models Based on Poisson Processes . . . 81

A Optimization Methods **84**
A.1 Karush-Kuhn-Tucker Condition . . . 84
A.2 Optimizing Non-Smooth and Non-Continuous Functions . . . 84
A.3 Model Constraints . . . 84
A.4 Optimizations for the Independent Model . . . 85

B Modeling **87**
B.1 Relaxation Approach to Modeling . . . 87
B.2 Restrictive Approach to Modeling . . . 87

C Hypothesis Testing **89**
C.1 Non-Comparative Hypothesis Testing 89
C.2 Comparative Hypothesis Testing 89

D Game-Theoretic Analysis **90**

E Estimation Theory **91**

1 Introduction

Cryptography's aim is to enable correspondents to communicate in the presence of an adversary. The correspondents generally need some form of advantage over the adversary so that they can overcome the adversary. This advantage almost always includes one or more keys known to at least one of the corresponds but unknown to the adversary. These keys are called secret (or private) keys. Most cryptographic protocols rely on such secret keys because if the adversary knew the secret key(s), then the adversary would know as much as the correspondents and could undermine the security of the protocol.

Secrecy of the keys corresponds to the lack of information that the adversary knows about the keys. Information is measured in entropy. So, the keys must have some amount of secret entropy. In general, the type of entropy appropriate for cryptography is min-entropy, which measures the difficulty of guessing the information (see §3.1.1, [X9.82], or [Lub96]).[1] In certain situations, other types of entropy are appropriate for cryptography, such as working entropy (see §3.1.5) and contingent entropy (see §3.2.2).

The entropy needed for secret keys is obtained from a *source*. Sources that have been in used or proposed for obtaining cryptographic entropy include a ring oscillator, a noisy diode, mouse movements, variances in disk read times, or even system process usages. Generally, one or more samples are obtained from one or more sources. In many cryptographic systems, these samples are accumulated into something called an entropy pool, using a deterministic process. A pool may just be a concatenation of all the values accumulated, but generally due to memory restrictions, some compression process is applied. The compression process may be as simple as a group addition, or may involve a cryptographic hash function, or may involve a randomness extraction. At some point, a value called a *seed* is extracted from this pool in order to generate a secret key. Key generation often involves a pseudorandom number generator, which takes as input the seed. All the processing from the source samples to the secret key is deterministic and cannot be deemed to add any entropy, because the deterministic algorithms in a cryptographic system cannot be kept sufficiently secret and because it can be difficult to assess the entropy of an algorithm.

In order for the samples obtained from a source to have entropy, the source should adhere to a probability distribution. This report addresses the situation that the probability distribution of the source is not exactly known. What is assumed is that the source adheres to a probability model, which means that its probability distribution at least belongs to some known set of probability distributions. Given this assumption, statistical inference is applied to assess the cryptographic entropy provided by the source.

In non-cryptographic statistical inference, an unknown probability distribution, or some characteristics thereof, is to be inferred from some sample values obtained from that distribution. Often, many assumptions about the unknown probability distribution are made, such as its being a normal distribution. Such assumptions help to make meaningful inference. Such assumptions are justified by more extensive previous statistic inferences and by the Central Limit Theorem (which usually assumes a sum of independent distributions).

1.1 Further Motivation

This section gives further motivation of how entropy is used and generated in cryptography.

1.1.1 Roles of Entropy in Cryptography

This section gives some examples of the role that entropy assessment might play in typical cryptographic applications.

1.1.1.1 Seeding Pseudorandom Number Generators A cryptographic system should typically use a well-seeded and well-designed deterministic pseudorandom number generator to generate random numbers, especially keys. The initial seed provides the cryptographic entropy to the numbers generated.

A well-designed pseudorandom number generator should ensure that the numbers generated

- appear as indistinguishable from uniform as needed,
- cannot feasibly be used to recover the internal state of the pseudorandom number generator,

[1] Shannon entropy, another type of entropy often used in communication theory, measures the compressibility of information, which is not relevant for avoid cryptographic attacks on keys (see §3.1.2) or [MvOV97].

- cannot feasibly be used, together with internal state of the pseudorandom number generator to determine past internal states. This is called backtracking resistance [NIST 800-90].

These are among the goals that pseudorandom number generators in [NIST 800-90], which, in one case, seem to met under certain assumptions [BG07].

Remark 1.1. Backtracking resistance can be necessary for forward secrecy of key agreement schemes.

Remark 1.2. Unclear responsibility for properly seeding of pseudorandom number generators can result in major problems. Suppose a manufacturer of cryptographic software implements a pseudorandom number generator but does not provide a source of entropy. If the manufacturer sets the seed to a default value, and user of the software may mistakenly generates "random" values with default seed, unwittingly believing that random number generator includes a source of entropy, then the outputs of the pseudorandom number generator should be considered to have zero entropy. Bringing attention to the task assessing entropy may at least prevent such a severe failure.

Initial seeding is often done in a fairly ideal setting such as a manufacturing site. This should enable very thorough entropy assessment.

1.1.1.2 Runtime Refreshment of Pseudorandom Number Generators

If the internal state of a deterministic pseudorandom number generators is somehow revealed to an adversary, then all its future outputs can be determined by the adversary, unless the pseudorandom number generator is refreshed with new entropy.

The property obtained by frequent refreshing is called prediction resistance in [NIST 800-90] (where refreshing is called reseeding). Barak and Halevi [BH05] call it forward security.

The entropy needed for forward security generally must be obtained during deployment in the field. In many cases, entropy in the field should be regarded as scarce. For this reason, entropy assessment is appropriate.

Entropy assessment on the sources that will be used in the field can be done both ahead of time before deployment, and also done during deployment.

Remark 1.3. It has been pointed out, in [JJSH98, BH05], that decisions made related to entropy assessment risk leaking information. As far as possible, such leakage should be incorporated into the entropy assessment. See §6.1.12, for a simplified example.

1.1.1.3 Prospective and Retrospective Assessment

A sample from a source can be used to infer something about its distribution. In some cases, the sample is just discarded, and the inference about the source is used to assess its future ability to generate entropy. This approach is *prospective* assessment.

In others cases, the sample is used for some cryptographic application, such as forming some of the input to derive a secret key. Reasons for doing this may be that entropy is believed to be so scarce that is not affordable to discard it. In this case, the assessment is *retrospective*.

Remark 1.4. Retrospective assessment can leak information to an adversary, so contingent entropy must be assessed in this case, as noted in Remark 1.3.

1.1.1.4 Computationally-Secure and Information-Theoretic Keys

Most keys deployed in cryptography undergo multiple use in such a way that one observation of such usage provides enough information to determine, using unlimited computation, the key which could then be used to compromise the remaining usage of the key.

For example, in many forms of public-key cryptography, a public key determines uniquely its corresponding private key. As another example, suppose a typical stream cipher (which attempts to generate a one-time pad from a finite key), such as the Advanced Encryption Standard used in counter mode (AES-CTR), is used once, but on a message, part of which is known to the adversary and part of which is unknown. If the known part of the message has sufficient length then, the adversary, given unlimited computation, could determine the stream cipher key, and then the whole message.

By contrast, some cryptographic protocols offer information-theoretic security. Shannon's one-time pad is the most famous example. These protocols attempt to resist an adversary with unlimited computational power. To

achieve this, they often require a very large cryptographic key, which in many cases needs to be nearly uniform. This requirement often makes them impractical.

Keys whose continued security rely on computational assumptions generally have the property of *confirmability*. An adversary who has the candidate key can confirm the key's correctness by observing the actual use of key. This means that what one considers as the entropy of key must account for an adversary who will exhaustively search for keys. The notion of working entropy from §3.1.5 can account for this.

1.1.1.5 Full and Partial Entropy Keys Some types of computational-security keys, such as public keys, are such that purely computational attacks are strictly faster than exhaustive search of all possible values of the keys.

For example, discrete logarithm keys, such as those used in Diffie-Hellman key agreement or ElGamal signatures, keys may be positive integers less than some prime q. Algorithms, such as Pollard's rho algorithm, can compute the private key about $\sqrt{q}$ steps. Schnorr [Sch01] gives strong evidence that if the private key is chosen from a random set of size $\sqrt{q}$, which allows for exhaustive search of $\sqrt{q}$ steps, no significant improvement of generic algorithms, such as Pollard rho, is any faster than about $\sqrt{q}$ steps. In other words, discrete logarithm private key seem only to require about half as much entropy as the bit length.

Other types of computational-security keys, such as symmetric encryption keys, are such that the best known computational attacks have cost similar to exhaustive search. For example, consider the block cipher Advance Encryption Standard (AES), with a key size of 128 bits. The best known attacks on the use of block cipher are exhaustively searching each possible key. Accordingly, AES-128 is generally considered to provide 128 bits of security. But providing 128 bits of security seems to require the key be perfectly uniform, meaning it has 128, or almost 128, bits of entropy. Such considerations have created enormous incentive to generate very uniform keys. Creating a nearly uniform distribution from biased entropy obtained from actual sources in environments where entropy is scarce may be too expensive.

Nevertheless, suppose that AES-128 was used with keys having only 100 bits of entropy. Presumably, then only 100 bits of security would be provided. Some chance exists that such keys could be weak. But this would seem unlikely if the keys were selected pseudorandomly, such as by the output of a hash. If 100 bits of security provide adequate protection, then the burden of producing a uniform key is lifted, and one can concentrate on providing adequate entropy.

1.1.1.6 Third Party Evaluation When a first party supplies a cryptographic product to a second party, the second party values a third party evaluation, such as [FIPS 140-1], of the cryptographic product. Third party evaluations of entropy have some difficulties:

- Proper entropy assessment requires direct access to the sources, which normally, should be not available in the cryptographic product.
- The first party has an incentive to supply the output of a deterministic pseudorandom number generator as the claimed source. To a third-party evaluator, the effect of this would be the source appears to adhere to a uniform distribution.

1.1.1.7 Organization-Level and User-Level Entropy An organization may wish to provide its members secret keys for encryption purposes, but retain a backup copy of the secret keys. In this case, the organization might use a deterministic pseudorandom number generator to generate all member secret keys. The organization may need to be quite sure about the security of the secret keys, so would likely invest considerable resources into using sufficient entropy for the seed.

Some cryptographic applications, such as personal privacy and non-repudiation, require that a user's secret key be truly secret to the user. In this case, some entropy for the user's secret key must be generated on the user's local system.

1.1.1.8 Passwords User-remembered passwords are values that a user must recall and enter into a device, usually to authenticate the user for reasons such as to obtain permission to access certain privileged information. Such passwords are typically too short to contain enough entropy to be used as cryptographic secret key in the sense

of being able to render exhaustive search infeasible. This shortness is partially based on the belief that users cannot remember high-entropy passwords.

Because of their low entropy, any data value which would allow off-line confirmation of the password guesses, such as the hash of a password, should not be made public. Otherwise password-dictionary could be applied. Password-authenticated key agreement schemes, such as SPEKE, are designed to avoid such off-line attacks. (The restriction on the exposing user-remembered passwords to off-line guessing attacks applies to both user-selected and system-generated passwords.)

Despite such usage restrictions passwords still need some entropy in order to avoid on-line guessing attacks, where an attacker can confirm password guesses on-line. To thwart on-line password attacks, usually limits on the number of failed password attempts is enforced.

Formally, the notion of working entropy, §3.1.5, can be used to reconcile the differing levels of entropy between passwords and cryptographic secret keys in more complete system. The measures used with passwords ensure that an adversaries targeting the password have a low workload, whereas for cryptographic secret keys, the adversaries can have a high workloads. The working entropy of the password and the secret keys, under low and high workloads respectively, should be similar.

1.1.2 Entropy Source Examples

This report is about assessing the cryptographic entropy of entropy sources. So, for the sake of concreteness, some examples of entropy sources are briefly discussed, which the techniques of this report could be applied to.

1.1.2.1 Operator System Processes For software to have an entropy source, one common practice is to examine the the set of processes running on the system. In complex operating systems where multiple processes are sharing processor time, it might be hoped that this kind of system information, such as the list of processes with amount of processor time each has used, has some entropy. For example, some processes may need to write to a hard disk, and disk seek times are known to vary depending on where data is located on the hard disk and other factors.

An advantage of such entropy sources is the lack of special hardware or user action.

1.1.2.2 Environmental Conditions Some systems have inputs which could be used as an entropy source. For example, a microphone can monitor the sound in the local environment.

An advantage of such an entropy source is the lack of special hardware or user action. A possible disadvantage is any adversary close enough may also have partial access to the entropy source.

1.1.2.3 User Inputs A user often supplies inputs to system, such as mouse movements or keyboard strokes. These inputs may be used as an entropy source. The inputs used for entropy may be gathered incidentally through normal use, or through a formal procedure where the user is requested to enter inputs with the instruction to produce something random.

In addition to treating user inputs as an entropy source from which entropy can be extracted to derive a secret cryptographic key, a system often relies user to directly provide a secret value, in form of a user-selected password, as in §1.1.1.8.

Passwords still require entropy, so entropy assessment of user-selected passwords is still warranted.

System-generated passwords apply a deterministic function to the output of random number generator. The deterministic function renders the random value in a more user-friendly format, such as alphanumeric. The result is a password and still needs some entropy, but in this case, the source of entropy could be some other entropy source, and still needs assessment.

1.1.2.4 Coin Flipping Perhaps the archetypal entropy source is the coin flip. A coin is thrown by a person into the air, with some rotation about an axis passing nearly through a diameter of the coin. The coin is either allowed to land on some surface or to be caught by hand. A result is either heads or tails, determined by which side is facing up.

Coin flips are often modeled such that each result is independent of of all previous results. Furthermore, for a typical coin, it is often modeled that heads and tails are equally likely.

A sequence of coin flips can be converted to a bit string by converting each result of head to a 1 and each tail to 0.

In the simples model above, the resulting bit string is uniformly distributed among all bit strings of the given length.

More skeptical models may be formulated. Firstly, it may be noted that a dishonest coin flipper could potentially cheat in certain ways. For example, the cheater may not rotate the coin on the correct axis, but rather an axis at 45° to the plane of the coin, which may cause the coin to appear to rotate, but always maintain one side closest to a particular direction in space. For another example, a cheater, with juggling skills, may be able to toss the coin with a given speed and rotation (of proper type) such that either the coin can be caught with an intended side up, or perhaps land on a surface with higher probability of landing on an intended side.

If one considers that cheating is possible, then consider that an honest coin flipper may inadvertently introduce bias in the coin flips. Indeed, in a cryptographic application relying only on coin flips for entropy, a user may need to flip a coin at least 128 times. As the user then becomes tired of repeated flips, the user may start to become repetitive and perhaps suffer from such bias.

To account for this, one could formulate a more pessimistic probability model for the coin flippers, and then do some statistical analysis comparing the pessimistic model with the actual sample of coin flips.

1.1.2.5 Dice Dice, usually cubes with numbers on the sides, have long been used in games of chance. Provided that adequate procedures are used in the rolling, the number that ends up at the top of the die, when its motion has ceased, is believed to at least be independent of previous events.

On the one hand, the roll of a die seems governed mainly by the deterministic laws of mechanics once it is released, so it may seem that all the randomness is supplied by the hand that rolled the die. On the other hand, it seems apparent that the rollers of dice cannot control the results of the die rolls,[2] and in particular source of the randomness is the actual rolling process.

The following explanation may account for this discrepancy. Each collision of the die with the ground causes it to bounce. Because the die is tumbling as it bounces, some of the rotational energy of the die may be converted into translational energy of the die, or vice versa. This conversion depends very much on the orientation of the die as it impacts the surface upon which it rolls. With each bounce, the resulting translational energy affects the amount of time before the next bounce. The amount of time between bounces affects the amount of rotation of the die, and therefore its orientation. This may mean that a small difference in orientations at one bounce results in a large difference in orientations at the next bounce. It may be that a butterfly effect is in effect, which each bounce magnify the effect of the orientation and rotation, so that the outcome of the die roll, as determined by the final orientation of the die depends on the extremely fine details in the initial orientation and motion of die. Such process are known as chaotic processes. Although technically deterministic, physical chaotic processes are hard to predict, partly because it is too difficult to obtain the necessary precision on the initial conditions to determine the final condition.

Rolling dice may be a practical way to seed a random number generator that will say, be used to generate organizational level secret keys. Rolling dice may be fairly impractical for user-level secret keys. Rolling dice is infeasible for runtime sources of entropy.

1.1.2.6 Ring Oscillator Ring oscillators have been studied as sources of entropy. See, for example, Sunar, Martin and Stinson [SMS07] or Baudet, Lubicz, Micolod, and Tassiaux [BLMT11].

Ring oscillators are essentially odd cycles of delayed not-gates. Whereas even cycles of delayed not gates can be used for memory storage, ring oscillators tend to oscillate between 0 and 1 (low and high voltage) at a rate proportional to the number of gates in the oscillator.

Since the average oscillation rate can be calculated from the number of gates and general environmental factors, such as temperature, it is only the variations in the oscillation that should be regarded as the entropy source.

[2] For example, otherwise, many games of chance would be adversely affected. That such games of chance still seem to work suggests that most people cannot control the roll of die, which suggests that some butterfly effect is occurring.

Ring oscillators are not typically available in general purpose computer systems. But they can be included in custom hardware, or even in field programmable gate arrays (FPGA).

Remark 1.5. Sunar, Martin and Stinson [SMS07] do not explicitly use the approach of this report.

1.1.2.7 Radioactive Decay Some smoke detectors use the radioactive element americium which emits alpha particles. The same method could be perhaps be used as a cryptographic entropy source, such as for the generation of organization-level secret keys.

1.1.2.8 Hypothetical Muon Measurer For the purposes of hypothetical discussion, consider an entropy source in the form a muon[3] measurer. The muon measurer provides a 32-bit measure of the speed of each muon passing through the device. On average, one muon passes through the detector per minute. Because of the underlying physics of muons, this entropy source may be viewed as providing a very robust entropy source, whose rate of entropy cannot be reduced by an adversary. [4]

This hypothetical source illustrates the task of assessing entropy. Consider the following situation. A cryptographic module testing lab receives a vendor submission of such a muon-based source. The lab accepts the general theory supplied by the vendor that each muon 32-bit speed measurement is an independent random variable with some stationary probability distribution. The lab spends about one work day to obtain 1024 speed measurements from the submitted muon detector. All speed measurements are distinct except for a single pair with the same speed.

This hypothetical example is treated formally in §6.3. For a simplified analysis, consider the following. Artificially assume that the muon speed measurements are uniformly distributed within some fixed, but unknown, subset of all possible 32-bit speed measurements. Even more simplistically, further assume just three hypotheses:[5] that this subset has size 2^{10}, 2^{30} or 2^{20}. In the first hypothesis of a 2^{10}-uniform distribution, one would have actually expected many more repetitions than just one. In the second hypothesis of a 2^{30}-uniform distribution, one would not have expected repetitions. In the third hypothesis of a 2^{20}-uniform distribution, one expects about one repetition after 2^{10} samples. Therefore, the third hypothesis seems, at least intuitively, the most consistent with the sample collected.[6]

1.1.2.9 Quantum Particle Measurement The theory of quantum mechanics implies that quantum particles, such as photons or electrons, can exist in a superposition of states under which measurement causes a wave function collapse. The theory states that wave function collapse is a fully random process independent of all other events in the universe. Under this theory, an entropy source derived from such wave function collapse would not leak any information, which is highly useful for cryptography.

Jennewein *et al.* [JAW+00] devised such a device using an attenuated light source, a beam splitter and two single photon detectors.

1.2 Previous Work

Past work on assessing cryptographic entropy has been inadequate.

[3] A muon is an elementary particle in the standard model of physics. Essentially, it is heavier version of an electron. Muons are a form of ionizing radiation, so are easily detectable, and were discovered even before the neutron. Muons are deemed difficult to produce artificially, but do occur naturally on Earth, originating from background cosmic rays (high energy protons) colliding with atoms in the atmosphere. They travel near light speed. Because of their speed and mass, they are highly penetrating, and are detectable through hundreds of meters of rock. Muons are fairly frequent at ground level

[4] This entropy source may succumb to an attack if an adversary surrounds it by other muon detectors, in which case it may be able to obtain similar speed measurements of all muons passing through the entropy source. However, this is meant only as a hypothetical example.

[5] Each of the three hypotheses is an instance of the subuniform probability model discussed in §2.3.1, but taking all three together can considered as a restriction of the independent probability model in §2.3.2.

[6] In the formal view of this report, what this simplistic analysis has done is to: assume a formal probability model, although an artificial one; use a sample statistic (§5), the number of repetitions; make a statistical inference (§4), using maximum statistic-induced likelihood, which is the that distribution with 2^{20} value is the most likely of the three distribution in the model. See §6.3.5.1 for a similar treatment.

1.2.1 Hypothesis Testing

Much past work on the assessment of randomness in cryptology, such as [FIPS 140-1] and [Mau90], has taken the form of hypothesis testing. Hypothesis testing fails to assess cryptographic entropy in several respects:

1. Zero entropy values can be contrived that pass given hypothesis tests, such as by taking the output of secure stream cipher or pseudorandom number generator, such as one of those in [NIST 800-90]. If contrived zero-entropy values can pass hypothesis tests, then it is possible that zero-entropy or too-low-entropy values can accidentally be generated that pass tests.
2. The outcome of a hypothesis test is binary: it is either a *pass* or a *fail*, not a quantity of formally assessed entropy.
3. In the formal framework of this report, hypothesis testing is the use of statistical inference in the singular uniform probability model of §2.3.1. Assuming this model is problematic because
   (a) As a singular model, meaningful inference is difficult (hence the pass or fail outcome).
   (b) Assuming the uniform model is a too strong and unrealistic assumption.
   (c) The uniform model is subject to the tying effect Remark 5.6 which requires the use of sample statistic to overcome tie-breaking effects. Poorly-chosen sample statistics rely on poorly-formulated assumptions about potential divergences from a uniform distribution.

Some developers of "true" random number generators have relied on hypothesis testing as follows. The build an entropy source with some tunable parameter. For certain values of the tunable parameter, the source may fail the hypothesis tests. For other values of the tunable parameter, the source may pass the hypothesis. The developers tune the parameters such that the entropy source has desirable properties (perhaps efficiency) and such that it passes the hypothesis tests. The entropy of such an entropy source has not been assessed.

Although hypothesis testing in cryptology has mainly been applied to the uniform model, it can be applied to any model, and as such can serve a purposes other than entropy assessment. Hypothesis testing is further discussed in an appendix to this report §C.

1.2.2 Randomness Extraction

Other past work in cryptology, such as [JJSH98], has studied how to extract almost uniformly random bit strings from random but biased bit strings. This process is called *randomness extraction* (though uniformity extraction would have been a more appropriate term).

Randomness extraction does not solve the problem of assessing entropy. Actually, randomness extraction is a step that only makes sense to be applied after entropy is assessed, because randomness extraction takes as input values with some amount of entropy.

In the general framework of this report, the entropy after randomness extraction should also be assessed using applied entropy §3.2.1.

1.2.3 Entropy Assessment

The following previous work comments on entropy assessment.

1.2.3.1 ANSI X9.82-2 The ANSI accredited standards committee X9's working group F1 recognized the need for entropy assessment. F1 began a draft standard ANS X9.82-2 [X9.82] that covers entropy sources. The author was a member of the working group F1 during this time, although not an editor of ANSI X9.82-2. The content [X9.82] varied considerably as it was edited and the working group discussed it.

No versions of ANSI X9.82-2 formalized a notion of a probability model as in this report (§2). Some drafts of ANSI X9.82-2 mentioned the hidden Markov model (see §2.3.4 for a description in this report), but this was later removed. The latest drafts restrict the probability model to the independent identically distributed model (see §2.3.2 in this report).

Statistical inference is used various drafts ANSI X9.82-2. For example, maximum likelihood estimates, with a requirement on large sample size, is used. Hypothesis testing is also used, based on *ad hoc* sample statistics, but in this case to test the hypothesis of the independent identically distributed model.

The document [X9.82] targets not only developers of entropy sources but also third party assessors, such as cryptographic module testing laboratories who have generally reported results as pass or fail. Perhaps this target audience explains why ANSI X9.82-2 was gradually simplified.

1.2.3.2 Barak and Halevi

Barak and Halevi [BH05] state:

> *... entropy estimation in general is an inherently impossible task.*

The context in which they claim impossibility of entropy estimation may not be the same as the context in which [X9.82] and this report attempt to assess entropy. Nonetheless, the strength of their statement seems to contradict at least the beliefs of the X9F1 working group.

However, even in Barak and Halevi's model [BH05], the entropy source is just assumed to have a minimum amount of entropy. This seems to be an entropy estimate of some form. Indeed, they also suggest a

> *very low static estimate for the entropy (e.g. such as 1/2 entropy bit per sample [bit])* ,

which seems inconsistent with their previous statement about the inherent impossibility.

1.3 Overview of this Report

1.3.1 Contributions

The main contributions of this report are:

- Formalization of probability models for application to cryptography.
- Several new forms of entropy that are appropriate for cryptography.
- Statistical inference methods that are appropriate for assessing cryptography entropy in a general setting.
- A paradigm that makes assumptions used to assess entropy as clear as possible.

1.3.2 Organization

The subsequent sections cover topics as follows:

- Section 2 gives formal definitions and examples of probability models.
- Section 3 gives formal definitions of cryptographic entropy.
- Section 4 gives formal definitions and examples of general statistical inference.
- Section 5 gives formal definitions and examples of, sample statistics and the resulting induced inference.
- Section 6 provides some examples of assessing entropy.
- Appendix A discusses various results from optimization theory that may be applicable to the inference methods.
- Appendix B discusses briefly some approaches to formulating a suitable probability model.
- Appendix C discusses the special case of hypothesis testing.
- Appendix D discusses the case where the adversary can influence the probability distribution.
- Appendix E discuss estimation theory, a method to assess any given inference method.

Remark 1.6. Throughout this report are scattered various remarks, such as this one. Generally these remarks are tangential to the main topic, or may refer to concepts outside the current scope, or to concepts later in the this report.

2 Probability Models

As Shannon showed, the study of information is founded on probabilities. Statistical inference involves a family of probability distributions called a probability model. This report uses statistical inference in the assessment of entropy. So, this section formally defines the notion of a *probability model* relevant to cryptography. Previous work in cryptology does not define such a general notion.

Many different probability models can be formulated under the notion of this report. Statistical inference depends on choice of probability model. Because the formal entropy assessment in this report is stated with respect to a probability model, the actual formal assessment must include the formal description of the probability model.

The formal entropy assessment can only be as appropriate as the probability model is appropriate for the given entropy source.

Remark 2.1. In this report, probabilities are used to measure an adversary's pre-existing lack of knowledge about a value which the adversary wishes to guess. An adversary may acquire extra knowledge about a specific value, which leads to the modifications of the entropy defined in §3.2, such as contingent entropy from §3.2.2 accounts for an adversary with available extra information about the outcome of a probabilistic event. Conversely, the cryptographer have more knowledge than the adversary regarding a particular, in which case eventuated entropy from §3.3.2 can be used to account for an adversary having less information about the probabilistic event than the cryptographer.

2.1 Formal Definition of Probability Models

A *probability space* Π and a *sample space* X are sets. In cryptological contexts, X is usually finite but Π is often uncountably infinite. The sample space X will be assumed to be finite, unless otherwise noted. An element $p \in \Pi$ is called a *probability distribution*, or just *distribution*, for short. An element of $x \in X$ is called a *sample*. A *probability function* for Π and X is a function

$$P : \Pi \times X \to [0,1] : (p,x) \mapsto P_p(x) \tag{2.1}$$

such that for all $p \in \Pi$, the following sum holds

$$\sum_{x \in X} P_p(x) = 1. \tag{2.2}$$

A *probability model* is a triple (Π, X, P), where Π is a probability space, X is a sample space, and P is a probability function.

Remark 2.2. For given $p \in \Pi$, write P_p for the function such that $P_p : X \to [0,1] : x \mapsto P_p(x)$. When clear from context, the function P_p may also be called a probability function.

Remark 2.3. For task of assessing entropy, the probability theory notions of an *event* and a *random variable* do not play a significant role, for the following reasons.

- An *event* corresponds to a subset of X, and a probability distribution defines the probability of an event. If X is discrete, and $E \subseteq X$, then probability of the event, under distribution p, is $\sum_{x \in E} P_p(x)$, using this report's formalism for a probability model. Because only discrete sample spaces are relevant to cryptography, the notion of event is a derivable from our formal definition of a probability model, and thus redundant.

  Usually entropy depends on the probability of a single sample, not the probability of an event. The notion of an event is incorporated to the definitions of certain kinds of entropy, such as eventuated entropy from §3.3.2, but the formal definition of probability can be stated without reference to the notion of event,
- A *random variable* is a variable taking values in the sample space X, with probabilities given by a given probability distribution p. If X is discrete, then notions such as the expected value of random variables can be expressed as $\sum_{x \in X} P_p(x)x$ using this report's formalism of a probability model. Because only discrete sample spaces are relevant to cryptography, the notion of a random variable is derivable from our formal definition of a probability model, and thus redundant.

  Usually entropy depends on the probability of a single sample, not on the expected value of random variable. Indeed, generally the values of samples have no bearing on the entropy.

  A possible role for the notion of random variables is in non-categorical probability models, see §2.5.3, where the sample values have structure that is useful in making statistic inference by way of sample statistics §5.

Remark 2.4. In cryptology, the notation $P(x)$ is often used for the probability of an event X occurring. In the notation of this report, a subscript p has been added to reflect the fact that the probability distribution p is unknown.

Remark 2.5. Cryptology also models the adversary. Three relations between the adversary and the distribution to be inferred are:

1. The adversary does not know the distribution p.
2. The adversary does know distribution p.
3. The adversary chooses the distribution $p \in \Pi$.

The three levels grant the adversaries successively more power. Some adversaries may be something intermediate between these three levels.

Remark 2.6. This report mainly focuses the second level adversarial model, where the adversary knows p, because this is most important and realistic.

Remark 2.7. The first level, which is more optimistic for the cryptographer than the second level, can be treated formally as an instance of the second level if the adversary's lack of knowledge about the distributions in the first level can be formulated in terms of probability. This would result in a new model at the second level, in which the distributions formally model the distributions of the first of the level combined with a distribution on the distributions. See Remark 2.53 for an example.

Remark 2.8. In contrast to the adversary, the cryptographer does not know p, but instead tries to infer p. So, the adversary actually has more power than the cryptographer. This may be realistic if the adversary has more access to the entropy source and can spend more effort on better statistical inference.

Remark 2.9. Over and above knowing the distribution p, an adversary may also be able to learn some information about a sample x drawn from the distribution from p. This can be accounted by using contingent entropy §3.2.2.

Remark 2.10. The third level from Remark 2.5 is discussed briefly in §D. In this case, the probability model is not controlled by the adversary, only the probability distribution. But in the formalism of choosing a probability model, the model should be chosen to encompass all the possible distribution that the adversary may be able cause to occur. The formalism need not give the influence over x, because the adversary can exert influence by exerting influence on p.

Remark 2.11. Cryptology deals with finite or discrete sample spaces X. Nevertheless, sometimes it is useful to consider continuous sample spaces X, such as a precursor model which gets subjected to a discretizing transformation. Working in the continuous model may actually simplify statistical inference, because the discretizing transformation may be discontinuous and non-smooth, making it awkward to optimize (optimization arises in the statistical inference process).

Remark 2.12. When X is a continuous space, equipped with a measure μ, then (2.2) is replaced by

$$\int_X P_p d\mu = 1, \tag{2.3}$$

and furthermore, we may extend the range of the probability function as follows:

$$P : \Pi \times X \to [0, \infty] : (p, x) \mapsto P_p(x), \tag{2.4}$$

so now $P_p(x)$ can exceed one. In this case, the function $P_p : X \to [0, \infty] : x \mapsto P_p(x)$ is called a *probability density* function.

Remark 2.13. For greater generality, do not suppose that X has a pre-existing measure. Let $M(X)$ be the collection of all measures on X. Then the model is defined by some function

$$P : \Pi \to M(X) : p \mapsto \mu_p \tag{2.5}$$

and impose the condition:

$$\int_X d\mu_p = 1. \tag{2.6}$$

In the previous example, we have $\mu_p = P_p\mu$. In the case of a finite or countably infinite set X, then the measure μ_p can be defined from the usual probability function P_p via

$$\mu_p(Y) = \int_Y d\mu_p = \sum_{y\in Y} P_p(y). \tag{2.7}$$

2.2 Equivalence, Isomorphism and Restriction

If (Π, X, P) is a probability model then two probability distributions $p, q \in \Pi$ are *equivalent* in the model (Π, X, P) if $P_p(x) = P_q(x)$ for all $x \in X$, which can be written $p \equiv q$.

Given two probability models (Π, X, P) and (Θ, Y, Q), the models are *isomorphic* if there exists functions $\beta : \Pi \to \Theta$ and $\gamma : \Theta \to \Pi$ and a bijective function $b : X \to Y$ such that for all $(p, x) \in \Pi \times X$ it is true that $P_p(x) = Q_{\beta(p)}(b(x))$ and for all $(q, y) \in \Pi \times Y$ it is true that $Q_q(y) = P_{\gamma(q)}(b^{-1}(y))$. In other words, if one simply relabels the elements of probability space and the sample space, one obtains an isomorphic model. Also addition or removal of equivalent distributions yields isomorphic models.

Remark 2.14. Entropy, see §3, of a probability distribution p is invariant under isomorphism. Therefore, strictly speaking, from a cryptological perspective, it suffices to only consider probability models up to isomorphism. That said, certain probability models may include the possibility of numeric relationship between components of x, in which case, an arbitrary isomorphism would render this relationship arbitrary, and therefore likely more difficult to process, and in particular, to make inferences about.

Henceforth, models will be considered only up to isomorphism, unless otherwise noted.

Remark 2.15. If (Π, X, P) is probability model and $z \in X$ is such that $P_p(z) = 0$ for all $p \in \Pi$, then we say that z is *non-occurring*. Otherwise z will be said to be *occurring*. Modifications of models by addition or removal of non-occurring sample values may be considered weakly isomorphic.

Given two probability models (Π, X, P) and (Θ, Y, Q), the latter is a *restriction* of the former if $Y = X$ and $\Theta \subset \Pi$ and, for all $p \in \Theta$ and $x \in Y$, it is true that $Q_p(x) = P_p(x)$. Conversely, (Π, X, P) is a *relaxation* of (Θ, Y, Q). Similarly, (Θ, Y, Q) is more restrictive than (Π, X, P), and (Π, X, P) is less restrictive than (Θ, Y, Q).

If (Π, X, P) is a probability model, and $p \in \Pi$ and $x, y \in X$ and $P_p(x) = P_p(y)$, then x and y are said to be *equiprobable* at distribution p.

Remark 2.16. Equiprobable distributions have equal typicality, §4.4.2.

If x and y are equiprobable at all $p \in \Pi$, then x and y are said to be *equilikely* in the model.

Remark 2.17. All non-occurring sample values are equilikely.

Remark 2.18. Likelihood functions are defined in §4.4.1. Equilikely sample values x and y have the same likelihood functions: $L_x = L_y$.

2.3 Examples of Models

Statistical inference can be conducted over any probability model. For the sake of concreteness, some example models are given in this section.

Some of these example models are cryptographically appropriate models for certain entropy sources.

2.3.1 Singular, Uniform, and Deterministic

Excluding the degenerate case of $\Pi = \varnothing$, the most restrictive a probability model can be is if $|\Pi| = 1$. In this case, the probability model is *singular*.

An example of a singular probability model is the *uniform* probability model where $P_p(x) = 1/|X|$ for all x. More generally, any model isomorphic to the uniform model is also called a uniform model. Also, given any finite set X,

there is a uniform model on X, which will be written as $u(X)$. Up to isomorphism, the uniform model is determined by the cardinality of X, so this uniform model may be referred to as the $|X|$-uniform model. For example, the 6-uniform model implies a uniform model with $|X| = 6$, a model sometimes attributed to a roll of a cube die.

When clear from context, *uniform* is applied applied to distributions, not just models. Specifically, for any probability model (Π, X, P), as distribution $p \in \Pi$ is the uniform distribution if $P_p(x) = 1/|X|$ for all $x \in X$. If (Π, X, P) has a uniform distribution, then it is a relaxation of the uniform model $u(X)$.

Remark 2.19. The uniform distribution p is generally the most cryptographically secure probability distribution on the sample space, because it has the maximum possible min-entropy, $\log_2 |X|$, see §3.1.1, of all distributions on the space X, and because it is usable as one-time pad.

Another important example of a singular probability model is a *deterministic* model. In the case, $\Pi = \{p\}$ and there is some $x_0 \in X$, such that $P_p(x_0) = 1$ and $P_p(x) = 0$ if $x \neq x_0$.

As with the term *uniform*, when clear from context, the term *deterministic* applies to individual probability distributions, not just models. Specifically, if (Π, X, P) is a model, $p \in \Pi$, and $P_p(X) = \{0, 1\}$, then p is a deterministic distribution. If p is deterministic and $P_p(x) = 1$, then the notation $p = p_x$ will sometimes be used. So, $P_{p_x}(x) = 1$ and $P_{p_x}(y) = 0$ for $y \neq x$.

Remark 2.20. A deterministic distribution is the least cryptographically secure distribution, because a deterministic distribution has zero min-entropy, which means that an adversary knowing the distribution can guess the sample value.

Remark 2.21. For a given probability model, it is worth being well aware of the set of deterministic distributions that it contains, since when one obtains a sample value x such that the deterministic distribution on x belongs to the model, inferring that the distribution could be deterministic is very compelling, since sample x and deterministic distribution on x are as perfect a fit between a sample and distribution as can be.

Remark 2.22. A *pseudodeterministic* model is a model that contains a deterministic distribution p_x for each $x \in X$. Inference in a pseudodeterministic model can be problematic, because, given sample x, the distribution p_x is the best inference, which is deterministic and has zero entropy. Any inference method that includes p_x among the inference to be made from x and takes the minimum min-entropy of the inferred distributions gives an inferred min-entropy of zero.

Remark 2.23. A *fatalistic* model on a sample space X is the most restricted pseudodeterministic model: $\Pi = \{p_x : x \in X\}$ with $P_{p_x}(x) = 1$ and $P_{p_x}(y) = 0$ for $y \neq x$. The fatalistic model is also the least restricted model in which all distributions have zero min-entropy.

The fatalistic model is worse than a deterministic model because it cannot be rejected by hypothesis testing. For example, if a deterministic model with $\Pi = \{p_x\}$ is wrong, then it is possible that a sample obtained can be y with $y \neq x$, in which case the model will be seen to have been wrong.

Assuming a model that is not fatalistic can be strongly justified if, upon scrutiny by a real adversary, the adversary gains no advantage. A weaker justification for a non-fatalistic model for an entropy source is hypothesis testing on a non-fatalistic model..

Remark 2.24. Intermediate to uniform and deterministic are a family of singular probability models, called *subuniform* models. For integers m, N with $1 \leqslant m \leqslant N$, a (m, N)-subuniform model is such that $|X| = N$, and $P_p(x) \in \{0, 1/m\}$ for all $x \in X$, which implies that P_p is nonzero on a subset of cardinality m, and it is constant on this subset. The N-uniform model is the (N, N)-subuniform model. The deterministic model is the $(1, N)$-subuniform model.

Similarly, subuniform distributions are distributions p in any probability model (Π, X, P) such that $P_p(X) = \{0, 1/m\}$.

Remark 2.25. Singular models, especially the uniform model, have been used in hypothesis testing, such as the previous work [FIPS 140-1].

Statistical inference, §4, is the process of inferring something about p from a given value of x. In a singular model, only one value of p is possible. The inference to be made in a singular model therefore takes the form of a pass or fail, or perhaps some grading of the fit between an observed sample x and the model's single distribution.

Singular models are generally inappropriate for assessing cryptographic entropy, because they are generally already assume a value of the entropy and because the limited form of the inference (pass or fail).

Even if the uniform model has reasonable belief for some source, such as the entropy source devised by Jennewein *et al.*

[JAW+00], and if hypothesis testing is the one's only goal (so, for some reason, one is not trying formally assess entropy), the uniform model is still somewhat unsuitable in a formal sense, as discussed below.

The formal unsuitably is because the uniform model requires the use of sample statistics, see §5, to overcome the tying effect in uniform distributions, see Remark 5.6. As such, sample statistics when applied to hypothesis, are essentially trying to detect the possibility that hypothesis is false. In other words, the sample statistic is testing if some other hypothesis is more realistic. But sample statistic do not formally state what the alternative hypothesis.

This report therefore proposes an alternative approach to hypothesis testing, which is outlined in §6.5 and §B.

2.3.2 Independent (Identically Distributed)

Another probability model is the (m, N)-*independent (identically distributed)* model:

$$\Pi = \left\{p = (p_0, \ldots, p_{m-1}) : p_i \in [0,1], \sum p_i = 1\right\} = [0,1]_1^m \tag{2.8}$$

$$X = \{x = (x_0, \ldots, x_{N-1}) : x_i \in \{0, 1, \ldots, m-1\}\} = \mathbb{N}_m^N \tag{2.9}$$

$$P_p(x) = \prod_{i=0}^{N-1} p_{x_i}. \tag{2.10}$$

In the abbreviated notations given above: $[0,1]$ means the interval of real numbers between 0 and 1, inclusive; $\mathbb{N}_m$ means $\{0, 1, \ldots, m-1\}$; S^m means the set of m-tuple with entries in S; and S_1^m means the subset of S^m such that the sum of the entries in the m-tuple is one.

The parameter m is called the *width*, and the parameter N is called the *length*. A distribution in (m, N) may be referred to as an *independent* distribution on the sample space $\mathbb{N}_m^N$.

In this model, the parts x_i of x are restricted to be individual random variables with *identical and independent distributions*. There is no restriction, however, on the this common distribution.

Remark 2.26. The (m, N)-independent model is a relaxation of the m^N-uniform model because taking $p = (1/m, \ldots, 1/m)$ causes $P_p(x) = 1/m^N$ for all $x \in X$.

Remark 2.27. In reference to this model, we may sometimes call p a probability vector and x a sample vector.

Remark 2.28. The $(2, N)$-independent probability model may to be an appropriate way to model a coin tossed N times if the coin's probabilities of landing heads or tails are independent and stable.

Remark 2.29. The independent model is also a relaxation of the deterministic model in the following sense. Fix some $i \in \{0, \ldots, m-1\}$. If $p_i = 1$, then $P_p(x) = 1$ if $x = (i, i, \ldots, i)$ and otherwise $P_p(x) = 0$. These are the only deterministic distributions in the independent model.

Remark 2.30. The $(m-1, N)$ independent model is a restriction, up to isomorphism, of the (m, N) independent model.

Remark 2.31. For $N \geqslant 2$, the independent model is not pseudodeterministic.

Remark 2.32. Given the (m, N)-independent model, it is natural to consider the following function $f : X \to [0,1]^m$ defined by the relation $f(x)_i = |\{j : x_j = i\}|/N$. This is the *frequency* function, and it is easily seen to be the maximum likelihood inference (§4.3.1 and §4.4.1) $\hat{p}(x)$ for x.

Also, the $P_p(x)$ is a function of $f(x)$, so if $f(x) = f(y)$, then x and y are equilikely. Furthermore, x and y are equilikely only if $f(x) = f(y)$.

The number of different value that f takes is $\binom{m+N-1}{m} = \frac{(m+N-1)!}{m!(N-1)!}$. For $m \gg N$, this number is approximately $\frac{m^{N-1}}{(N-1)!}$, so the average size of an equilikely class is about $\frac{(N-1)!}{m}$. For $N \gg m$, this number is approximately $\frac{N^m}{m!}$, so the average size of an equilikely class is about $\frac{m!m^N}{N^m}$.

Remark 2.33. Randomness (uniformity) extraction is a known method in cryptology, an example of which follows. Suppose that (Π, X, P) is an (m, N)-independent model and that f is the frequency function defined above. Define a function $g : X \to \mathbb{Z}$ as follows: $g(x)$ is the index of x amongst the list of y with $f(y) = f(x)$, with the list being sorted lexicographically. Let

$e(x) = (f(x), g(x))$, and let $Y = [0,1]^m \times \mathbb{Z}$. Then $e : X \to Y$ is an injection. Define a probability model (Π, Y, Q) such that $Q_p(e(x)) = P_p(x)$ for all $x \in X$. The probability model is (Π, Y, Q) is partially subuniform in the following sense: for a fixed $f \in [0,1]^m$, the set $\{Q_p(f, g) : g \in \mathbb{Z}\}$ contains zero and has cardinality at most two. As such, what one can do is *extract* the value $g(x)$ from x, and essentially ensure that it appears to abide by the uniform model, of a size that may be calculated from x using multinomial coefficients. During the process, considerable valuable entropy contained in x may be lost, with the gain in uniformity usually being a theoretical goal. Entropy is often more important than uniformity, and in some systems, entropy is too scarce to sacrifice.

Remark 2.34. Uniformity extraction can be more generally viewed as taking advantage of the presence of equilikely sample values. Given a sample value x, one may know that, in the assumed probability model, that x is equilikely with some number of other sample values. It follows that the index of x among this set of equilikely values has a uniform distribution.

Remark 2.35. A relaxation (Π', X, P') of the independent model (Π, X, P) can be formed by taking $\Pi' = \Pi \times \Sigma_X$ where Σ_X is the set of all permutations of X. Then let $P'_{(p,s)}(x) = P_p(s(x))$. This relaxation implies allows an arbitrary structure on the sample space, and then distributions which are independent identically distributed within the given structure. So, call this model the *structureless* independent model.

This relaxed model has many equivalent distributions. For example, if t is any permutation of the set $\mathbb{N}_m$ and is adapted to X by application to each entry, and adapted to Π by re-ordering of the entries, then $(p, s) \equiv (t(p), t \circ s)$. It may be that for almost of all of the space Π', the latter equivalences determine the entire equivalence classes, because the function $P'_{(p,s)}$ determines (p, s) up to the transformation by t as described above, but there are exceptions. For example, if p is corresponds the uniform on $\mathbb{N}_m$, then $(p, s) \equiv (p, t)$ for any permutation s and t of X.

This structureless independent model is pseudodeterministic, so inference in this model is infeasible. But a (common) product, as in §2.4.5, of structureless independent models may not be pseudodeterministic, allowing distributions to be inferred.

2.3.3 Markov

Another probability model commonly considered is the (m, N) *Markov* model. The sample space $X = \mathbb{N}_m^N$, which is the same as the (m, N) independent model. The probability space Π consists of pairs $p = (v, M)$, where $v : \{0, 1, \ldots, m-1\} \to [0,1]$ whose values sum to one, and $M : \{0, 1, \ldots, m-1\}^2 \to [0,1]$, whose values, when summed with any fixed first argument, total to one. More compactly, $\Pi = [0,1]_1^m \times ([0,1]_1^m)^m$. We may view v and M as a vector and a matrix. Then

$$P_p(x_0, \ldots, x_{N-1}) = v(x_0)M(x_0, x_1)M(x_1, x_2) \ldots M(x_{N-2}, x_{N-1}) \tag{2.11}$$

This allows x_{i+1} to depend on x_i according to the matrix M. As in the independent model, the parameter m will be called the width and the parameter N will be called the length.

Remark 2.36. The $(m-1, N)$ Markov model, is a restriction, up to isomorphism, of the (m, N) Markov model.

Remark 2.37. The (m, N) Markov model is, up to isomorphism, a relaxation of the (m, N) independent model. Distributions $(v, M) \in \Pi$ such that all the rows of M are identical to v give rise to distribution equivalent to a distribution in the (m, N) independent model with $p = v$.

Remark 2.38. The deterministic distributions in the Markov model's probability space are those that take constant values x such that x has the form $u(vw)^y v \in X$, where ab indicates concatenation of sequence and the sequence uvw has no repeated elements.

Remark 2.39. In the $(2, 4)$ Markov model, the distributions $(0, 1, 0, 0)$ and $(0, 0, 1, 0)$ are equilikely. More generally, in the $(2, N)$ Markov model, define a function $f : X \to \mathbb{Z}^3$, such that

$$f(x) = \left(x_0, \quad \sum_{i=0}^{N-1} x_i, \quad \sum_{i=0}^{N-2} |x_i - x_{i+1}| \right). \tag{2.12}$$

If $f(x) = f(y)$, then x and y are equilikely. A function with the same property for the (m, N) Markov model is given in §5.5.

2.3.4 Hidden Markov

The (h, m, N) hidden Markov model may be thought of as being built on top of a Markov model. The sample space $X = \mathbb{N}_m^N$, as in the (m, N) independent model. The probability space Π has elements p that are triples (v, M, Q) of functions:

$$\begin{aligned} v &: \{0, 1, \ldots, h-1\} \to [0, 1] \\ M &: \{0, 1, \ldots, h-1\}^2 \to [0, 1] \\ Q &: \{0, 1, \ldots, h-1\} \times \{0, 1, \ldots, m-1\} \to [0, 1] \end{aligned} \tag{2.13}$$

such that the values of v sum to one, and such that for each fixed value of the first argument, the values of M sum to one, and likewise for Q. In abbreviated notation, $\Pi = [0, 1]_1^h \times ([0, 1]_1^h)^h \times ([0, 1]_1^m)^h$.

As in the independent model, the parameter m will be called the width and the parameter N will be called the length. The parameter h will be called the *height*.

An auxiliary set S, consisting of *hidden states*, is defined as the same set as the sample space in the (h, N) independent model, so $S = \mathbb{N}_h^N$. The probability function is defined as:

$$P_p(x) = \sum_{s \in S} v(s_0) Q(s_0, x_0) \prod_{i=1}^{N-1} M(s_{i-1}, s_i) Q(s_i, x_i) \tag{2.14}$$

Remark 2.40. The hidden states are distributed according to a Markov model. Each sample component is distributed from a component of the hidden state, essentially by taking one step in a different Markov model but modified so that initial state of the modified model is determined by the hidden state.

Remark 2.41. The (h, m, N) hidden Markov model is a restriction of the (h', m', N) hidden Markov model if $h' \geqslant h$ and $m' \geqslant m$.

Remark 2.42. If $m \geqslant h$, then the (h, m, N) hidden Markov model is a relaxation of the (h, N) Markov model.

Remark 2.43. The deterministic distributions in the (h, m, N) hidden Markov model are those that have non-zero probability at just one value x of the form $x = u(vw)^y v \in X$, where ab indicates concatenation of sequences a and b and uvw is a sequence of length at most h.

The number of terms in the sum (2.14) is h^N, which may be too many for practical computations, even for modestly small values of h and N. However, $P_p(x)$ can be computed efficiently, by an algorithm known as the forward-backward algorithm, or forward-Viterbi algorithm.

A main idea of this forward algorithm is to use matrix multiplication. Let M also denote the $h \times h$ matrix naturally corresponding to the function M. Let V be an $h \times h$ matrix which is a diagonal matrix whose entry in position (s_0, s_0) is given by $v(s_0)$. Similarly, let Q_i be another diagonal $h \times h$ matrix whose entry in position (s_i, s_i) is given by $Q(s_i, x_i)$. Then compute the matrix product

$$P = VQ_0MQ_1M \ldots MQ_{N-1} \tag{2.15}$$

The sum of all of its entries gives $P_p(x)$. This formulation is generally more efficient than (2.14), because each matrix multiplication uses about h^3 multiplications, so the total number of multiplications is at most about Nh^3 rather than the Nh^N use in the literal formulation of (2.14).

Remark 2.44. The general Viterbi algorithm also takes x and computes the hidden state s of highest probability on the condition of resulting in x. The Baum-Welch algorithm [BPSW70] uses the Viterbi algorithm to infer, from x, the maximal likelihood estimate for the parameters (v, M, Q).

Remark 2.45. If the $h \geqslant m, N + 1$, then the (h, m, N) hidden Markov model is pseudodeterministic.

2.3.5 Unrestricted

The unrestricted model is defined as follows. Identify each probability distribution p with its probability function P_p. (Recall, p and q are equivalent distributions if and only if $P_p = P_q$, so this identification characterizes p up to isomorphism.) For a finite set X, define the unrestricted model $U(X) = (\Pi, X, P)$ on X, by setting Π to be the set of all legal probability functions on X.

Remark 2.46. The $(m, 1)$ independent probability model is an unrestricted probability model. Similarly, the $(m, 2)$ Markov model is also an unrestricted model.

Remark 2.47. An artificial restriction of the unrestricted model (Π, X, P), is to restrict the probability space to a subset Π' of those distributions p whose min-entropy, §3.1.1, is at least some desired threshold. More generally, such a restriction can be applied to any given model. In the approach of this report, this is like assuming what one wants. Such a model may describe a goal of a cryptographic system, but does not truly describe an actual source.

In the approach of this report, starting from a given model (Π, X, P), the choice of p is beyond the cryptographer's control. In particular, no action of the cryptographer, even given information about a sample value x, can force p to some subset Π'. The cryptographer may be able to take advantage against an adversary of properties of the observed sample x, and these can be accounted for in the modified definitions of entropy from §3.2, but they still do not change p.

2.4 Combining and Transforming Models

This section gives examples of transforming and combining models.

2.4.1 Applied Models

Let $f : X \to Y$ be a function from sample space X to sample space Y. Given probability model (Π, X, P), the function f creates an *applied* model (Π, Y, Q) given by

$$Q_p(y) = \sum_{x:f(x)=y} P_p(x) \tag{2.16}$$

A special case is when f is a permutation of X to itself. In this case, the resulting model is a permuted model.

Remark 2.48. A hidden Markov model can be viewed as a restriction of the applied model derived from a Markov model as follows. Specifically, the (h, m, N) hidden Markov model is an applied model derived from a subset of the $(hm, N+1)$ Markov model. In the Markov model the sample space consists of sequences of pairs $((s_0, x_{-1}), (s_1, x_0), \ldots, (s_N, x_{N-1}))$. The applied function sends this sample to $(x_0, \ldots, x_{N-1})$. The values s_j correspond to the hidden states in the hidden Markov model. A Markov distribution (v, M) can be chosen to give the resulting applied distribution for each distribution in the hidden Markov model. (Some Markov distributions will give a distribution that is not described by Markov model.)

Remark 2.49. Remark 2.48 may suggest that an approach to making inferences in the hidden Markov model by making inferences in a Markov model in order to make inferences . Some potential difficulties with such an approach are:

- The parameters of the Markov model may be much larger,
- The observed samples from which one wants to make an inference are the resulting applied function to the actual samples,
- The probability space is actually a restriction of the Markov model distribution, which must be accounted for in making appropriate inference.

Remark 2.50. The (m, N) Markov model may also be viewed as an applied model derived from a restriction of the $(m^{(m+1)}, N)$ independent model. Consider the samples in the independent model to be a sequence $(g_0, \ldots, g_{N-1})$ where each entry g_i is a function $g_i : \{-1, 0, \ldots, m-1\} \to \{0, 1, \ldots, m-1\}$. The applied function f is:

$$f(g_0, \ldots, g_{N-1}) = (x_0, \ldots, x_{N-1}) \tag{2.17}$$

where $x_0 = g_0(-1)$ and $x_j = g_j(x_{j-1})$ for $j \geqslant 1$. If the entries g_i in the independent model are selected as functions distributed in a way consistent with the Markov distribution (v, M), then the $f(g_0, \ldots, g_{N-1})$ has the distribution equivalent to that given by the Markov distribution.

2.4.2 Unions of Models

If (Π, X, P) and (Θ, X, Q) are models with a common sample space, then a left union of their models is the model $(\Pi \cup \Theta, X, R)$ can be formed as:

$$R_p(x) = \begin{cases} P_p(x) & \text{if } p \in \Pi \\ Q_p(x) & \text{if } p \notin \Pi \end{cases} \tag{2.18}$$

Left unions are non-commutative: they are sensitive to the order of the two models over which the left union is applied. When the two models are consistent on the intersections of their probability spaces, their left union is same regardless of the order.

The disjoint union of an indexed family of sets S_i for $i \in I$ is defined as

$$\biguplus_{i \in I} S_i = \{(i, s_i) : i \in I, s_i \in S_i\}. \tag{2.19}$$

Given an indexed family of models, (Π_i, X, P^i) with a common sample space and index set I, their disjoint union is the model $(\biguplus_{i \in I} \Pi_i, X, Q)$, where

$$Q_{(i,p)} = P^i(p, x). \tag{2.20}$$

Remark 2.51. As an example, take any model (Π, X, P) with X finite. One can take the family of models which are permutations of this model. So, the indices are the $|X|!$ permuation of the set X. The disjoint union of these model is the *dispermuted* model of derived from the original model (Π, X, P).

Suppose that one has one has a source whose samples derived from some unobservable samples underlying source. Suppose that the underlying source is known with certainty to adhere to a given model. The way the observed samples are derived from the underlying unobserved samples is by measurement process, which is known to be deterministic and lossless. This derivation process is otherwise completely unknown to the cryptographer, but may be known to the adversary. The dispermuted model describes these circumstances.

If the original model has any deterministic distributions, then the resulting dispermuted model is pseudodeterministic. For example, the dispermuted model derived from the independent model is pseudodeterministic.

2.4.3 Vacuous Extensions

If (Π, X, P) is a probability model, and $X \subset Y$, the vacuous extension of model to Y is the model (Π, Y, Q) such that:

$$Q_p(y) = \begin{cases} P_p(y) & \text{if } y \in X \\ 0 & \text{if } y \notin X. \end{cases} \tag{2.21}$$

Vacuous extensions together with disjoint unions allow mixture of models with different sample space sizes.

2.4.4 Hulls and Composite Models

Given a family of distributions, and a distribution on the family, one can formulate the *weighted mean* of the family of distributions. Precisely, if (Π, X, P) and (Θ, Π, Q) are models such that all $q \in \Theta$ are *discrete* in the sense the set $\text{supp}(q) = \{p : Q_q(p) > 0\}$ is countable, then a model (Θ, X, R), the *hull* model, can be defined with probability function R

$$R_r(x) = \sum_{p \in \Pi} Q_r(p) P_p(x) \tag{2.22}$$

and we can think if r as a distribution on X under probability function R acting like the *weighted mean* of the distribuitons p in Π. When necessary to distinguish the role of distribution r in the hull model (Θ, X, R) from the role of r in the model (Θ, Π, Q), we may write $\bar{r}$ for the distribution in the hull model, and refer to $\bar{r}$ as the mean distribution.

Remark 2.52. If (Θ, Π, Q) admits non-discrete distributions, then the probability function Q is replaced by a probability measure, and the hull model can defined using a measure, provided that the function $p \mapsto P_p(x)$ is measurable under measure $q \in \Theta$.

Remark 2.53. Remark 2.7 referred to converting a first level model of adversary who does not know the distribution into an adversary who does know the distribution. Formalizing this conversion requires formulation of what the adversary does not know about the distribution in the first level model. So, a second level probability model on the first level probability space is what is needed in our formalism. The resulting model on the sample space at the second level can be achieved using the hull of models.

Remark 2.54. For a specific example of the conversion in Remark 2.53, suppose that the first level model is the $(2, N)$-independent model. The probability vectors $(p_0, p_1) \in \Pi$ are characterized by the value of $p_1 \in [0, 1]$. If the adversary does not know p_1 in the formal sense that from the adversary's perspective p_1 is uniformly distributed in the interval $[0, 1]$, then we can formalize this into a model at the second level, in which the adversary knows the distribution. In this case, because we supposed a single distribution for p_1, namely the uniform distribution on the interval $[0, 1]$, we obtain with a singular model, at the second level. (If we had suppose a family of distributions for p_1, we would have obtained a non-singular model.)

Formally, the first level model (Π, X, P) is the $(2, N)$ independent model, while model on the probability space is (Θ, Π, Q) is a singular model, continuous sample space model, and is the a continuous variant of uniform model. Becuase this second model is singular, it has just a single distribution, which will write as $\bar{p}$. The distribution $\bar{p}$ is not discrete. The hull model is $(\Theta, X, R) = (\{\bar{p}\}, \{0, 1\}^N, R)$ for R defined as follow. The sum from (2.22) must be replaced by an integral, because $\bar{p}$ is not discrete. Using p_1 to represent elements of Π, the formula for R is given by the integral

$$R_{\bar{p}}(x) = \int_0^1 p_1^W (1 - p_1)^{N-W} dp_1 \tag{2.23}$$

where W is the sum of the entries of x, its Hamming weight. This integral evaluates to $Q_{\bar{p}}(x) = \frac{1}{N+1} \frac{1}{\binom{N}{W}}$, which can be proven using integration by parts.

In the model (Θ, X, R) the adversary's best strategy is to guess $x = (0, 0, \dots)$ or $x = (1, 1, \dots, 1)$. The min-entropy of the distribution is $\log_2(N + 1)$, which is rather low compared bit length N of x.

The fact this model is singular means that it may be rather unrealisitic. If however, the evidence of this singular model is very strong for the given source, then it should be deemed to provide low entropy.

The *composite* model derived from model (Π, X, P) and model (Θ, Π, Q) is defined as model $(\Theta, \Pi \times X, S)$ with probability function:

$$S_r(p, x) = Q_r(p) P_p(x) \tag{2.24}$$

The hull model may be obtained from the composite model as an applied model, using the function $f : \Pi \times X \to X : (p, x) \mapsto x$. When necessary distinguish the role of distribution r in the composite model $(\Theta, \Pi \times X, R)$ from the role of r in the model (Θ, Π, Q), we may write $\hat{r}$ for the distribution in the hull model, and we refere $\hat{r}$ as the *composite* distribution.

2.4.5 Products of Models, Multiple Sources, and Repeated Sampling

Given two probability models with a common probability space, (Π, X, P) and (Π, Y, Q), their product probability model can be formed as $(\Pi, X, P) \times (\Pi, Y, Q) = (\Pi, X \times Y, R)$ defined by

$$R_p(x, y) = P_p(x) Q_p(y) \tag{2.25}$$

Iterating such product, as in a power, starting from the unrestricted model, allows (m, N) independent probability model to be expressed as $U(\mathbb{N}_m)^N$ where $\mathbb{N}_m = \{0, 1, 2, \dots, m - 1\}$.

Remark 2.55. In many applications of probability, one works with repeated samples, also known as Bernoulli trials, of some distribution. By formalizing the product model above, then we can make this assumption about the independence of repeated trials explicit.

For example, suppose a cryptographic source is built from a ring oscillator. We may perhaps model individual runs, starting from system boot, of the ring oscillator by the hidden Markov model. To model the fact that at each system re-boot, the ring oscillator starts afresh and independently, then we can take the product of the hidden Markov models.

Remark 2.56. In cryptology applications, where we mainly care about min-entropy, then it can be noted that the min-entropy of p in the product model is the sum of its min-entropy in the underlying models.

A different type of product of models is as follows: the product of (Π, X, P) with (Θ, Y, Q) is the model $(\Pi \times \Theta, X \times Y, R)$ where $R_{(p,q)}(x, y) = P_p(x)Q_q(y)$. To distinguish this product from the previous, one can call this product the *mixed* product, and call the previous product the *common* product.

Generally, inference over a mixed product of models is equivalent to doing inference in each model separately. By comparison, inference over a common product, because it is a more restrictive model, cannot be so separated. In particular, mixed products do not lead to stronger inferences, where as common products do.

Remark 2.57. In cryptology applications, to derive a key, one may often attempt to use multiple different sources. If one models these sources as independent and unrelated, then one can use the mixed product model to jointly model these independent sources.

Remark 2.58. When two models have a common probability space, then both the common and mixed product are definable. The common product will be a restriction of mixed product, and the mixed product will be a relaxation of the common product.

Remark 2.59. The mixed product is actually a common product of the models $(\Pi \times \Theta, X, P')$ and $(\Pi \times \Theta, Y, Q')$ where $P'_{p,q}(x) = P_p(x)$ and $Q'_{p,q}(y) = Q_q(y)$. These two models in the common product are equivalent to the original two in the mixed product.

Remark 2.60. The mixed product of pseudodeterministic models will be pseudodeterministic, whereas the common product may not be.

2.5 Models with Extra Structure

In the general definition of probability model (Π, X, P), the probability space Π and the sample X are treated as sets with no structure. In non-cryptographic applications it is often convenient or desirable to equip Π or X with some additional structure. Even if such additional structure does not have immediate application to entropy assessment, it may be useful in the process of establishing evidence for a given probability model. So, such structures are discussed briefly in this subsection.

2.5.1 Measurable Models, Bayesian Models and Prior Probabilities

We may sometimes need to have a measure μ on the probability space. This measure can then be considered as a supplementary component of the probability model. Sometimes we may also want measures also to be defined on various subsets of Π, such as lower dimension slices.

For the independent probability model, the Markov model and the hidden Markov model, the probability spaces are defined as intersections of a real hypercube with certain hyperplanes. Therefore, one possible family of measures can be built on these probability space inherited from the Euclidean metric.

More generally, since the models we consider are defined over a fixed sample space, then we may regard probability distributions as probability vectors. Therefore, the probability space can be mapped to a subset of $\mathbb{R}^{|X|}$ (with equivalent distributions mapped to the same point). Then various metrics and measures available on $\mathbb{R}^{|X|}$ can be induced onto the probability space Π without any theoretical difficulty.

Finally, one may want to normalize the measure μ so that $\mu(\Pi) = 1$. Then the measure μ may be thought of as defining probabilities on the probability distributions. These probabilities of probabilities are often used in *Bayesian inference*, in which case they are sometimes called the *prior probabilities*.

Remark 2.61. For a probability model (Π, X, P) that is a restriction of model (Σ, X, Q) which also has an associated measure μ, then it may be possible to induce a measure on $\Pi \subseteq \Sigma$. This induced measure can then be normalized so to give a total of one over the whole set Π.

Remark 2.62. For a probability model (Π, X, P) that is a restriction of model (Σ, X, Q), then one may also define a measure μ on (Σ, X, Q), such that $\mu(S) = 0$ if S is disjoint from Π. In other words, the measure μ on (Σ, X, Q) is imposing a restriction to the model (Π, X, P).

Remark 2.63. A possible objection to measures, and more specifically prior probabilities, is that prior probabilities must be assumed. However, any probability model must be assumed. Indeed, assuming prior probabilities, as illustrated above, is a generalization of assuming a specific probability model.

Remark 2.64. A probability model with associated prior probabilities, that is, a quadruple (Π, X, P, μ) where μ is a measure on Π such that $\mu(\Pi) = 1$, can sometimes be converted into a singular probability model (Σ, X, Q) as follows, by taking the hull model from §2.4.4.

Explicitly, the probability space Σ is singular with a single element σ, and the probability function is computed as follows

$$Q_\sigma(x) = \int_\Pi L_x d\mu \tag{2.26}$$

where L_x is the function defined on Π such that for $p \in \Pi$ we have $L_x(p) = P_p(x)$. For this to be well-defined at x, the function L_x must be measurable and integrable over Π. In this view, the original probability distributions may be viewed as hidden state associated with the singular model (Σ, X, Q), much like the hidden states in the hidden Markov model.

Remark 2.65. The probability space of the unrestricted model $U(X)$ can be parametrized in a measure-preserving way by elements of the unit hypercube of dimension $|X| - 1$. (In terms of a metric, the probability space $U(X)$ is a simplex, so no metric-preserving transformation can map it to a hypercube.) Mapping the probability space to a hypercube may be a convenient transformation for certain algorithms, even if measure-preservation is not a goal. Other probability models, including the independent model and Markov model, have probability spaces which are spanned by subspaces similar to the unrestricted model.

For example, if $X = \{0,1\}^t = \{(x_0, \ldots, x_{t-1}) : x_i \in \{0,1\}\}$ and $U(X) = (\Pi, X, P)$, then Π with the natural measure can be mapped to $H = [0,1]^{\bigcup_{j=0}^{t-1}\{0,1\}^j} = \{(u_{()}, u_{(0)}, u_{(1)}, \ldots, u_{\underbrace{(1,1,\ldots,1)}_{t-1}}) : u_y \in [0,1]\}$ while preserving measure, as follows. Use the notation $x \oplus u$ for $x \in \{0,1\}$ and $u \in [0,1]$ to mean u if $x = 0$ and $1 - u$ if $x = 1$. Let $\pi : H \to \Pi$ be defined such that:

$$P_{\pi(u_{()}, u_{(0)}, \ldots)}(x) = \prod_{j=0}^{t-1} x_j \oplus u_{(x_0, \ldots, x_{j-1})}. \tag{2.27}$$

For another example, suppose that $X = \{0, 1, \ldots, d\}$ and the unrestricted model is $U(X) = (\Pi, X, P)$. Let $H = [0,1]^d = \{(u_1, \ldots, u_d) : u_i \in [0,1]\}$. Define $\pi : H \to \Pi$, by:

$$P_{\pi(u_1, \ldots, u_m)}(x) = (1 - u_x^{1/x}) \prod_{y=x+1}^{d} u_y^{1/y} \tag{2.28}$$

with the convention that $(1 - u_0^{1/0}) = 1$.

2.5.2 Metric Models

One can associate a metric with a probability model: so that some probability distributions may be then viewed as closer to each other than others.

Remark 2.66. In some cases, a metric on a space can be used to build a measure on the space. So a metric model can be converted into a measurable model.

Remark 2.67. This report allows models in which Π is infinite, even a continuum (for the example, the independent model). The statistical inference methods used to assess entropy in this report produce an optimization problem defined over the probability space Π.

Optimization methods over a continua generally make use of some coordinatization of the infinite set. Such a coordinatization generally implies a metric. Furthermore, optimization methods over continua generally make use of gradients, which are defined with respect to a metric.

Sometimes one wants a metric on the probability space for purposes other than just applying optimization methods. In particular, previous work in cryptology, such as [Lub96], has used the following metric, often known as

statistical distance:

$$d(p,q) = \tfrac{1}{2} \sum_{x \in X} |P_p(x) - P_q(x)|. \tag{2.29}$$

This metric gives the maximum probability for any algorithm to correctly guess whether a single sample x originates from probability distribution p or q. In previous work on cryptology, this has been used as a measure of some candidate distribution p, say for a cryptographic key, closeness to an ideal distribution q, say a uniform distribution. The idea is that an adversary of unlimited computational power will not be able to distinguish from p and q, except with probability bounded by the statistical distance. In most cases, the application of this notion is take some source of biased entropy and produce from it a distribution p that is close to some ideal q.

The metric (2.29) can be regarded as based on an L^1 norm. Other previous work, such as [BH05] in cryptography has considered a metric based on L^∞ norm, which can be written as

$$d_\infty(p,q) = \max_{x \in X} |P_p(x) - P_q(x)|. \tag{2.30}$$

2.5.3 Non-Categorical and Poisson Models

Most of the example models above (uniform, independent, Markov and hidden Markov) would be called *category data* models because the only structure of the sample values x upon which the probability model depends is the division of x into a sequence of components whose values have no significance, in the sense of isomorphism of models. In particular, although the components in the above models were stated in terms numerical values, the models themselves make no assertions to be made about any numerical relations. Just to be clear, an individual distribution may treat the numerical values differently, but the model as a whole does not. In other words, the models have automorphisms permuting the orders of the numerical values of the components (and also the order of the components in the case of the independent and uniform models).

In category data models, it makes no sense to perform arithmetic operations on the numerical values of the categories. In non-categorical models, such as heights of people, operations such as expected (average) values of sample for a given distribution make sense. Much of statistics is, naturally, devoted to such non-category models. For example, the central limit theorem suggests that in, say, the independent model the average of the sample components has a distribution that approaches a normal curve. However, in a category data model, such an average makes little sense.

An informal reason to focus in cryptology on category data models is that entropy of a source is more important that the structure of the sample values. It is not numerical relations between components of the sample that are important, but rather the entropy of the distribution. As such, numerical patterns may be irrelevant to main cryptological goal of assessing entropy. More precisely, rather than attempting to find a numerical pattern, for example, a trend towards linear growth in a sequence of sample values, which may be of tremendous importance in non-cryptological studies, in cryptology we would rather adopt a Markov model, to accommodate the possibility of a evolving pattern, for the purposes of assessing entropy. So, in cryptology, we certainly want to recognize patterns, as indicators of lower entropy, but we wish to pay no heed to whether the patterns have numerical significance.

Nevertheless, certain cryptographic sources may be reasonably expected to have components that have numerical values and relationships. By incorporating such numerical relations directly into the probability models, one may be able to make better inferences. Therefore, despite the consideration above, one may still want to use non-category data models in cryptology.

Remark 2.68. Just as an example, suppose that $\Pi = \mathbb{R}$ and that $X = \{0,1,2\}^N$ and that $P_p(x) = 1$ if $x_i = \lfloor 1 + \sin(pi) \rfloor$ for all i, and otherwise $P_p(x) = 0$. As motivation, suppose that a (poorly designed) ring oscillator follows such a probability model.

Remark 2.69. In the model from Remark 2.68, all distributions are deterministic. The most sensible inference (§4) ought to, given x return the one distribution p with non-zero probability on x. (Given only partial information, say y, about x, the one might infer a set of distributions which have non-zero probability on the possible x for the given y.)

Because each distribution is deterministic, the entropy is zero. Formally, such source provides no security at all in this model.

Intuitively, one might argue that there is entropy to be found in the choice of p, which, before generation of x, is unknown not only the cryptographer but also to the cryptanalyst. Therefore, there may be some security.

But in this case, to formally attribute min-entropy, one must assign probabilities to each probability distribution. This may be an example of where one wants a measure on the probability space.

More appropriate would be to revise the probability model so that former probability space becomes some intermediate state value over which one assigns probabilities, a little like the Hidden Markov model has a hidden state. Although the hidden state here is a continuum, and therefore requires a continuous probability distribution, it nevertheless induces a discrete probability distribution on the finite sample space, and therefore the min-entropy of the samples is well-defined (whereas the min-entropy of the hidden state would be infinite). (See Remark 2.64.)

2.5.3.1 Poisson Models The *Poisson* probability model is the model (Π, X, P) in which the probability space is $\Pi = [0, \infty)$ (the set of non-negative real numbers), and the sample space is $X = \mathbb{Z}_{\geqslant 0} = \{0, 1, 2, 3, \dots\}$ (the set of non-negative integers), and the probability function is given by the formula:

$$P_p(x) = \frac{e^{-p} p^x}{x!}. \tag{2.31}$$

Each individual distribution in a Poisson model is called a Poisson distribution. Poisson distributions are well-studied distributions in probability theory. Poisson distributions have the property that the sum of two independent random variables with Poisson distributions p and q gives another random variable with Poisson distribution $p + q$.

Another closely related model is the *Poisson process* model. This model has probability space $\Pi = [0, \infty)$ as in the Poisson model, and sample space X' consisting of the countable subsets of real numbers. The uncountable size of the sample space makes this is continuous model, so it does not have a probability function, but rather a probability measure. To each distribution q there is associated a measure μ_q on the space X'. The cryptographically relevant properties of this measure are as follows.

- For any interval $[a, b]$, define a function

  $$c_{a,b} : X' \to X : x \mapsto |x \cap [a, b]| \tag{2.32}$$

  where $X = \mathbb{Z}_{\geqslant 0}$ as in the Poisson model. The function $c_{a,b}$ is called the count function because it measures how many of the values resulting the Poisson process land in the given interval. Then $c_{a,b}$ can be used to define a discrete model, where for $x \in X$, probability function is defined as

  $$P_q(x) = \mu_q(c_{a,b}^{-1}(x)), \tag{2.33}$$

  so the measure of the set of all those $x' \in X'$ with count x. The Poisson process model is such that for distribution $q \in \Pi$ in the Poisson process model, the resulting distribution is a Poisson distribution on X. Moreover, it is the Poisson distribution with $p = q(b - a)$.

- For any two disjoint intervals $[a, b]$ and $[c, d]$, the two resulting Poisson distributions obtained from the two corresponding count functions are independent of each other.

A Poisson process model can be used to model radioactive decay sources, for example.

3 Entropy Parameters

A *probability parameter*, or *distribution parameter*, or just *parameter* when clear from context, on a probability model (Π, X, P) is a function $r : \Pi \to R$. The set R can be called the *parameter space*, and elements of R can be called *parameter values*, or when clear from context just *parameter*.

A *sample-dependent parameter* on a probability model is a function $f : \Pi \times X \to R$. The set R will also be called a *parameter space*.

Remark 3.1. In general applications of statistics, a parameter of the probability distribution may be some particularly important unknown quantity figuring in some random process. The important quantity may need to be separated from some less important components of the probability distribution, in which case the parameter may be called a signal, and the remaining contribution to the probability distribution may be called the source. For a specific example, consider a poll of voters. The signal may be the proportion of the total population's voting preferences, and the noise (source) may be the method used to select the sample poll and the accuracy of the poll responses.

In cryptology, certain parameters are crucially important. Unlike typical applications of statistical inference, the focus of cryptology is not on the nature of some unknown quantity nominally related to the sample space, but rather on the probability distribution itself. More precisely, rather than being able to make some useful assertions about the nature of the random variable modeled by x, the goal in cryptology is for p to be such that making predictions about x is difficult.

So, the cryptologically relevant parameters of p are measures of how difficult p makes predicting x. Generally, such a measure quantifies the amount of information that the adversary lacks about x. Measures of information are called *entropy*. Several different types of entropy are discussed below.

3.1 Entropy

This section defines various types of entropy.

3.1.1 Min-Entropy

In cryptology, the parameter of greatest interest is *min-entropy*. For a probability model (Π, X, P), and given probability distribution $p \in \Pi$, the min-entropy of p is defined to be

$$H_\infty(p) = -\log_2 \max_x P_p(x) = \min_x(-\log_2 P_p(x)). \tag{3.1}$$

The units of min-entropy are called *bits*.

Remark 3.2. An adversary who knows p, and wants to guess x, should guess the value of x for which $P_p(x)$ is maximal.

Remark 3.3. The 2^N-uniform distribution p has N bits of min-entropy.

Remark 3.4. If $H_\infty(p) = u$ in the model (Π, X, P) and $H_\infty(p) = v$ in the model (Π, Y, Q), then $H_\infty(p) = u + v$ in the product model $(\Pi, X \times Y, P \times Q)$.

Remark 3.5. In the (m, N) independent model, a distribution $p = (p_0, \ldots, p_{m-1})$ has min-entropy $H_\infty(p) = -N \log_2(\max_{j=0}^{m-1}(p_j))$.

Remark 3.6. In the (m, N) Markov model, a distribution $p = (v, M)$ has min-entropy

$$H_\infty(p) = -\log_2 \left(v \odot \underbrace{M \odot M \odot \cdots \odot M}_{N-1 \text{ copies of } M} \odot u \right) \tag{3.2}$$

where: $\odot$ is defined just like normal matrix multiplication except that instead of computing dot products of rows with columns by summing the pairwise products of the elements, one takes the maximum of the pairwise products[7]; and u is the vector with all entries equal to one.

[7] For example, $\begin{pmatrix} 3 & 2 & 1 \end{pmatrix} \odot \begin{pmatrix} 3 \\ 5 \\ 7 \end{pmatrix} = 10$.

This formula may be viewed as a special case of the Viterbi algorithm. The naive approach would have been to compute $P_p(x)$ for all m^N possible values of x and compute the maximum. Formula (3.2) can be computed with cost proportional to at most m^2N or $m^3 \log_2(N)$.

Remark 3.7. Using the fact that, for any finite set of non-negative real numbers $\{a, b, \ldots, c\}$, the maximum is expressible as a limit: $\max(a, b, \ldots, c) = \lim_{r\to\infty} \sqrt[r]{a^r + b^r + \cdots + c^r}$, allows the modified matrix multiplication $\odot$ from Remark 3.6 to expressed using conventional matrix multiplication and a limit by the formula:

$$H_\infty(p) = -\log_2 \lim_{r\to\infty} \sqrt[r]{v_r (M_r)^{N-1} u} \tag{3.3}$$

where for a vector or matrix A, the notation A_r means the corresponding vector or matrix with all entries raising to the power of r.

Remark 3.8. Under composition of distributions (2.24), min-entropy obeys the inequality

$$H_\infty(\hat{r}) \geqslant H_\infty(r) + \max_p H_\infty(p) \tag{3.4}$$

using the notation from §2.4.4.

3.1.2 Shannon Entropy

Although *Shannon* entropy is not suitable for formally assessing cryptographic entropy, it appears often in previous work, such as means to characterize a uniform distribution. The definition of Shannon entropy is included in this section and its unsuitability as a form of cryptographic is explained.

For a probability model (Π, X, P), and given probability distribution $p \in \Pi$, the Shannon entropy of p is defined to be

$$H_1(p) = -\sum_{x\in X} P_p(x) \log_2 P_p(x). \tag{3.5}$$

Remark 3.9. Suppose that $|X| = 2^m + 1$ and for some $x_0 \in X$, the probability distribution p is such that $P_p(x_0) = 1/2$ and $P_p(y) = 1/2^{m+1}$ for $y \neq x_0$.. Then $H_1(p) = 1 + m/2$, but $H_\infty(p) = 1$, which illustrates the cryptological unsuitability of Shannon entropy for assessing non-uniform distributions.

Remark 3.10. Suppose that $X = \{0, 1, \ldots, 2^{128} - 1\}$, that $P_p(0) = 2^{-7}$, and that $P_p(x) = \frac{1-2^{-7}}{2^{128}-1}$ for $x \neq 0$. Then $H_\infty(p) = 7$, but $H_1(p) \approx 127.066$. So, the distribution p has one bit less of Shannon entropy less than the uniform distribution on X. If a key were generated from this distribution, an adversary would have probability 2^{-7} of determining the key by deriving it from $x = 0$. If about 2^7 keys are generated independently from under this, then the probability is about 0.63 that one of the keys will derived from $x = 0$.

Remark 3.11. Suppose that $X = \{0, 1, \ldots, 2^{128} - 1\}$, that $P_p(0) = 2^{-15}$, and that $P_p(x) = \frac{1-2^{-15}}{2^{128}-1}$ for $x \neq 0$. Then $H_\infty(p) = 15$, but $H_1(p) \approx 127.997$. So, the distribution p has 0.003 bits less of Shannon entropy less than the uniform distribution on X. If a key were generated from this distribution, an adversary would have probability 2^{-15} of determining the key by deriving it from $x = 0$. If about 2^{15} keys are generated independently from under this, then the probability is about $1 - e^{-1} \approx 0.63$ that one of the keys will derived from $x = 0$.

Remark 3.12. From an adversary's perspective it may seem relevant to calculate the expected number of guesses need to determine the sample value. The distribution in Remark 3.9 has fairly high expected value for the number of guesses. But this high expected value does not reflect the risk to the user of choosing a weak value. A better notion is working entropy (§3.1.5).

Remark 3.13. Many references in cryptology refer to Shannon entropy, but do not mention min-entropy.

Remark 3.14. The 2^N-uniform distribution p has N bits of Shannon entropy. More generally, a distribution p on X is uniform if and only if $\log_2 |X|$ bits of Shannon entropy.

Remark 3.15. If $H_1(p) = u$ in the model (Π, X, P) and $H_1(p) = v$ in the model (Π, Y, Q), then $H_1(p) = u + v$ in the product model $(\Pi, X \times Y, P \times Q)$. So, like min-entropy, Shannon entropy is multiplicative.

Remark 3.16. Shannon entropy is useful for non-cryptographic parts of information theory, as in the following examples.

- Shannon entropy measures how compressible a sequence of values x distributed according to p is. (In other words, taking an a probability distribution in the independent model.) The idea is to encode x using approximately $-\log_2 P_p(x)$ bits, which can be realized using something like arithmetic encoding.
- Shannon entropy is useful to measure the amount of error rates that can be detected or corrected with error correcting codes.

Remark 3.17. Shannon entropy is preserved under composition of distributions (2.24) in the sense that

$$H_1(\hat{r}) = H_1(r) + \sum_p Q_r(p) H_1(p) \tag{3.6}$$

using the notation from §2.4.4.

3.1.3 Renyi Entropy

A common generalization to min-entropy and Shannon entropy is *Renyi* entropy [Rén60]. For a probability model (Π, X, P), given probability distribution $p \in \Pi$, and real number $t > 0$ with $t \neq 1$, the Renyi entropy at order t for probability distribution p is defined to be

$$H_t(p) = \frac{1}{1-t} \log_2 \sum_{x \in X} P_p^t(x). \tag{3.7}$$

As $t \to 1$, Renyi entropy approaches Shannon entropy. As $t \to \infty$, Renyi entropy approaches min-entropy. Therefore min-entropy and Shannon entropy can be considered as special cases of Renyi entropy, of orders ∞ and 1, respectively. Renyi entropy is known to be a decreasing function of t. In particular, min-entropy is always at most Shannon entropy.

Remark 3.18. As $t \to 0$, Renyi entropy tends to $\log_2 |\{x : P_p(x) \neq 0\}|$, sometimes called the *Hartley* entropy.

Remark 3.19. Sometimes the term Renyi entropy refers just to H_2, Renyi entropy of order two. This is sometimes known as the collision entropy, because it is related to the rate at which the distribution, when taken over two samples, repeats.

Remark 3.20. If $H_t(p) = u$ in the model (Π, X, P) and $H_t(p) = v$ in the model (Π, Y, Q), then $H_t(p) = u + v$ in the product model $(\Pi, X \times Y, P \times Q)$.

Remark 3.21. Although, as noted above, Renyi entropy decreases with t, bounds within a factor exist in the other direction, with the bound $\frac{1}{2} H_2 \leqslant H_\infty$ being of some interest to cryptology.

Remark 3.22. For any fixed value of H_∞, there exists probability distributions with H_1 arbitrarily large, provided that the size of the sample space X is sufficiently large, as noted in Remark 3.9.

3.1.4 Generating Series of a Distribution

If (Π, X, P) is a probability model and X is a finite or countable set, the distributive generating series for distribution p is given by:

$$D(p; z) = \sum_{x \in X} z^{-\log_2(P_p(x))} \tag{3.8}$$

The series $D(p; z)$ may be viewed as a function of z or as an element of the ring of Hahn series $\mathbb{Z}[[z^{\mathbb{R}}]]$, whose elements have the form $\sum_{w \in W} a_w z^w$ where W is a well-ordered subset of the non-negative reals, and a is an arbitrary function from W to non-zero integers.

The expression $D(p; z)$ on the left side of (3.8) does not refer explicitly to the probability function P. When this it is necessary to distinguish such series for different probability model, a subscript can be used as $D_P(p; z)$.

The condition that $\sum_{x\in X} P_p(x) = 1$ implies that $D(p; -\frac{1}{2}) = 1$. The distributive generating series determines Renyi entropy of order $t \neq 1$ via

$$H_t(p) = \frac{1}{1-t} \log_2 D(p, 2^{-t}). \tag{3.9}$$

Conversely, the Renyi entropies $H_t(p)$ as a function of t determine the distributive generating series $D(p; z)$ as a function of z.

In a mixed product of two models, as in §2.4.5, a distribution has the form (p, q) where p and q are distributions in the models over which the product is taken. In this case:

$$D((p, q); z) = D(p; z)D(q; z) \tag{3.10}$$

In a common product, a single distribution p is associated with different probability functions, P, Q and R in the notation of §2.4.5, and

$$D_R(p; z) = D_P(p; z)D_Q(p; z) \tag{3.11}$$

Given a distribution p which is known to be an unknown permutation of a common N^{th} power of another unknown base distribution p, taking the N^{th} root of the distributive series of the powered distribution, which can be done using the binomial theorem in the Hahn series formula, can be used to determine the distributive series of the base distribution.

Remark 3.23. Suppose that we somehow know with certainty that source is governed by a model (Π, X, P) is some permutation of the $(3, 2)$ independent model for some fixed but unknown permutation. Further suppose that we are able obtain very many independent samples from the model (Π, X, P) and thereby infer with fairly strong confidence that the distribution p describing the source satisfies:

x	$P_p(x)$
00	0.112
01	0.1089
02	0.1155
10	0.1155
11	0.1056
12	0.1056
20	0.112
21	0.1225
22	0.1024

(3.12)

to some approximation. The distributive series is then for p in the presumed model is:

$$D(p, z) = z^{3.28771} + 2z^{3.24332} + z^{3.19892} + 2z^{3.15843} + 2z^{3.11404} + z^{3.02915} \tag{3.13}$$

Note that the min-entropy is given by the exponent of the leading term 3.28771, but question is here is more general than to infer the min-entropy.

The base distribution has distributive series $\sqrt{D(p, z)}$, which in this example can be calculated directly by insepction by halving the exponents of the terms with coefficient 1.

$$\sqrt{D(p, z)} = z^{1.64386} + z^{1.59946} + z^{1.51457} \tag{3.14}$$

Exponentiating the exponents gives that the underlying $(3, 2)$ independent model is $p = (0.35, 0.33, 0.32)$, or it could be some other permuation of p.

The permutation π relating our given model to the $(3, 2)$ independent model with distribution p given above, in the sense that $P_p(x) = R_p(\pi(x))$ where R is the probability function of the $(3, 2)$ independent model is:

x	$\pi(x)$
00	20
01	11
02	01
10	10
11	12
12	12
20	02
21	00
22	22

(3.15)

The utility of determining π may be for somthing like uniformity extraction. In this case, it would have been possible to determine p without actually computing $\sqrt{D(p,z)}$.

Remark 3.24. The general method of using the binomial theorem to compute $\sqrt{D(p,z)}$, would works as follows:

$$\begin{aligned}\sqrt{D(p,z)} &= z^{\frac{3.02915}{2}}(1+2z^{0.0848889}+2z^{0.129283}+z^{0.169778}+2z^{0.214172}+z^{0.258566})^{1/2}\\ &= z^{1.51457}\sum_{n=0}^{\infty}\binom{1/2}{n}(2z^{0.0848889}+2z^{0.129283}+z^{0.169778}+2z^{0.214172}+z^{0.258566})^n\\ &= z^{1.51457}(1+\frac{1}{2}(2z^{0.0848889}+2z^{0.129283}+\dots)+\dots)\\ &= z^{1.51457}+z^{1.59946}+z^{1.64386}\end{aligned} \tag{3.16}$$

where the ... above would all cancel. A practical implementation would need to establish some criteria for ignoring certain terms.

3.1.5 Working Entropy

Another generalization of min-entropy is *working entropy*[8] defined by

$$H_{(w)}(p) = \min_{x_j}\left(-\log_2\sum_{j=1}^{\lfloor 2^w\rfloor}P_p(x_j)\right) \tag{3.17}$$

where the minimum is taken over arbitrary sequences x_j of distinct values, and, as a convention, let $P_p(x)=0$ if $x\notin X$, which allows the sum to be well-defined for all w. (Unlike earlier notation in this report, the index j in x_j above here does not refer to the j^{th} entry in a sample vector x but rather the j^{th} sample vector in a sequence of sample values.) The variable w is the workload and is measured in bits. Min-entropy is working entropy at workload of zero bits.

Remark 3.25. A uniform distribution over a set of size 2^h has working entropy of $h-w$ bits at a workload of $w\leqslant h$ bits for each positive integer 2^w. At workloads of $w\geqslant h$, the working entropy is zero.

Remark 3.26. For any distribution p on a finite (or countable) sample space, $H_{(\infty)}(p)=0$, so working entropy is zero at infinite workload.

Remark 3.27. For a finite sample space $H_{(H_0(p))}(p)=0$, so working entropy is zero at workload equal to the Hartley entropy.

Remark 3.28. Working entropy is a non-increasing function of w.

Working entropy is most relevant in the situation where an adversary to a cryptographic scheme sees cryptographic operations resulting from application of the secret entropy, and can gain some value by determining such entropy, usually in the form of compromising past or future cryptographic operations. The seen operations may permit the adversary willing to make calculations approximately proportional 2^w to confirm 2^w guesses at the secret value, such as by attempting the cryptographic computation with guessed value of the value.

Remark 3.29. Consider a bit string x of length 128 with the following nearly uniform distribution. The probability that $x=0^{128}$ is 2^{-80}, and probability of any other given value of x is $\frac{1-2^{-80}}{2^{128}-1}$. This distribution has min-entropy of only 80 bits, which is 48 bits less than the min-entropy of the uniform distribution on the same sample space. At a workload of 48 bits, the working entropy of this distribution is about 79 bits, which is only one about bit less than that of the uniform distribution. So, the effect of aberrant spikes in probability distribution on working entropy is reduced at high workloads.

[8] Being a natural generalization of min-entropy, working entropy has probably been defined in previous work.

Remark 3.30. Working entropy as defined in (3.17) is not a continuous function of w. The following variant is more continuous as a function of w, by adding an extra term to the sum, as follows.

$$H'_{(w)}(p) = \min_{x_j} -\log_2 \left((2^w - \lfloor 2^w \rfloor)\, P_p(x_{\lfloor 2^w \rfloor + 1}) + \sum_{j=1}^{\lfloor 2^w \rfloor} P_p(x_j) \right) \tag{3.18}$$

Remark 3.31. A remarkable property of Renyi entropies is additivity over (products of) independent distributions (Remarks 3.4 and 3.20. A weaker property of working entropy may be:

$$H^{(\Pi, X\times Y, P\times Q)}_{(v+w)}(p) \lessapprox H^{(\Pi,X,P)}_{(v)}(p) + H^{(\Pi,Y,Q)}_{(w)}(p) \tag{3.19}$$

where the superscripts indicate the model over which the working entropy is defined.

Remark 3.32. If a bound in the opposite direction to the bound in (3.19) held, then assessing the working entropy of multiple independent sources might be feasible.

3.2 Modifications of Entropy

This section describes some modified versions of entropy, which are useful to address certain realistic cryptographic circumstances.

3.2.1 Applied Entropy

In some cryptographic applications, there is a function $f : X \to Y$ such that given a sample value x, only the value $f(x)$ is used as a key. In this case, the adversary need merely guess $f(x)$. If $f(x)$ is easier to guess than x, then high min-entropy of x does not suffice for security for $f(x)$.

Remark 3.33. For example the function f could be: a key derivation function, a hash function, a debiasing function, a uniformity extractor, part of an entropy pooling function, a way to leave some of the sample x available for other use, or just a way to allow the possibility that an adversary may have a side channel to the source that depends on not just the used portion but a large portion.

A function $f : X \to Y$ and probability model (Π, X, P) induce the *applied* model (Π, Y, Q) where

$$Q_p(y) = \sum_{x : f(x)=y} P_p(x) \tag{3.20}$$

The *applied* entropy of the p in the model (Π, X, P) under the function f is the entropy of p in the applied model (Π, Y, Q). One can consider applied min-entropy or applied working entropy. The applied min-entropy works out to be:

$$H_{f(\infty)}(p) = \log_2 \max_{y\in Y} \sum_{x : f(x)=y} P_p(x) \tag{3.21}$$

Remark 3.34. Cachin [Cac97] introduced the notion of *smooth entropy*. This notion blends the notions of entropy and randomness (uniformity) extraction. In the terminology of this report, consider each function $f : X \to Y$ for each size of Y, and forms the applied model. Moreover, each applied model is equipped with a metric on the probability space. The smooth entropy is parametrized by some distance value, a *smoothness bound* in the metrics. The *smooth entropy* of p at distance d, is the highest entropy of a uniform distribution that is within distance d of an applied distribution obtained from p. Smooth entropy expresses the potential amount of uniform entropy that can be extraced from a source randomness (uniformity) extraction.

In the view of this report, the adversary knows the entropy extraction algorithm. Therefore, the adversary's ability to guess the applied value is still best described by the applied entropy, not the smooth entropy.

3.2.2 Contingent Entropy

In many cryptographic applications, given a sample value x, there exists some function f such that $z = f(x)$ is revealed to the adversary. This section defines *contingent* entropy to address this situation.

Remark 3.35. As an example, suppose that x is modeled by a Markov model, and that a nonce value, such as the initialization vector in the cipher-block chaining mode, is derived from x and sent in the clear. If the nonce does not reveal the whole of x, then it may still be possible to use x to derive a key. Contingent min-entropy measures the upper possible limit of how securely this can be done.

Remark 3.36. One may also assume that there is some side-channel leaking information about x. So the adversary can learn $f(x)$.

First some preliminaries are given. Two function $f : X \to Z$ and $g : X \to Y$ are said to be supplementary if the function $f \times g : X \to Z \times Y : x \mapsto (f(x), g(x))$ is injective.

Remark 3.37. The intuition is that the function g expresses at least enough information about x to recover x from $f(x)$.

The *contingent* entropy of distribution p in the model (Π, X, P) on the condition that the adversary learns $f(x)$ is the infimum of the applied entropy of p in the applied models (Π, Y, Q) over all g supplementary to f. One can consider contingent min-entropy or contingent working entropy.

A function g supplementary to f that seems to minimize entropy (of most types), is the following. Assume that X is sorted in manner such that P_p is a non-increasing function in the order. Let $g(x)$ be the index of x in the set $f^{-1}(f(x))$. This supplementary g induces a model where p has min-entropy:

$$H_{\infty|g}(p) = -\log_2 \sum_{z \in Z} \max_{x : f(x) = z} P_p(x). \tag{3.22}$$

Henceforth we will take (3.22) as the definition of contingent min-entropy.

Remark 3.38. An intuitive explanation for (3.22) is that whatever z turns out to be, the adversary will choose the most likely x corresponding to that z. So, the sum of maxima represents, over the choice of x, the adversary being successful.

Remark 3.39. Equations (3.22) with (3.21) only differ essentially in that maximization and summation operator have been swapped.

Some cryptographic applications, such as public-key cryptography, reveal an *injective* function f of the private key, such as the public key. (This often in symmetric-key cryptography, when a known message longer than the secret symmetric key is encrypted.)

Theoretically, this means that the contingent entropy of the key goes to zero. This does not imply zero security, because such functions f are generally believed to be one-way functions: inverting f is believed to computationally infeasible.

Incorporating a measure of infeasibility would allow one to define a notion of *computational min-entropy*. In this report, focus is deliberately restricted to the task of gathering sufficient entropy before the application of any cryptographically complex function, such as one-way functions.

3.2.3 Contingent Applied Min-Entropy

Let $f : X \to Y$ and $g : X \to Z$ be functions. Let (Π, X, P) be a probability model. Suppose that an adversary will learn $z = g(x)$ and but that only $f(x)$ will be used in the generation of a key. The *contingent applied min-entropy* of $p \in \Pi$, under f and g, is defined to be:

$$H_{f(\infty)|g}(p) = -\log_2 \sum_{z \in Z} \max_{y \in Y} \sum_{x \in f^{-1}(y) \cap g^{-1}(z)} P_p(x). \tag{3.23}$$

To generalize (3.23) from min-entropy to working entropy of workload w, let the maximization operator be indexed over $\lfloor 2^w \rfloor$-element subsets y of Y. An alternative formula for contingent applied min-entropy is as follows:

$$H_{f(\infty)|g}(p) = -\log_2 \max_{\alpha : Z \to Y} \sum_{x : \alpha(g(x)) = f(x)} P_p(x). \tag{3.24}$$

The function α represents the following strategy of an adversary: given $z = g(x)$ guess that $f(x) = \alpha(z)$. The sum in (3.24) is the probability that the adversary guesses $f(x)$ correctly from $z = g(x)$. The entropy is then the negative base two logarithm of highest success probability of any adversary. The alternative formula equals the original because:

$$\begin{aligned}\max_{\alpha:Z\to Y}\sum_{x:\alpha(g(x))=f(x)}P_p(x) &= \max_{\alpha:Z\to Y}\sum_{z\in Z}\sum_{x\in g^{-1}(z)\cap f^{-1}(\alpha(z))}P_p(x)\\ &= \sum_{z\in Z}\max_{y\in Y}\sum_{x\in g^{-1}(z)\cap f^{-1}(y)}P_p(x).\end{aligned} \tag{3.25}$$

3.2.4 Filtered Entropy

Suppose that (Π, X, P) is a probability model; and suppose that $Y \subset X$ is a subset of the sample space. Suppose that a sample x is drawn from with some distribution $p \in \Pi$. If $x \notin Y$ then x is rejected. Otherwise x is accepted. If $0 \neq \sum_{y\in Y} P_p(y)$, then the result is the Y-*filtered* probability model (Π, Y, Q) is defined by

$$Q_p(y) = \frac{P_p(y)}{\sum_{z\in Y} P_p(z)}. \tag{3.26}$$

The Y-*filtered* entropy of p is the entropy of p in the filtered model. Explicitly, the filtered min-entropy of distribution p is

$$\left(\min_{y\in Y} -\log_2 P_p(y)\right) + \log_2\left(\sum_{y\in Y} P_p(y)\right) \tag{3.27}$$

Remark 3.40. From an implementation perspective, filtered entropy fails to account for the cost of rejecting values of the source, and it fails to account for the fact that with some probability, rejection may occur in which case no entropy is provided. Arguably, the source could be sampled repeatedly until it is not rejected. This assumes that independent identically distributed samples could be obtained. The rate at which entropy would depend on the rate of rejection. Filtered entropy does not depend on the rejection rate.

Remark 3.41. A possible alternative to contingent entropy is is to define the averaged filtered entropy. Using (3.26), gives

$$\left(\sum_{z\in Z} R_p(z)\left(\min_{x:f(x)=z} -\log_2 P_p(x)\right)\right) - H_{1,(\Pi,Z,R)}(p) \tag{3.28}$$

where (Π, Z, R) is the probability model defined by $R_p(z) = \sum_{x:f(x)=z} P_p(x)$, and $H_{1,(\Pi,Z,R)}(p)$ is the Shannon entropy of p in this model.

Remark 3.42. As noted previously, Shannon entropy is generally not the right type of entropy in cryptographic applications, because it is sometimes too optimistic by overestimating the entropy needed for cryptographic applications.

In (3.28), Shannon entropy appears as a negated term, so its being too optimistic translates into too pessimistic.

Remark 3.43. The following example shows that the average filtered entropy is not very appropriate for cryptography. Suppose that p is a uniform distribution on $X = \{0,1\}^N$, and the function f determining what adversary learns $f(x)$ about x is the function

$$f(x) = \begin{cases} x & \text{if } x_0 = 0\\ 1 & \text{if } x_0 = 1\end{cases} \tag{3.29}$$

If x drawn from p is to be used to be used a key, then over $1/2$ the time the adversary could guess x. Yet the average contingent entropy is about $\frac{N}{2}$.

3.3 Sample-Dependent Entropy Parameters

Sample-dependent parameters are distinct from the previous parameters in that they vary with the sample value. For a probability model (Π, X, P), a sample-dependent parameter is a function of the form

$$r : \Pi \times X \to R. \tag{3.30}$$

Sample-dependent parameters can be useful in cryptography when one is assessing the entropy of a value x which is to be used. In other words, in retrospective entropy assessment, sample-dependent parameters may be important.

Remark 3.44. Each of the grading function from §4 could also be considered of sample-dependent parameters, but these are not necessarily related to entropy.

3.3.1 Sample Entropy (Information Content or Self-Information)

The most obvious sample-dependent parameter is probability function P. It has parameter space $R = [0, 1]$. Although the probability function does not fit well to other notions of entropy, merely applying a logarithm, so $I = -\log_2 \circ P$, meaning

$$I(p, x) = -\log_2(P_p(x)) \tag{3.31}$$

with the convention that $-\log_2(0) = \infty$, yields a sample-dependent parameter with the same units as other notions of entropy.

Previous names for this sample-dependent parameter are *information content* and *self-information*. This report shall use an alternative name *sample entropy*, emphasizing that it is a type of entropy dependent on the sample.

Remark 3.45. The min-entropy of distribution p is the minimum of the sample entropy $I(p, x)$ over all $x \in X$. Explicitly:

$$H_\infty(p) = \min_{x \in X} I(p, x) \tag{3.32}$$

In particular, sample entropy is always at least min-entropy: $I(p, x) \geqslant H_\infty(p)$.

Remark 3.46. Shannon entropy $H_1(p)$ of distribution of p is the expected value of $I(p, x)$ for random $x \in X$ distributed according to distribution p. Explicitly:

$$H_1(p) = \sum_{x \in X} P_p(x) I(p, x). \tag{3.33}$$

Remark 3.47. Sample entropy is additive over independent distributions in the following sense. Suppose that (Π, X, P) and (Π, Y, Q) are models and the $(\Pi, X \times Y, R)$ is their common product. Then $I(p, (x, y)) = I(p, x) + I(p, y)$. (In a mixed product, $I((p, q), (x, y)) = I(p, x) + I(q, y)$.)

In prospective entropy assessment, sample entropy cannot be used, because one does not know the sample x, so one cannot compute the sample-entropy. Therefore, use of sample entropy is only applicable in retrospective assessment. In some cryptographic applications, entropy may be so scarce that one may wish to rely on sample entropy rather than just min-entropy.

Remark 3.48. The sample entropy of a bit string can be greater than its length, whereas this is not true for min-entropy or Shannon entropy. More generally, sample-entropy can exceed the min-entropy of the uniform distribution.

Remark 3.49. The fact that the sample-entropy can exceed the min-entropy of the uniform distribution, makes sample-entropy inappropriate for cryptographic applications, because an adversary, can always guess the value of x using a uniform distribution. Although when the adversary knows the distribution and the distribution is non-uniform, a uniform guess at x is the adversary's optimal strategy, a cryptographer should hesitate to assume that the adversary will adopt a particular strategy.

Remark 3.50. In certain probability models, if one is inferring the distribution p based from the sample x, it will be generally be the case that the inferred p favors x over other sample values and therefore that the sample-entropy entropy at the inferred distribution will not exceed the the min-entropy of the uniform distribution on the sample space. Thus, the difficulty described in Remark 3.49 will not arise.

Remark 3.51. A general way to view the adversary in the contexts above is as follows. We assume that the adversary knows the distribution p of the sample x. The adversary can adopt some probabilistic strategy to guess x, which will described by another distribution. The adversary's expected success rate is $p \cdot q$, thinking of p and q as probability vectors of dimension $|X|$. Write q_x (resp. p_x) for the probability that distribution q (resp. p) assigns to x. Write $q = d^x$ for the distribution q such that $q_x = 1$ and $q_y = 0$ for $y \neq x$.

Let z maximize p_z. The min-entropy of p is thus $-\log_2 p_z$. Four strategies for the adversary and their expected success rates are given below.

1. Given p, the optimal strategy is $q = d^z$, then the adversary's expected success rate is $p \cdot d^z = p_z$. The negative base-two logarithm of the success rate with this strategy is the min-entropy of p.
2. If the adversary chooses $q = d^x$, for some other x, then the expected success rate is $p \cdot d^x = p_x$. In particular, the negative base two logarithm of the expected success rate of strategy $q = d^x$ is the sample entropy of x under the distribution p.
3. If the adversary chooses q as the uniform distribution on X, then $p \cdot q = 1/|X|$. The negative base two logarithm of the expected success rate is then $\log_2 |X|$.
4. If the adversary choose $q = p$, then the negative base two logarithm of the expected success rate is the Renyi entropy of order two of the distribution p.

In the event that the sample actually takes the value x, then adversary's actual success rate is different from the expected success rate. In particular, the strategy $q = d^x$ has success rate one. But the adversary has no information with which to help choose this particular strategy.

Remark 3.52. Another potential pitfall of using sample entropy is that that in distributions p where P_p takes distinct values for every x, the sample entropy determines x. In this case, if the sample entropy is somehow leaked to an adversary, then the adversary learns x. To avoid a contingent entropy of zero, the sample entropy must not be leaked.

3.3.2 Eventuated Min-Entropy

Let (Π, X, P) be a probability model. Let $E \subseteq X$. Suppose that the event $x \in E$ has occurred. Then the *eventuated min-entropy* associated with E is a sample-dependent parameter given by

$$H_{\infty\|E}(p, x) = -\log_2 \max_{x' \in E} P_p(x'). \tag{3.34}$$

for $x \in E$. For $x \notin E$, this parameter's value does not matter, so can artificially be set to ∞.

Remark 3.53. Eventuated min-entropy is the sum of filtered min-entropy and applied sample entropy.

More explicitly, given f and $y = f(x)$, the eventuated min-entropy is the sum of the $Y = f^{-1}(y)$ filtered min-entropy and the sample entropy of the $f(x)$ in the f-applied probability model.

Remark 3.54. Eventuated min-entropy, like min-entropy and sample entropy, is additive over independent distributions.

3.3.3 Applied Eventuated Min-Entropy

Let (Π, X, P) be a probability model. Let $E \subseteq X$. Suppose that the event $x \in E$ has occurred. Let $f : X \to Y$ be a function, such that only $f(x)$ will be used, say, as part of secret key. Then the *applied eventuated min-entropy* of $f(x)$ associated with E and f is a sample-dependent parameter given by

$$H_{f(\infty)\|E}(p, x) = -\log_2 \max_{y \in f(E)} \sum_{x' \in f^{-1}(y)} P_p(x') \tag{3.35}$$

for $x \in E$. For $x \notin E$, this parameter's value does not matter, so can artificially be set to ∞.

Applied eventuated min-entropy is the eventuated min-entropy of the event $y \in f(E)$ in the f-applied model.

3.3.4 Contingent Eventuated Min-Entropy

Let (Π, X, P) be a probability model. Let $E \subseteq X$. Suppose that the event $x \in E$ has occurred. Let $g : X \to Z$ be a function, such that $g(x)$ will be learnt by an adversary. Then the *contingent eventuated min-entropy* of x associated with E under leakage of $g(x)$ is a sample-dependent parameter given, for $x \in E$, by

$$H_{\infty|g\|E}(p, x) = \min_{f:X\to Y} H_{f(\infty)\|E}(p, x) \tag{3.36}$$

where f ranges over all function supplementary to g. For $x \notin E$, this parameter's value does not matter, so can artificially be set to ∞.

4 Statistical Inference

The general problem that we need to consider is the *(statistical) inference* problem, which is the following. The knowns are (Π, X, P) and $x \in X$, and the unknown is p. We wish to infer something about which p could have given rise to x. In cryptology, we especially wish to infer something about the various parameters (especially entropy) of the distribution p.

Remark 4.1. Inference necessarily involves guessing, because x generally gives incomplete information about p. There may be many different distributions $p \in \Pi$ consistent with x in the sense that that $P_p(x) \neq 0$, and it is not strictly possible to know which of these p is the correct. So inference is imperfect.

Remark 4.2. Because of the intrinsic imperfection and incompleteness of inference, we can only hope to obtain an inference method that is reasonable. Reasonable is perhaps definable, but we will not formally define it in this report.

Remark 4.3. In this report, the view is taken that, to be reasonable, an inference method must at least be well-defined on all, or a very large class, of probability models. A reasonable inference method should also be invariant under isomorphism of models. Furthermore, a reasonable inference method should not just be well-defined, but should also agree with intuition on very restrictive models.

Remark 4.4. As stated earlier, using an inference method that is general in the sense above avoids crafting an inference method too specific to a particular source and probability model. Although, perhaps there are reasonable inferences methods specific to a given probability model, the view taken in this report is that all assumptions about the source should be confined to, or concentrated in, the probability model. An inference method specific to a probability model or source contradicts this view, and has the disadvantage of dispersing the assumptions made about the source in distinct areas.

4.1 Inference functions

An *inference function* for (Π, X, P) is a function that takes input of X and outputs some *assertion* about the unknown distribution p. In this report, we first consider three types of direct assertions about p. Later, we will discuss how such inferences about p may be converted into inferences about the cryptologically important parameters of p.

4.1.1 Point-valued inferences

An inference function i for model (Π, X, P) is *point-valued* if it is a function of the form $i : X \to \Pi$. That is, to each sample value x it assigns a probability distribution, which can be called the *inferred* distribution.

4.1.2 Set-valued inferences

Let (Π, X, P) be a probability model. Let $[\Pi]$ be the set of all subsets of Π. An inference function for model (Π, X, P) is *set-valued* if it is a function of the form $i : X \to [\Pi]$. That is, to each sample value x it assigns a set of distributions, which we will call the *inferred set* of distributions, or set of *inferred* distributions.

Remark 4.5. Generally, we expect set-valued inference functions to respect equivalence of distributions: if $p \in i(x)$ and $p \equiv q$, then $q \in i(x)$. Otherwise, the set-valued inference functions should be deemed unreasonable.

Remark 4.6. In this report, we will consider several specific set-valued inference functions.

Remark 4.7. Often, the inferred sets contain just a single distribution. In such cases, we may think of the set-valued inference as being point-valued. Sometimes a sample value $x \in X$ has an inferred set containing many distributions, even infinitely many. For such a sample, a set-valued inference function is unable to prefer one distribution over the other in the inferred set. For cryptographic applications, the most cautious choice of distribution can be inferred, as described in §4.5.2.

Remark 4.8. The inferred set is sometimes empty.

4.1.3 Grading-valued inferences

A *grading* on the model (Π, X, P) is a function $g : \Pi \to [0, \infty)$, meaning a non-negative real-valued function on the probability space. The set of gradings can be written $\Gamma(\Pi)$.

A *strict grading* is a function $g : \Pi \to [0, 1]$; so a strict grading taking value at most one. Usually, the gradings considered in this paper will be strict gradings. The set of strict gradings can be written $[0, 1]^{\Pi}$.

A *binary grading* is function $g : \Pi \to \{0, 1\}$; so a binary grading is a strict grading taking only integer values. The set of binary gradings can be written $\{0, 1\}^{\Pi}$. A binary grading can be regarded as equivalent to subset of Π by the relation $g \equiv g^{-1}(1)$.

A *grading-valued* inference function for model (Π, X, P) is a function of the form $i : X \to \Gamma(\Pi)$; so i assign each sample value x a *grading* $g = i(x)$ of all probability distributions in the probability space Π. The intention of a grading-valued inference function is that a higher value of the inferred grading is intended to indicate better consistency of the sample value x the distribution p.

Remark 4.9. Generally, we expect grading-valued inference functions to respect equivalence of distributions: if $g = i(x)$ is an inferred grading function and $p \equiv q$, then $g(p) = g(q)$.

If $i(x) = g$, then $g(p)$ is called the *grade* of p at x. In strict notation the grade is $i(x)(p)$, but to avoid the double argument, the notations $i(x, p)$ or even $i_x(p)$ may be used when clear from context. So, a grading-valued inference function may be thought of as a bivariate function $i : X \times \Pi \to [0, \infty)$. With a slight re-use of terminology, we may also call such a bivariate function a *general grading* function. So, a grading-valued inference function can be thought of we can define a general grading function, and vice versa.

Remark 4.10. We can also fix a probability distribution p and a grading-value inference function i, we get a function $i_p : X \to [0, 1] : x \mapsto i(x, p)$. So, if $i(x) = g$, then $i_p(x) = g(p)$.

Remark 4.11. Unlike probability functions, there is no requirement that the grading function $g : \Pi \to [0, 1]$ sums to one over Π, in the sense that $\sum_{p \in \Pi} g(p) = 1$ if, say, Π is finite. Indeed, in general, for infinite probability spaces (for which no measure has been assigned), such summation to one is not even a well-formulated requirement. If Π is equipped with a measure, it may be convenient for g to be a measurable function, ideally whose total integral over Π is finite.

Remark 4.12. A set-valued inference functions may be encoded as grading-valued inference functions as follows. The set $[\Pi]$ of subsets of Π may be encoded as the set $\{0, 1\}^{\Pi}$ of functions $g : \Pi \to \{0, 1\}$ under that convention that a subset of Π is encoded as function taking value 1 on the subset and value 0 off the subset. The set $\{0, 1\}^{\Pi}$ is a subset of the set $[0, 1]^{\Pi}$, so therefore this encoding is also an encoding of $[\Pi]$ inside of $[0, 1]^{\Pi}$. Applying this encoding to the outputs of set-valued inference function, we get a grading-valued inference function.

Remark 4.13. The output of a grading-valued inference function for a probability model with infinite probability space describes an infinite amount of information (but will typically be described with a finite formula, of course). For the purposes of cryptography, what is needed is a single estimate of entropy, a single real number, or perhaps even just a single bit: a decision to accept or reject. As such, grading-valued inference functions seem not to be directly usable for the cryptographic applications. Nevertheless, we may derive from grading-valued inference functions other inferences, such as set-valued inferences, so, it will turn out that grading-valued inference functions can serve as intermediate steps in cryptographic applications.

4.1.3.1 Expectation of General Gradings

It can be informative to consider the expected value of a general grading function $g : X \times \Pi \to [0, \infty)$, at a given probability distribution. Precisely, this is defined as

$$E_g(p) = \sum_x g(x, p) P_p(x) \tag{4.1}$$

This value $E_g(p)$ gives a way to calibrate a grading value $g(x, p)$, especially one p has be inferred from x using the grading g. If $g(x, p)$ is many magnitudes lower than $E_g(p)$, then one may even infer that the probability model itself is not accurate.

Remark 4.14. For the likelihood grading, §4.4.1, $E_{g_L}(p) = \sum_x P_p(x)^2$. For uniform distribution, and more generally, distributions with high min-entropy, this value can be quite low. In fact, taking the negative logarithm of the expected likelihood

grading, we get $-\log_2(E_{g_L}(p)) = H_2(p)$, the Renyi entropy of order two. Renyi entropy of order two is known to be at most twice min-entropy, so it can be quite small.

Remark 4.15. For typicality grading, §4.4.2,

$$\begin{aligned}
E_{g_k}(p) &= \sum_x g_k(x,p)P_p(x) \\
&= \sum_{P_p(y)<P_p(x)} P_p(y)P_p(x) + k \sum_{P_p(y)=P_p(x)} P_p(y)P_p(x) \\
&= \frac{1}{2}\left(\sum_{P_p(y)<P_p(x)} P_p(y)P_p(x) + \sum_{P_p(y)>P_p(x)} P_p(y)P_p(x) + 2k \sum_{P_p(y)=P_p(x)} P_p(y)P_p(x) \right) \\
&= \frac{1}{2}\left(\sum_{x,y} P_p(y)P_p(x) + (2k-1) \sum_{P_p(y)=P_p(x)} P_p(y)P_p(x) \right) \\
&= \frac{1}{2}\left(\left(\sum_x P_p(x) \right)^2 + (2k-1) \sum_{P_p(y)=P_p(x)} P_p(x)^2 \right) \\
&= \tfrac{1}{2} + (k-\tfrac{1}{2}) \sum_x P_p(x)^2 Q_p(x)
\end{aligned} \tag{4.2}$$

where $Q_p(x)$ is the number of y such that $P_p(x) = P_p(x)$.

Therefore, the expected value of the balanced typicality is exactly one half. Inclusive typicality averages to more than one half, and exclusive to less. If P_p takes distinct values for all x, then the expected inclusive and exclusive typicality differ from one half by the expected value of the likelihood grading. If p is a uniform distribution, then the expected value of inclusive typicality is one: indeed inclusive typicality is one: all values of x are equally typical. Conversely, the expected value of exclusive typicality is zero for a uniform distribution p.

Therefore, inclusive and balanced typicality are inherently calibrated in the sense that no matter what p is, the expected typicality is at least one half. Furthermore, if x is such that $g_1(x) \ll \frac{1}{2}$ or $g_{1/2}(x) \ll \frac{1}{2}$ for probability distributions p in the probability model, then one can infer from x that perhaps the probability model is not valid.

Remark 4.16. The expected value of all generalized typicalities, §4.4.3 are $\frac{1}{2}$.

Remark 4.17. The expected value of the Bayesian grading §4.4.6 is

$$E_B(p) = \frac{1}{\int_\Pi L_x d\mu} \sum_{x \in X} P_p(x)^2 \tag{4.3}$$

and is a scaled version of the expected value of the likelihood grading.

4.2 Inference Methods

An *inference method* is a function that takes a probability model (Π, X, P) and outputs an inference function for (Π, X, P). We say an inference method is point-valued if all inference functions it produces are point-valued. Similarly, an inference method is set-valued if it only outputs set-valued inference functions, and grading-valued if it only output grading-valued inference functions.

Remark 4.18. A grading-valued inference method I is a function to functions to functions: on input of a probability model (Π, X, P), inference method I outputs a grading-valued inference function i, which, in turn, is a function with domain X and range of functions from Π to $[0, 1]$.

A grading-valued inference function for the probability model (Π, X, P) has an associated general grading function $g : X \times \Pi \to [0, 1]$. Any function that maps a model to such general grading function on the model, is called a general grading method. Each grading-valued inference method defines a general grading method, and each general grading method defines a grading-valued inference method.

4.3 Set-Valued Inference From Grading-Valued Inferences

In this section, we give two ways to derive a set-valued inference method from a grading-valued inference method. Collectively, we will call such methods *graded inference methods*.

4.3.1 Maximally Graded

Suppose that g is a general grading on model (Π, X, P), associated with a grading-valued inference function i_g. The *maximally graded inference* associated with grading g is a set-valued inference $i_{\max g}$ function generally defined as follows:

$$i_{\max g}(x) = \{p : g(x,q) \leqslant g(x,p) \forall q \in \Pi\} \tag{4.4}$$

We may think of $i_{\max g}$ as derived from g or from i_g.

Remark 4.19. In some cases g is discontinuous and such a maximum p may not exist. In these case, an alternative definition may sometimes be available. Consider the supremum of gradings values at x, written $s_x = \sup_{p\in\Pi} g(x,p)$. Define sets $S_\epsilon = \{p : g(x,p) \geqslant s_x - \epsilon\} \subseteq \Pi$, which are nested according to the size ϵ.

As a matter of convenience, we may define the following. Let $\bar{S}_\epsilon$ be the closure of S_ϵ in some natural topology on Π. If $i_{\sup g}(x) = \bigcap_{\epsilon>0} \bar{S}_\epsilon$ is non-empty (which is true if Π is given a compact topology), may serve as a suitable substitute for an empty set $i_{\max g}(x)$, even if values of $g(x,p) < s_x$ for $p \in i_{\sup g}(x)$.

In cryptographic applications, we are more interested in entropy parameters than in distributions themselves. If the parameters are continuous then the definition of $i_{\sup g}(x)$ above will provide the desired answer for the parameters. For discontinuous parameters the parameters on the $i_{\sup g}(x)$ may not be what is desired. In this case, we should think of $i_{\sup g}(x)$ not as the intersection of the chain of sets of $\bar{S}_\epsilon$, but rather as the limit of the chain of sets S_ϵ. This enables us to consider limits of parameters on S_ϵ, which may differ the value of parameters on the intersection.

Remark 4.20. In many cases, the inferred set $i_{\max g}(x)$ is a single element (singleton) set. In these case, the inference is much like a point-valued inference function. However, there are often some values of x for which several, possibly infinitely many, different distributions p attain the maximal value.

If G is a general grading method or I_G is grading-valued inference method, then it is possible to derive a set-valued inference method $I_{\max G}$ using the inference functions above.

Remark 4.21. Maximally graded inferences are model-dependent in the sense that (4.4) include Π. A potential consequence of this model-dependence is that the maximally graded inference in restriction (Θ, X, P) of the model (Π, X, P), may not have a given relation with the maximally graded inference in the model (Π, X, P).

4.3.2 Threshold Graded and Confidence Levels

Suppose that g is a general grading on a model (Π, X, P). Let $t \in [0,1]$ and call this value the *threshold level*. The *threshold graded inference* function $i_{g>t}$ is a set-valued inference function defined by

$$i_{g>t}(x) = \{p : g(x,p) > t\} \tag{4.5}$$

If $t > u$, then $i_{g>t}(x) \subseteq i_{g>u}(x)$, so the sets obtained are shrinking or stable in size as a function of the threshold. A high threshold may lead to a narrow, perhaps even empty, inference, while a low threshold may lead to a broad inference.

The value $c = 1 - t$ may sometimes be called the *confidence level* of the inference. As confidence increases, the breadth of the inference may increase (or stay stable). This reflects the intuitive notion that as we can generally make the sacrifice of broadening our inference set in order to gain a more confident inference.

Gradings are best thresholded when the distribution of the grading, for fixed p and varying x, has some resemblance to the uniform distribution on $[0,1]$, because the confidence level has a better meaning. In the following sections, some gradings will have such a property, and others will not.

Remark 4.22. Threshold graded inferences are not model-dependent in the sense of Remark 4.21 provided that the grading is not model-dependent. In particular, if the set $i_\Theta(x)$ threshold graded inference in restriction (Θ, X, P) of the model (Π, X, P),

and $i_\Pi(x)$ is the threshold graded inference in the model (Π, X, P), then

$$i_\Theta(x) = \Theta \cap i_\Theta(x). \tag{4.6}$$

When using such a threshold graded inference and taking the infima of parameters, as in §4.5.2, then restriction of the of model cannot decrease the inferred parameter, and relaxing model cannot increase the inferred parameter.

Remark 4.23. As note in Remark 2.5, we may sometimes presume that an adversary can influence the choice of p in Π. If an adversary has such power over p, then a maximally graded inference has little value. For appropriate gradings, a high-confidence threshold grading would still have some value.

4.4 Example Gradings

In this section, we describe some important grading methods.

4.4.1 Likelihood

The likelihood grading g_L is defined by

$$g_L(x, p) = P_p(x). \tag{4.7}$$

We also write for the associated inference function $L_x = i_{g_L}(x)$. Therefore $L_x(p) = P_p(x)$.

The term *likelihood*, instead of *probability*, is used here to avoid thinking that L_x has the properties of probability function. For example summing (or integrating) the values of L_x over all probability distributions is not guaranteed to yield 1.

Remark 4.24. For likelihood, we make an exception about the general grading's name, since the general grading is simply the probability function, we will call the grading the likelihood grading.

Remark 4.25. Likelihood is, of course, a well-known and fundamental notion in statistical inference.

4.4.2 Typicality

For a given *inclusivity level* $k \in [0, 1]$, we define the *typicality grading* g_k as follows:

$$g_k(x, p) = \left(\sum_{y : P_p(y) < P_p(x)} P_p(y) \right) + k \left(\sum_{y : P_p(y) = P_p(x)} P_p(y) \right) \tag{4.8}$$

In this report, we will only consider $k \in \{0, \frac{1}{2}, 1\}$, which give rise to *exclusive*, *balanced* and *inclusive* typicality.

Remark 4.26. Inclusive typicality $g_1(x, p)$ is the probability that a random sample y is at most as probably as x. Exclusive typicality $g_0(x, p)$ can also be defined as the probability that a random sample y is less probable than x. Balanced typicality is the average of inclusive and exclusive typicality, in other words, it is half-way between inclusive and exclusive.

Remark 4.27. Typicality, unlike likelihood, when used for inference, attempts to captures the notion of how a sample compares in probability to other samples under the same probability distribution. This notion was used in the intuitive reasoning for the loose inference in §1. We argued that 2^{10}-uniform was not to be inferred, because more repetitions would have been expected in the sample.

Remark 4.28. For a fixed distribution p, ranking sample values x by typicality or likelihood gives the same ranking. For fixed x, and varying p, the rankings induced by typicality may differ from those by likelihood.

Remark 4.29. When p is a uniform distribution on X, then typicality is constant for all x, and takes on the value k, the inclusivity. When p is an almost uniform distribution on X, then for the most probable x, it takes value approximately $1 - (1 - k)/|X|$. For $k < 1$, this will be larger than the typicality at the uniform distribution.

Remark 4.30. When p is subuniform on X, then

$$g_k(x,p) = \begin{cases} k & \text{if } P_p(x) > 0 \\ 0 & \text{if } P_p(x) = 0 \end{cases} \tag{4.9}$$

As such, in models that admit subuniform distributions, any inference based on typicality treats them equally. This effect may be called subuniform distribution have tied typicality. We call this the *tying effect*.

Some models may admit distributions with higher typicality than all subuniform distributions, in which case some useful inferences can be made.

In some case, sample statistics (§5) may serve as *tiebreakers* between subuniform distributions.

Remark 4.31. Inclusive typicality is always at least as large as likelihood:

$$g_1(x,p) \geqslant g_L(x,p) \tag{4.10}$$

but balanced and inclusive typicality could be less. Similarly, $1 - g_0(x,p) \geqslant g_L(x,p)$. A stronger fact is that the gap between exclusive and inclusive typicality is always at least the likelihood.

$$g_1(x,p) - g_0(x,p) \geqslant g_L(x,p) \tag{4.11}$$

Remark 4.32. The notion of *typicality* is based on well-known notions in statistics of *significance level*, *p-value* (also known as *percentile* or *quantile*, depending on the units) and *cumulative probability function*.

The general notion of *significance level* refers to a value of the *p-value*. The general notion of *p-value* is sample statistic (see §5) that takes uniform distribution on $[0,1]$, at least under the *null hypothesis*.

A p-value statistic may be formed for continuous distributions by taking cumulative probability function with respect to some function f defined on the sample space. Any choice function f yields a p-value. So, the p-value of x is probability that $f(y) \leqslant f(x)$, for y drawn from the same distribution.

A common use of p-values is when the distribution is a normal distribution and the function f is the identity, then p-value is related to the Gauss error function.

In cryptographic applications, distribution are typically discrete, so achieving a proper p-value uniformly distributed in $[0,1]$ cannot be strictly. Nevertheless one can try to approach such a distribution.

In cryptographic applications, the parameters of interest, entropy and so on, depend primarily on the distribution, not on specific properties structure in x.

Typicality is the cumulative probability with respect to the probability function.

Remark 4.33. Typicality ranks the sample space by probability. Such ranking is often implicit in rankings of popularity, such as music sales charts, election results, and lists of popular names. The game show *Family Feud* involves such a ranking.

Remark 4.34. For an almost uniform distribution (with no equal probabilities), the distribution typicality is almost uniformly distributed in $[0,1]$, much like a p-value.

Remark 4.35. Randomized typicality $g_?(x,p)$ is a random variable defined with the same equation (4.8) as typicality, except that variable k is selected uniformly at random from the interval $[0,1]$. If distribution p is fixed, and x has the probability distribution given by p, then $g_?(x,p)$ is uniformly distributed in $[0,1]$.

Remark 4.36. The notion of *typicality* adapts the notion of significance level towards the task of assessing min-entropy, in that it ranks the samples according to their probabilities.

4.4.3 Generalized Typicality and Adjusted Likelihood

Let $\sigma : [-1,1] \to [-1,1]$ be an odd, non-decreasing function.

Remark 4.37. Odd means that $\sigma(-x) = -\sigma(x)$ and non-decreasing means that if $x < y$ then $\sigma(x) \leqslant \sigma(y)$.

Remark 4.38. We can write $\sigma(x) = xf(x^2)$, where $f : [0,1] \to [0,\infty)$ is a function such that for $x > 0$, it holds that $f(x) < 1/\sqrt{x}$.

Generalized typicality is parametrized by such functions σ. So, σ-typicality is defined by

$$t_\sigma(x,p) = \tfrac{1}{2}\left(1 + \sum_y P_p(y)\sigma\left(P_p(x) - P_p(y)\right)\right) \tag{4.12}$$

This is a generalization of balanced typicality in the following senses.

1. Putting $\sigma(x) = x/|x|$ for $x \neq 0$ and $\sigma(0) = 0$, which is to say, putting $\sigma(x)$ equal to the sign of x, then σ-typicality is balanced typicality.
2. For any σ meeting the stated conditions, σ-typicality is bounded, like balanced typicality, such that $t_\sigma(x,p) \in [0,1]$ for all x and p. (This is due to the values of σ belonging to $[-1,1]$.)
3. The expected value (see §4.1.3.1) of σ-typicality is $1/2$, just like balanced typicality. (This is due to σ being an odd function.)

Remark 4.39. When p is subuniform on X, then

$$t_\sigma(x,p) = \begin{cases} \frac{1}{2} & \text{if } P_p(x) > 0 \\ \frac{1}{2} - \frac{\sigma(P_p(y))}{2} & \text{if } P_p(x) = 0 \text{ and } P_p(y) > 0 \end{cases} \tag{4.13}$$

As such, in models that admit subuniform distributions, any inference based on generalized typicality treats all subuniform distributions equally. In such cases, sample statistics (§5) may serve as tiebreakers.

Remark 4.40. For threshold-graded inference at confidence levels of $\frac{1}{2}$ or higher, all subuniform distributions in the model must be considered, because of the upper case in (4.13).

Remark 4.41. For threshold-graded inference at confidence levels strictly higher than $\frac{1}{2}$, given sample x, some subuniform distributions p in which $P_p(x) = 0$ may be inferred, because of the lower case of (4.13). Call this an *aberrant* inference. Aberrant inferences can be viewed as a strong discrepancy with the likelihood grading.

Balanced typicality does not exhibit aberrant inferences because the σ is the sign function, which causes the lower term to be zero in the lower case of (4.13).

Despite aberrant inferences, generalized typicality can still be useful for assessing entropy. For example, at a given confidence level, only those subuniform, among those allowed within the probability model, with a given support of a given size but not containing the given sample x will be inferred.

If the infimum entropy is always attained at an aberrantly inferred subuniform distribution, then at least the inferred entropy decreases with confidence level, decreasing to 0 as the confidence level approaches 1.

Remark 4.42. Generalized typicality sheds some light on a potential difficulty of balanced typicality: its discontinuity. The discontinuity is made clearly attributable to the discontinuity of the sign function σ when viewed of as a case of generalized typicality.

Balanced typicality has the disadvantage of being discontinuous, which can make bounding and optimizing balanced typicality difficult. This disadvantage can perhaps be overcome by using the observation that balanced typicality is σ-typicality with σ equal to the sign function. The idea is then to use an alterntiva σ function, which is continuous, or even smooth, but still similar enough the sign function to so that the resulting σ-typicality inherits the desired properties of balanced typicality.

The simplest continuous form of generalized typicality is given the choice $\sigma_1(x) = x$. In this case, formula (4.12) simplifies to:

$$t_{\sigma_1}(x,p) = \tfrac{1}{2}\left(1 + P_p(x) - \sum_y P_p(y)^2\right). \tag{4.14}$$

As a function of x, with p fixed, the varying term in the sum, $\frac{1}{2}P_p(x)$, is a scaling of the likelihood grading. When p varies, the term $\sum_y P_p(y)^2$ varies, but independently of x. We call this special case of generalized typicality, *adjusted likelihood*.

Remark 4.43. Adjusted likelihood can be related to order two Renyi entropy by $t_{\sigma_1}(x,p) = \frac{1}{2}\left(1 + L_x(p) - 2^{-H_2(p)}\right)$.

Remark 4.44. Adjusted likelihood, and other generalized typicalities, can, for certain distributions p, even distributions moderately distant from uniform distributions, be very close to $\frac{1}{2}$, for two reasons. First, both the likelihood and adjustmust terms can be quite small even for quite non-uniform distribution. Second, the difference between the likeilhood and the adjustment term can very small. Calculations done with such typicalities may require considerably high degrees of precision.

Adjusted likelihood may also be expressed as

$$t_{\sigma_1}(x,p) = \frac{1}{2}\left(\frac{5}{4} - \left(P_p(x) - \frac{1}{2}\right)^2 - \sum_{y\neq x} P_p(y)^2\right) \tag{4.15}$$

which is an affine transformation of the square of the Euclidean distance (as would be natural to define in the unrestricted model) between the distribution p and the pseudo-distribution h_x that has probability 1/2 of taking value x, and probability 0 otherwise. So, $t_{\sigma_1}(x,p) = \frac{5}{8} - \frac{1}{2}\|h_x - p\|^2$.

Two other families of σ, generalizing σ_1, seem reasonable to consider.

1. Let $\sigma_{1/m}(x) = \sqrt[m]{x}$ for m odd. At $m = 1$, the function is $\sigma_1(x) = x$, which gives the adjusted likelihood that we have already seen. Taking $m \to \infty$, these functions approach the sign function. These functions $\sigma_{1/m}$ are continuous, but have infinite slope at 0, which may make some optimization algorithms difficult.

2. Let
$$\sigma_m(x) = x\sum_{n=0}^{m-1}\binom{-1/2}{n}(x^2-1)^n, \tag{4.16}$$
for integers $m \geqslant 1$. At $m = 1$, the function is $\sigma_1(x) = x$ (and is the same as $\sigma_1(x)$ above). As $m \to \infty$, the functions approach the sign function.

Remark 4.45. The series (4.16) comes the from the Taylor series expansion of $1/\sqrt{x}$ at $x = 1$, and using polynomial prefixes of these series as a function $f(x)$ and defining $\sigma = xf(x^2)$.

Remark 4.46. The function $\sigma_2(x) = \frac{x}{2}(3 - x^2)$. The function $\sigma_3(x) = \frac{x}{8}(15 - 10x^2 + 3x^4)$. Also, $\sigma'_m(0) = O(\sqrt{m})$ seems to hold, with $2\sqrt{m/\pi}$ seeming to be a good approximation.

In generalized typicality, summation over all y results in some symmetry. For example, taking the polynomials σ_m above, we can write:

$$t_{\sigma_m}(x,p) = \sum_{n=0}^{m-1} Q_n(p)P_p(x)^n \tag{4.17}$$

where $Q_n(p)$ is a symmetric polynomial evaluated at all of the variables $P_p(y)$ (for each possible value of y). Expressing $Q_n(p)$ as a polynomial of symmetric polynomials, such as elementary symmetric polynomials, may give a sum with fewer terms than there are values of y, and as such, may make calculating with and optimizing σ-typicality easier.

Remark 4.47. In the independent model, $Q_n(p)$ can also be expressed as a related symmetric function of the components of the p vector.

4.4.4 Calibrated Typicality

Generalized typicality t_σ may have a tendency to be too close $\frac{1}{2}$, and as such be not too useful for establishing confidence levels. For a grading to be most meaningful in the sense of confidence levels, the distribution of the grading should be somehow nearly uniform on the interval $[0,1]$. More precisely, for each fixed p, the distribution $g(x,p)$ should be almost uniform.

Balanced typicality, and generalized typicality, approach such uniformity mainly in the sense that g has expected value $\frac{1}{2}$ and g is value within $[0,1]$. Another measure of closeness to uniformity could be to look at the variance. The

variance of the uniform variable in $[0,1]$ is $\frac{1}{12}$. So given a grading g with expectation $\frac{1}{2}$ and variance v, one could apply a linear transformation to get another grading $g' = \frac{1}{2} + \frac{1}{\sqrt{12v}}(g - \frac{1}{2})$, with expectation $\frac{1}{2}$ and variance $\frac{1}{12}$. A potential problem with g' is that it might have value outside the range $[0,1]$, in which confidence levels make little sense (a negative confidence level or confidence level higher than 1 makes little sense).

An alternative calibration method is as follows. Let $\kappa : [0,1] \to [0,1]$ be another odd function. Then κ-calibrated σ-typicality is defined to be:

$$t_{\sigma,\kappa}(x,p) = \tfrac{1}{2}\left(1 + \kappa\left(\sum_y P_p(y)\sigma(P_p(x) - P_p(y))\right)\right) \tag{4.18}$$

Balanced typicality is the special case of calibrate typicality in which κ is the identity function and σ is the sign function.

4.4.5 Agreeability Gradings

The notion of statistical distance from (2.29), which has sometimes been used in cryptography, can be adapted to act like a grading. Let p_x be the deterministic distribution, in the unrestricted model, that takes on sample value x with probability one, and thereby all other values with probability zero. One could define the *agreeability* grading as

$$g_a(x,p) = 1 - d(p_x, p) \tag{4.19}$$

but such an agreeability grading simplifies to

$$g_a(x,p) = P_p(x) \tag{4.20}$$

which is just the likelihood grading.

One could use other distance metrics, such as those based on the Euclidean metric (L^2 norm), as defined on the natural parametrization of the unrestricted model, In fact, adjusted likelihood is already related to such a metric. One could also use a distance based on the L^∞ norm to define an agreeability rating.

4.4.6 Bayesian Grading and Posterior Probabilities

The Bayesian grading $B_x : \Pi \to [0,\infty]$ can be defined when the probability space Π is equipped with a measure μ, and the likelihood grading L_x is measurable and integrable with respect to this measure. It is defined as:

$$B_x = \frac{L_x}{\int_\Pi L_x d\mu}. \tag{4.21}$$

We also write $B_x(p) = B(x,p)$.

Remark 4.48. The Bayesian grading is based on *Bayes' law* for conditional probabilities. Elaborating our probability notation slightly (to a notation so commonly used in much previous work that formal definitions will be omitted in this report), Bayes' law states that the conditional probability is $P(A|B) = P(A \cap B)/P(B)$. This implies that $P(B|A) = P(A|B)P(B)/P(A)$. For the problem at hand, we want the conditional probability of the hypothesis p given the evidence x, so $P(p|x)$, which is given by the formula $P(x|p)P(p)/P(x)$. The factor $P(x|p) = P_p(x) = L_x(p)$, by definition. The factor $P(x)$ is the marginal probability of x over all possible distributions p, using the associated prior probabilities, which is the integral in the denominator of (4.21). The factor $P(p)$ is set to 1 in (4.21) because it is really covered by the measure μ itself. In other words, we when we integrate the function B_x over the measure, the measure provides the contribution of $P(p)$, that is, the prior probabilities.

Remark 4.49. Define $\nu = B_x\mu$ as another measure on Π with the definition $\nu(S) = \int_S B_x d\mu$ for any subset $S \subseteq \Pi$. In fact, this resulting measure satisfies $\nu(\Pi) = 1$, so actually ν can be used to define probabilities of the probabilities distribution. These are known as the *posterior probabilities*.

4.5 Parameter Inference

Cryptographers are primarily interested in inference about the entropy parameters.

4.5.1 Converting Distribution Inference from Parameter Inferences

If i is an inference function for a model (Π, X, P) and $r : \Pi \to R$ is a parameter on the model, then it is sometimes possible to define a *indirect inference function* for R, which is function from X to assertions about R. The definition depends on the nature of the inference function, as given below.

4.5.1.1 Point-valued For a point-valued inference function $i : X \to \Pi$, the the naturally parameter induced inference function for R is defined as $j : X \to R : x \to r(i(x))$.

4.5.1.2 Set-valued For a set-valued inference function i mapping X to subsets of Π, the naturally parameter induced inference function j for R maps X to subsets of R in the following manner. For $T \subseteq \Pi$, define $r(T) = \{r(p) : p \in T\}$ and define $j(x) = r(i(x))$.

4.5.1.3 Grading-valued For a grading-valued inference function i mapping X to functions from Π to $[0, 1]$, then there is no naturally induced inference function for R, unless further information is available. If the probability space is equipped with an appropriate measure, we can defined *Bayesian parameter induced inference function* $j : R \to [0, 1]$, as follows. For $y \in R$, let $j(y)$ be the average value of i over the the subset $r^{-1}(y)$ of the probability space Π. The function j is only defined where the sets $r^{-1}(y)$ are measurable and i is measurable on this set.

4.5.2 Narrowing Set-Valued Entropy Inferences to a Point-Valued by Infima

Often, a narrower inference is desired than provided by a set-valued naturally parameter induced inference. In this case, one may want to apply a further function to the parameter inference $j(x)$. For example, if the parameter space R is an ordered set, then one can take the minimal (or infimum) value of r on the set $j(x)$.

In cryptology, one wants to be prudent, so taking a minimum, or infimum, value of min-entropy over a set of reasonably inferred possible probability distributions is best.

On the other hand, one does not want to waste entropy when it is scarce, so one needs to strike a balance. To this end, one may want to isolate conceptually the stage where one might underestimate entropy, specifically to this stage. In particular, if the inferred set of entropies includes both large and small values, perhaps application of second inference method is appropriate.

Remark 4.50. In the case of a Bayesian parameter induced inference function j where the parameter space R is also equipped with a measure, then one can take the weighted average of parameter values in the parameter space R using the function j as the weighting.

In general averaging is risky in cryptography because adversaries are not restricted to average behavior, so this method is not recommended for estimating min-entropy in cryptology.

5 Sample Statistics

A *sample statistic*, or just *statistic*, on a probability model (Π, X, P) is a function $s : X \to S$. A *statistic method* is a function from probability models to statistics on these models.

5.1 Induced Model

A statistic s defines an *induced probability model* (Π, S, Q) as follows:

$$Q_p(t) = \sum_{s(x)=t} P_p(x) \tag{5.1}$$

Two sample statistics for a given probability model may be regarded as *equivalent* if their induced probability models are equivalent.

Remark 5.1. We only occasionally refer S by a general name: *statistic space*.

Remark 5.2. The model induced by a sample statistic is the same as the applied model from §3.2.1. A distinction is being made between the two concepts because sample statistic are only used to help make inference, whereas the applied model treats what values are going to be used as a source of entropy.

Strictly speaking, choice of the sample statistic s is essentially arbitrary. But we will argue that, depending on the purpose, some sample statistics are preferable to others.

Remark 5.3. Any statistic s on a singular probability model induces another singular probability model. However, if the initial singular model is uniform, the induced model is not necessarily uniform. The resulting lack of uniformity can be regarded as a means whereby one can make some inferences more easily.

Remark 5.4. Any surjective statistic s on an unrestricted probability model $U(X)$ induces another model that is isomorphic to unrestricted probability model $U(S)$, because for any probability distribution on S can be induced from some probability distribution on X. If s is not surjective, then the induced model is effectively the unrestricted model on the image of s extended so that elements of S outside the image of s always have probability zero.

5.2 Induced Inference

Given a statistic s on (Π, X, P) and an inference method I, we define an *(statistic) induced inference function* $j = j_{I,s}$ for (Π, X, P) by defining

$$j_{I,s}(x) = I(\Pi, S, Q)(s(x)), \tag{5.2}$$

recalling that $I(\Pi, S, Q)$ is an inference function for (Π, S, Q), and thus a function from S to assertions about $p \in \Pi$. We will sometimes say that this inference is *based* on the statistic s. (Given a statistic method and an inference method, one can similarly define a *(statistic) induced inference method*.)

Remark 5.5. The function $j_{I,s}$ is an inference function for (Π, X, P) but not for the induced probability model (Π, S, Q). Also, the inference method I will generally produce an inference function $I(\Pi, X, P)$ for (Π, X, P) which is different from the inference function $j_{I,s}$ for (Π, X, P).

Remark 5.6. In models admitting subuniform distributions, some conceptually important gradings, such as typicality, treat all subuniform distributions equally, resulting a tying effect. Sample statistics can be used as a tiebreaker in such models.

Remark 5.7. Strictly speaking, an inference function may itself be viewed as a sample statistic, because these categories of functions share the common domain of the sample space X. We will usually regard inference functions and sample statistics as entirely distinct entities because of their different purposes. Despite this distinction, the same functions can be useful as both inference function and sample statistics.

Generally, given two inference method I and J and a probability model (Π, X, P) one can define an inference function $j_{I,J(\Pi,X,P)}$, where the inference function $J(\Pi, X, P)$ serves the role of the sample statistic. Doing this effectively defines an *inference-induced* inference method J_I, which on input (Π, X, P) returns inference function $j_{I,J(\Pi,X,P)}$.

Specifically, for some models, using a maximal likelihood inference function may be a useful sample statistic.

Remark 5.8. For each $x \in X$, an artificial statistic is $s_x : X \to \{0, 1\}$ such that $s_x(x) = 1$ and $s_x(y) = 0$ for $y \neq 0$. The s_x induced inclusive typicality of x equals the likelihood of x if the likelihood is less than $\frac{1}{2}$. The induced inclusive likelihood for $y \neq x$ is 1. So, if hypothesis testing is being done, there exits arbitrary statistics that can accept any model based on any observation. And there exist statistic that can be used to reject any model, unless the model has a very high maximum likelihood for each sample value, meaning that it approaches pseudodeterminism and must have low inferred min-entropy.

5.3 Model-Neutral Statistics

The approach in this report for formally assessing cryptographic entropy tries to isolate almost all assumptions to the probability model, and relies on the presumption of the validity of the probability model. Unfortunately, inference encounters some difficulties in the context, such as the tying effect. Sample statistics may be useful to overcome these difficulties.

The probability model may have been from hard-won inference using a extensive effort. The assessment process, therefore, should attempt not to discard the gains made in the determination of the probabilty model. To this end, this section describes criteria for sample statistic being consistent with the probaiblty model.

Remark 5.9. For hypothesis testing, the very proability model is in question, so it may be less important to use sample statistics that are consistent with the model. Neverthless, the approach in this report to hypothesis is to formulate alternative hypothesis models, and then use sample statistic that are consistent with alternative model.

Firstly, some concepts are introduced. A bijection from the sample space to itself will be called a *sample transformation*. A sample transformation $z : X \to X$ is said to be *neutral* for model (Π, X, P), if all $p \in \Pi$ and all $x \in X$, the condition $P_p(x) = P_p(z(x))$.

Remark 5.10. A neutral transformation z describes an isomorphism from the model to itself, in which the isomorphism acts as the identity on the distributions.

Remark 5.11. Some natural neutral transformation for the independent model are those that permute the positions of entries in the sample vector x.

Remark 5.12. The set of neutral transformations for the independent model is the set of functions that permute the preimages of the frequency statistic (§5.4.2).

A transformation $z : X \to X$ is said to be *invariant* for a model if, for all $p \in \Pi$ and $x, y \in X$, if $P_p(x) = P_p(y)$, then $P_p(z(x)) = P_p(z(y))$.

Remark 5.13. Any neutral transformation is also an invariant transformation.

Remark 5.14. Some natural non-neutral invariant transformation for the independent model are those that permute the values of entries in the sample vector x.

The set of neutral transformations and the set of invariant transformations are implied by the probability model. Because the probability model is an assumption, the set of neutral transformations and the set of invariant transformation are implied assumptions.

A sample statistic s is *model-neutral* if for all neutral transformations $z : X \to X$, all $p \in \Pi$, and all $x \in X$, it holds that $Q_p(s(x)) = Q_p(s(z(x))$ where Q_p is the induced probability function.

Remark 5.15. A statistic s is model-neutral if $s(x) = s(z(x))$ for all x and all neutral transformations z. In this case, we say the statistic is *strongly model-neutral*.

A sample statistic s is *model-invariant* if for all invariant transformations $z : X \to X$, all $p \in \Pi$, and all $x, y \in X$, if $Q_p(s(x)) = Q_p(s(y))$, then $Q_p(s(f(x)) = Q_p(s(f(y)))$.

Model-neutral and model-invariant sample statistics are attempts to not contradict the assumptions made in the probability model. By making induced inference based on model-neutral or model-invariant sample statistics, one is not interfering too much with assumptions that have been made, neither doubting them nor strengthening them.

Remark 5.16. Given any statistic $s : X \to S$ for a model (Π, X, P), a model-neutral statistic $\hat{s} : X \to \hat{S}$ can be derived from s as follows. Let $\hat{S}$ be the set of multisets with elements in S. Let

$$\hat{s}(x) = \{s(z(x)) : z \in Z(\Pi, X, P)\} \tag{5.3}$$

where $Z(\Pi, X, P)$ is the set of neutral transformations of the model.

Remark 5.17. Sample statistic method can include inference methods, as already noted in Remark 5.7. The gradings likelihood typicality and generalized typicality can act as strongly model-neutral sample statistics. Likewise, the maximally graded and threshold graded inferences associated to these gradings are strongly model-neutral as sample statistics.

Remark 5.18. In the uniform model, all transformations are neutral and invariant. Consequently, the model-neutral and model-invariant statistics are precisely those that induce a uniform distribution. So, model-neutral and model-invariant statistics for the uniform cannot be used as tiebreakers and cannot overcome the tying effect.

5.4 Sample Statistics for the Independent Probability Model

For the task of estimating the min-entropy in the uniform or independent model, the following statistics seem as though they may be appropriate, given their natural relation to the estimate of min-entropy that one gets by examining the frequencies of the sample.

5.4.1 Identity

Strictly speaking, the identity function itself is a sample statistic. We call this the *identity statistic*, but inference based on the identity statistic is the same as direct inference.

5.4.2 Frequency

A straightforward sample statistic that is the *frequency* statistic:

$$f : \{0, 1, \ldots, m-1\}^N \to \{0, 1, \ldots, N\}^m \tag{5.4}$$

such that

$$f(x)_i = |\{k : x_k = i\}| \tag{5.5}$$

so $f(x)_i = j$ is the number of k such that $x_k = i$ is j. Here, entries $f(x)$ are indexed starting from 0.

Remark 5.19. For example, if $(m, N) = (3, 4)$ then $f(0, 2, 2, 2) = (1, 0, 3)$. To see this, note $x = (0, 2, 2, 2)$ and $f(x)_0 = 1$ because that set of k such that $x_k = 0$ is simply the value $k = 1$ (if x is indexed starting from 1).

The frequency statistic is also related to related to the probability function of the independent model by the following formula:

$$P_p(x) = p^{f(x)} \tag{5.6}$$

where the notation a^b for vectors a and b of equal length, such as p and $f(x)$ which both have length m, means $\prod a_i^{b_i}$, where the product ranges over the set of indices of the vectors.

Remark 5.20. It follows from (5.6) that a sample transformation z is neutral for the independent model if and only if it satisfies $f(x) = f(z(x))$ for all x. Hence Remark 5.12.

Remark 5.21. It follow from (5.6) that a sample transformation z is invariant for the independent model if and only if $f(z(x)) = f(z(y))$ for all x, y such that $f(x) = f(y)$.

Remark 5.22. The maximal likelihood inference $\hat{p}(x)$ from x is related to this statistic by simple division $\hat{p}(x) = f(x)/N$.

The induced probability model has probability function which can be written as:

$$Q_p(v) = \binom{N}{v} p^v \tag{5.7}$$

using the same vector exponentiation notation as in (5.6), and multinomial notation where $\binom{N}{v} = \frac{N!}{\prod v_i!}$, where the product in the denominator again ranges over the set of indices of the vector.

Remark 5.23. For fixed x, the probability function Q_p is proportional to P_p. Consequently, the likelihood functions are proportional, and will give to rise to the same inferences under maximal likelihood inference.

Remark 5.24. The typicality grading will differ more because the factor $\binom{N}{f(x)}$ can change the ordering of the probabilities.

Remark 5.25. The frequency sample statistic is model-neutral, because any neutral transformation z preserves the value of $f(x)$, and the induced probability function is determined by $f(x)$ as in (5.7).

Remark 5.26. The frequency sample statistic is model-invariant, because $Q_p(f(x)) = Q_p(f(y))$ for all p, if and only if, $f(x) = f(y)$.

5.4.3 Partition

Another sample statistic for the independent model the *partition* statistic ϕ where $\phi(x)$ is $f(x)$ resorted in non-ascending order.

Remark 5.27. For example, if $(m, N) = (3, 4)$, then $\phi(0, 2, 2, 2) = (3, 1, 0)$. For another example with the same (m, N), we have $\phi(0, 0, 2, 2) = (2, 2, 0)$.

The partition statistic is also a statistic on the probability model induced by the frequency statistic. Thus we say that ϕ is a *coarser* statistic than f, or that ϕ is a *coarsening* of f. Let ξ be the sorting function, so that $\phi = \xi \circ f$.

The probability distribution induced by the partition statistic has sample space which is the set of partitions of N of length m (or at most length m, if we ignore entries of value 0.) For partition θ, the induced probability distribution has

$$Q_p(\theta) = \binom{N}{\theta} \sum_{v:\xi(v)=\theta} p^v = \binom{N}{\theta} m_\theta(p) \tag{5.8}$$

where m_θ is the monomial symmetric function, in the notation used by Macdonald [Mac95].

Remark 5.28. If z is neutral for the independent model, then by Remark 5.20, $f(x) = f(z(x))$. It follows that $\phi(x) = \xi(f(x)) = \xi(f(z(x)) = \phi(z(x))$. Therefore, $Q_p(x) = Q_p(z(x))$, so ϕ is a model-neutral statistic.

5.4.4 Mode

A another nontrivial statistic we consider is the *mode* statistic μ, which the maximal entry in the frequency vector (so first in the partition vector). For an example of the mode statistic, consider again $(m, N) = (3, 4)$ and $x = (0, 2, 2, 2)$. Then $\phi_1(0, 2, 2, 2) = 3$.

The maximum likelihood inference for min-entropy at x is $-\log_2(\mu(x)/N)$. Inference induced from the mode statistic are the same as those induced from using the maximal likelihood estimate for min-entropy as a sample statistic.

The mode statistic is a coarsening of the partition statistic. It is the coarsest of the statistics for the independent model described in this report.

5.5 Statistics for the Markov Model

5.5.1 Markov Frequency Statistic

A natural generalization to the Markov model of the frequency sample statistic for the independent model is the following (Markov) frequency statistic. Given $x \in X = \{0, 1, \ldots, m-1\}^N = (x_0, \ldots, x_{N-1})$, the Markov frequency

statistic is

$$F(x) = (e(x), U(x)) \tag{5.9}$$

where: $e(x) = e_{x_0}$ is a m dimensional vector all of whose entries are zero except for the entry in position x_0 whose value is 1 (vector entry indices run from 0 to $m-1$); and $U(x)$ is an $m \times m$ matrix with non-negative integer entries $U_{i,j}$ indexed by integers pair (i, j) such that $0 \leqslant i, j < m$ with:

$$U_{i,j} = |\{k | 1 \leqslant k \leqslant N-1,\ x_{k-1} = i,\ x_k = j\}| \tag{5.10}$$

In words, $F(x)$ marks the initial state and the number of transitions between the various states in x. Formally, F is a function $F : X \to S$, where S is the statistic space. We can take S to be the set of matrices of all pairs of vectors and matrices of dimension m, with non-negative integers, such that the sum of vector entries is 1 and the sum of the matrix entries is $N-1$.

Remark 5.29. We could also take the statistic space to this image, $S = F(X)$, which is a proper subset of all the non-negative integer m-dimensional vector-matrix pairs.. The resulting induced model is weakly isomorphic.

The induced model will have form (Π, S, Q) and the induced probability function $Q : \Pi \times S \to [0, 1]$ is

$$Q(p, s) = Q((v, M), (e, U)) = \pi(e, U) v^e M^U \tag{5.11}$$

where: $\pi(e, U)$ is an integer counting the number of sequences x such that $F(x) = (e, U)$; and the notation a^b and A^B for vector and matrix exponentiation indicates entry-wise exponentiation followed by taking the product over all of the entries (with the convention that $0^0 = 1$).

Formula (5.11) is because the original probability function may be also computed using the frequency sample statistic by the related formula:

$$P_p(x) = P_{(v,M)}(x) = v^{e(x)} M^{U(x)} \tag{5.12}$$

Remark 5.30. Formula (5.12) implies that sample values x and y are equilikely if and only if $F(x) = F(y)$.

Remark 5.31. The induced likelihood function $L_{F(x)}$ is proportional to the likelihood function L_x, by a factor $\pi(F(x))$. So, the induced model does not alter the maximal likelihood inference.

Such proportionality can fail between typicality and induced typicality, so using the Markov frequency statistic can alter typicality-based inferences.

A combinatorial description for $\pi(e, U)$ is as follows: $\pi(e, U)$ is the number of sequences in $(0, \ldots, m-1)^N$ that begin with i if $e_i = 1$, and that have $U_{j,k}$ occurrences of the two adjacent element subsequence (j, k).

Remark 5.32. A related description of $\pi(e, U)$ is as follows. For any matrix U with non-negative integer entries, define a matrix $V = \hat{U}$ such that $V_{i,j}$ is the number of sequences beginning with i and ending with j, and containing exactly $U_{k,l}$ consecutive entries in the sequence of the form (k, l). Then:

$$\pi(e, U) = e \hat{U} f, \tag{5.13}$$

where e is viewed as a row vector and f is a column vector with all entries equal to one.

Remark 5.33. The matrix operator $U \mapsto \hat{U}$ from Remark 5.32 has a role in matrix powering. Let X be any square matrix and let n be a non-negative integer. Then

$$X^n = \sum_U X^U \hat{U} \tag{5.14}$$

where the sum ranges over U with the same shape as X and non-negative integer entries summing to n, and X^U indicates, as above, applying entry-wise exponentiation and taking the product of the all the power entries. For example, if X is a 2 by 2 matrix, then:

$$X^2 = X^{\left(\begin{smallmatrix}2&0\\0&0\end{smallmatrix}\right)} \left(\begin{smallmatrix}1&0\\0&0\end{smallmatrix}\right) + X^{\left(\begin{smallmatrix}1&1\\0&0\end{smallmatrix}\right)} \left(\begin{smallmatrix}0&1\\0&0\end{smallmatrix}\right) + X^{\left(\begin{smallmatrix}0&1\\1&0\end{smallmatrix}\right)} \left(\begin{smallmatrix}1&0\\0&1\end{smallmatrix}\right) + X^{\left(\begin{smallmatrix}0&1\\0&1\end{smallmatrix}\right)} \left(\begin{smallmatrix}0&1\\0&0\end{smallmatrix}\right) + X^{\left(\begin{smallmatrix}1&0\\0&0\end{smallmatrix}\right)} \left(\begin{smallmatrix}0&0\\1&0\end{smallmatrix}\right) + X^{\left(\begin{smallmatrix}0&0\\1&1\end{smallmatrix}\right)} \left(\begin{smallmatrix}0&0\\1&0\end{smallmatrix}\right) + X^{\left(\begin{smallmatrix}0&0\\0&2\end{smallmatrix}\right)} \left(\begin{smallmatrix}0&0\\0&1\end{smallmatrix}\right) \tag{5.15}$$

So, taking a general $X = \left(\begin{smallmatrix}x_{00}&x_{01}\\x_{10}&x_{11}\end{smallmatrix}\right)$, which means $X^2 = \left(\begin{smallmatrix}x_{00}^2+x_{01}x_{10} & x_{01}x_{11}+x_{00}x_{01}\\ x_{10}x_{00}+x_{11}x_{10} & x_{11}^2+x_{10}x_{01}\end{smallmatrix}\right)$, the monomials, for example, $x_{00}^2 = X^{\left(\begin{smallmatrix}2&0\\0&0\end{smallmatrix}\right)}$ and $x_{01}x_{10} = X^{\left(\begin{smallmatrix}0&1\\1&0\end{smallmatrix}\right)}$ contribute to X^2 by scaling of $\left(\begin{smallmatrix}1&0\\0&0\end{smallmatrix}\right)$ and $\left(\begin{smallmatrix}1&0\\0&1\end{smallmatrix}\right)$ respectively.

Remark 5.34. In the case of two-by-two square matrices the matrix operator $U \mapsto \hat{U}$ from Remark 5.32 can be described as follows:

$$\widehat{\begin{pmatrix} a & b \\ c & d \end{pmatrix}} = \begin{cases} \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} & \text{if } |b-c| > 1 \\ \begin{pmatrix} 0 & \binom{a+c}{a}\binom{d+c}{d} \\ 0 & 0 \end{pmatrix} & \text{if } b = c+1 \\ \begin{pmatrix} 0 & 0 \\ \binom{a+b}{a}\binom{d+b}{d} & 0 \end{pmatrix} & \text{if } c = b+1 \\ \begin{pmatrix} \binom{a+c}{a}\binom{d+c-1}{d} & 0 \\ 0 & \binom{a+c-1}{a}\binom{d+c}{d} \end{pmatrix} & \text{if } b = c \end{cases} \tag{5.16}$$

with conventions $\binom{-1}{0} = 1$ and $\binom{m}{n} = 0$ if $m < n > 0$.

Remark 5.35. Goulden and Jackson [GJ83, Ex. 2.4.21] give a formula that determines $\pi(e, U)$. Suppose that $e_a = 1$. Let $k_j = \delta_{a,j} + \sum_{i=0}^{m-1} U_{i,j}$, where $\delta_{a,j} = 0$ if $j \neq 0$ and $\delta_{a,a} = 1$. If there exists some $b \in \{0, 1, \ldots, m-1\}$ such that $k_i = \delta_{j,b} + \sum_{i=0}^{m-1} U_{i,j}$, then

$$\pi(e, U) = \frac{\prod_{j=0}^{m-1}(k_j - 1)!}{\prod_{0 \leqslant i,j < m} U_{i,j}!} \det(K - U) \tag{5.17}$$

where K is the diagonal matrix with entry k_j at position (j, j). If no such b exists, then $\pi(e, U) = 0$.

This formula requires that all $k_i > 0$, but can easily be adapted to handle instances $k_i = 0$ by removing such i from all consideration, and re-indexing and re-computing for only those $k_i > 0$.

Remark 5.36. Goulden and Jackson's formula (5.17) determines $\hat{U}$. The conditions on the vector k mean that most entries in $\hat{U}$ are zero, while remainder can be computed as a determinant. Let f be the column vectors of all ones. Let U' be the transpose of U. Let $w = (U - U')f$. The conditions on the vector k implying the following.

1. If $w = 0$, then $\hat{U}$ will be a diagonal matrix.
2. If w has entry 1 in position i and entry -1 in position j with all other entries equal to 0, then $\hat{U}$ is a matrix with all entries equal to zero except the entry at position (i, j).
3. For any other value of w, the matrix $\hat{U}$ is all zeros.

Remark 5.37. Goulden and Jackson's formula is also related to the BEST theorem of de Bruijn, van Ardenne-Ehrenfest, Smith and Tutte on the number of Euler circuits in a directed graph.

5.5.2 Maximum Likelihood Markov Statistic

In general, in any probability model, the maximum likelihood inference can be used as a sample statistic. In the case of the Markov model, the maximum likelihood inference is closely related to the frequency statistic. With some exceptions, the maximum likelihood inference can be derived from the frequency rows by dividing each row of the frequency matrix by its sum. So,

$$\hat{p}(x) = (\hat{v}(x), \hat{M}(x)) \tag{5.18}$$

where $\hat{v}(x) = e(x)$ with $e(x)$ the first component of the frequency statistic $F(x)$, and

$$\hat{M}(x)_{i,j} = \frac{|\{k | (x_{k-1}, x_k) = (i, j)\}|}{|\{k | x_{k-1} = i\}|}. \tag{5.19}$$

The exceptional cases occur when some of the row sums of $U(x)$ are zero. In these exceptionsal cases, the corresponding rows of M have no effect on the probability of x. Therefore, the maximum likelihood inference in these cases is a set where the exceptional rows can take on any legal value.

Formally, the maximum likelihood statistic can viewed as the statistic $\hat{p} : X \to S$, with $S = [\Pi]$, meaning the set of all subsets of Π. Although the set S is uncountable, the image $\hat{p}(X) \subset S$ is a finite.

Remark 5.38. We could also take the statistic space to this image, $S = \hat{p}(X)$. The induced models are weakly isomorphic.

Although the Markov frequency statistic determines the maximum likelihood statistic, the converse can fail: the value of maximum likelihood statistic on the Markov model does not uniquely determine the frequency statistic. For example, in the $(3, 8)$ Markov model, consider the sample values

$$x = (1, 0, 2, 0, 0, 2, 0, 1), \tag{5.20}$$

$$y = (1, 0, 2, 0, 1, 0, 0, 2). \tag{5.21}$$

Their frequency statistics are:

$$F(x) = \left(\begin{pmatrix} 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 2 \\ 1 & 0 & 0 \\ 2 & 0 & 0 \end{pmatrix} \right), \tag{5.22}$$

$$F(y) = \left(\begin{pmatrix} 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 2 \\ 2 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \right), . \tag{5.23}$$

which are different. Their maximum likelihood statistics are

$$\hat{p}(x) = \hat{p}(y) = \left(\begin{pmatrix} 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} \frac{1}{4} & \frac{1}{4} & \frac{1}{2} \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \right). \tag{5.24}$$

Therefore, models induced by the frequency statistic and the maximum likelihood statistic may not be equivalent. In the earlier terminology, maximum likelihood is a coarser statistic than Markov frequency.

5.5.3 Runs Test

The nummber of runs of equal elements in a sequence x equals one plus the sum of the off-diagonal entries of the matrix $F(x)$. Because $P_p(x)$ is calculated from $(e(x), F(x))$, the number of runs in x is a model-neutral statistic in the Markov model.

5.5.4 Maximal Likelihood Min-Entropy Statistic

As above, inference methods may perhaps also appropriate as sample statistics. This suggests to use the inferred min-entropy over the maximum likelihood inference, specifically taking the infimum value of min-entropy over the inferred set of distributions.

6 Examples

This section provides some illustrative examples of entropy assessment in various models. In most of the examples, the optimization problems arises are easily solved. In a few examples, only the formulation of the optimization problem is given.

6.1 Toy Example in Independent Model

In this section the probability model is the $(2, 3)$-independent model from §2.3.2.

Remark 6.1. We may think of this model consisting of three coin flips, with 1 indicating a coin lead its head up, and 0 its tail up. Thus p_1 is the probability that a coin lands heads up.

Remark 6.2. A main reason for analyzing this toy model is to illustrate, with hand calculations, how the various entropy assessment approaches work.

Remark 6.3. Because this toy model is so small, the optimization problems arising from the process of statistical inference are generally easy to solve.

Remark 6.4. In certain real-world application, the optimization problems arising may be quite difficult to solve.

In all of the following examples, the sample will be $x = (0, 1, 1)$.

Remark 6.5. The amount of information in the sample is quite small. In other words, the sample size is small.

Remark 6.6. One effect of small sample size should be a lower confidence in the inference. Or, more precisely, a wide range of inferences at a given a confidence.

Remark 6.7. On the one hand, the expected wider range of inference may help highlights the differences between the various inference methods. On the other hand, because of the smallness of the sample, the results of various inference methods in this example should not be used as a means to evaluate or compare the various inference methods.

Remark 6.8. For prudence, cryptographers generally wish to take the minimum value of min-entropy. A widening of the range of distributions may lower this minimum value of min-entropy. In other words, a small sample size may result in an entropy estimate lower than the actual amount of entropy, for a given level of confidence.

6.1.1 Simplified Description of the Model

The probability space Π of the $(2, 3)$-independent model is the set

$$\Pi = \{(p_0, p_1) : p_0, p_1 \in [0, 1], p_0 + p_1 = 1\} \tag{6.1}$$

In the following examples, we will use a simpler but isomorphic model in which $\Pi = [0, 1]$. The isomorphism maps $p = (p_0, p_1)$ from the original model to $p = p_1$ in the simpler model. In the other direction, p in the simpler model maps to $(1 - p, p)$ in the original model.

Remark 6.9. The simpler model reduces the number of variables from two, namely p_0 and p_1, to one, p, and also avoids the use of subscripts. Variable p_1 has been chosen to map to p because p becomes the expected of each entry x_i. Less notation in the examples may better illustrate the ideas.

Remark 6.10. Such a simpler but isomorphic model could be used $(2, N)$ independent model. In the (m, N) independent model for larger m, there is less advantage.

6.1.2 Maximal Likelihood

For $x = (0, 1, 1)$, the likelihood grading function is

$$L_x(p) = (1 - p)p^2. \tag{6.2}$$

To help maximize L_x over the probability space $[0, 1]$, we can consider the derivative




Figure 1: Likelihood, Toy Example, with $x = (0, 1, 1)$

$$\frac{dL_x(p)}{dp} = L'_x(p) = 2p - 3p^2 = p(2 - 3p) \tag{6.3}$$

The critical points of the likelihood function, where $L'_x = 0$, are at $p = 0$ and $p = \frac{2}{3}$. For the purposes of maximization, we must also consider the boundary of the space Π, which occurs at $p = 0$, and $p = 1$. Therefore, we just need to evaluate L_x on each element of the vector $(0, \frac{2}{3}, 1)$, which gives $(0, \frac{4}{27}, 0)$. Therefore, the likelihood function attains its maximal value $\frac{4}{27}$ precisely at $p = \frac{2}{3}$.

The set-valued maximal likelihood inference for p is therefore for the set $\{\frac{2}{3}\}$. In a cryptographic application, we may want infer something about a probability parameter. Since the inferred set is a singleton set, for simplicity, we will speak of inferred values for the following discussions. The inferred probability distribution is $\hat{p} = \frac{2}{3}$.

The min-entropy $H_\infty(\hat{p})$ of the probability distribution $\hat{p}$ is as follows. Recall that the inferred probability distribution is $\hat{p} = \frac{2}{3}$. The value of x that maximizes $P_{\hat{p}}(x)$ is $\hat{x} = (1, 1, 1)$, and this gives $P_{\hat{p}}(\hat{x}) = \frac{8}{27}$. The min-entropy is therefore $-\log_2\left(\frac{8}{27}\right)$ which is approximately 1.75 bits of inferred min-entropy for p.

Remark 6.11. The sample used for inference $x = (0, 1, 1)$ and the sample $\hat{x} = (1, 1, 1)$ used to calculate min-entropy of the probability distribution inferred from x are not equal. In particular, the sample entropy (information content) of x is higher than the min-entropy of p.

6.1.3 Threshold Inclusive Typicality

The inclusive typicality with $x = (0, 1, 1)$, works out to be:

$$g_1(x, p) = \begin{cases} 3p^2 - 2p^3 & \text{if } 0 \leqslant p < \frac{1}{2}, \\ 1 & \text{if } p = \frac{1}{2}, \\ 1 - p^3 & \text{if } \frac{1}{2} < p \leqslant 1, \end{cases} \tag{6.4}$$

because: when $p > 1/2$, the only sample value y with $P_p(y) > P_p(x)$ is $y = (1, 1, 1)$, which has probability $P_p(y) = p^3$; when $p = 1/2$, the distribution is uniform so the inclusive typicality sums over all samples resulting in 1; when $p < 1/2$, the set of sample values y with $P_p(y) \leqslant P_p(x)$, is $\{(1, 1, 1), (0, 1, 1), (1, 0, 1), (1, 1, 0)\}$, whose sum of probabilities is $3p^2(1 - p) + p^3 = 3p^2 - 2p^3$. See Figure 2.




Figure 2: Inclusive Typicality, Toy Example, with $x = (0, 1, 1)$

Remark 6.12. The inclusive typicality evaluated at the maximum likelihood estimate for $\hat{p}$, from §6.1.2 therefore works out to be $\frac{19}{27}$. The corresponding confidence level is therefore $\frac{8}{27}$.

Remark 6.13. Inclusive typicality always takes value 1 at a the uniform distribution p, if the probability model admits a uniform distribution. More generally, it will take value 1 at subuniform distributions (see Remark 2.24. If inclusive typicality is used to formulate a maximally graded inference, then such uniform (and subuniform) distributions will always belong to the inferred set.

If the inferred set consists of only the uniform distribution, then it maximal inclusive typicality may seem to be a too optimistic inference method for use in cryptography. In other cases, the inferred set of distributions may contain other distributions, and by narrowing the inferred set of entropies by taking its infimum, one may not end up with an overly optimistic inference.

For each threshold value $t \in [0, 1]$, an inferred set of distributions may be given such that $g_{1/2}(x, p) > t$, as follows:

$$i_{g_1 > t}(x) = \begin{cases} \varnothing & \text{if } t = 1 \\ \{\frac{1}{2}\} & \text{if } \frac{7}{8} \leqslant t < 1 \\ \left[\frac{1}{2}, \sqrt[3]{1-t}\right) & \text{if } \frac{1}{2} \leqslant t < \frac{7}{8} \\ \left(q(t), \sqrt[3]{1-t}\right) & \text{if } 0 \leqslant t < \frac{1}{2} \end{cases} \tag{6.5}$$

where $q(t)$ is the unique value in $[0, \frac{1}{2})$ such that $3q(t)^2 - 2q(t)^3 = t$.

The corresponding inferred sets of min-entropy values are then given by:

$$H_\infty\left(i_{g_1 > t}(x)\right) = \begin{cases} \varnothing & \text{if } t = 1 \\ \{3\} & \text{if } \frac{7}{8} \leqslant t < 1 \\ (-\log_2(1-t), 3] & \text{if } 0 \leqslant t < \frac{7}{8} \end{cases} \tag{6.6}$$

Remark 6.14. Recall that $H_\infty(p)$ depends on the maximum value of $P_p(x)$, which will either by p^3 or $(1-p)^3$.

Cryptographers, in the interest of practicality may wish to narrow the inferred set of min-entropies to a single value, and, moreover, do so by taking the infimum of the set, for the sake of prudence. Instead of thresholds, one may consider confidence level $c = 1 - t$. For each confidence level $c \in [0, 1]$, the inferred $H_\infty(c)$ value is given by

$$H_\infty(c) = \begin{cases} \infty & \text{if } c = 0 \\ 3 & \text{if } 0 < c \leqslant \frac{1}{8} \\ -\log_2(c) & \text{if } \frac{1}{8} < c \leqslant 1 \end{cases} \tag{6.7}$$

Remark 6.15. In this example, the infima of the inferred sets do not actually belong to the sets.

Remark 6.16. By convention, the infimum of the empty set is (positive) infinity. An inference of infinite amount of entropy in a 3-bit random variable is clearly absurd, but this inference is only made at confidence level 0.

Remark 6.17. The sample entropy (information content) of $x = (0, 1, 1)$, which will be treated later, is certainly higher than the inferred min-entropy of p at high confidence levels. Intuitively, at a high confidence levels, close to 1, we must account for the possibility that p is high, and therefore $(1, 1, 1)$ would have been much more likely than $(0, 1, 1)$.

In this situation, cryptographers may recognize that the sample entropy in $(0, 1, 1)$ is at least $\log_2(3)$, because we have assumed the independent model, and the 0 bit could have occurred in any of the three locations. Such sample entropy is therefore useful in cryptography, so we will address its inference in full formality.

6.1.4 Threshold Balanced Typicality

The balanced typicality with $x = (0, 1, 1)$, works out to be:

$$g_{\frac{1}{2}}(x, p) = \begin{cases} \frac{3}{2}p^2 - \frac{1}{2}p^3 & \text{if } 0 \leqslant p < \frac{1}{2} \\ \frac{1}{2} & \text{if } p = \frac{1}{2} \\ 1 - \frac{3}{2}p^2 + \frac{1}{2}p^3 & \text{if } \frac{1}{2} < p \leqslant 1 \end{cases} \tag{6.8}$$




Figure 3: Balanced Typicality, Toy Example, with $x = (0, 1, 1)$

Remark 6.18. The third expression above may be derived as $(1-p)^3 + 3p(1-p)^2 + \frac{3}{2}p^2(1-p) = 1 - \frac{3}{2}p^2 + \frac{1}{2}p^3$.

Remark 6.19. The balanced typicality evaluated at the maximum likelihood estimate for $\hat{p}$, from §6.1.2 therefore works out to be $\frac{13}{27}$. The confidence level is therefore $\frac{14}{27}$.

Remark 6.20. Balanced typicality is maximized as p approaches $\frac{1}{2}$ from above. Thus typicality and likelihood, in this toy example, give much difference inferences under maximization. In particular, in this case, typicality may appear too optimistic. It would go too far to conclude from this toy example, however, that maximal typicality is always too optimistic.

For each threshold value $t \in [0, 1]$, an inferred set of distributions may be given such that $g_{1/2}(x, p) > t$, as follows:

$$i_{g_{\frac{1}{2}} > t}(x) = \begin{cases} \varnothing & \text{if } \frac{11}{16} \leqslant t \leqslant 1 \\ \left(\frac{1}{2}, q(t)\right) & \text{if } \frac{1}{2} \leqslant t < \frac{11}{16} \\ \left[\frac{1}{2}, q(t)\right) & \text{if } \frac{3}{16} \leqslant t < \frac{1}{2} \\ (r(t), q(t)) & \text{if } 0 \leqslant t < \frac{3}{16} \end{cases} \tag{6.9}$$

where $q(t)$ is the unique value in $(\frac{1}{2}, 1]$ such that $1 - \frac{3}{2}q(t)^2 + \frac{1}{2}q(t)^3 = t$, and $r(t)$ is the unique value in $[0, \frac{1}{2})$ such that $\frac{3}{2}r(t)^2(1 - r(t)) = t$.

The corresponding inferred sets of min-entropy values are then given by:

$$H_\infty\left(i_{g_{\frac{1}{2}}>t}(x)\right) = \begin{cases} \varnothing & \text{if } \frac{11}{16} \leqslant t \leqslant 1 \\ (-3\log_2(q(t)), 3) & \text{if } \frac{1}{2} \leqslant t < \frac{11}{16} \\ (-3\log_2(q(t)), 3] & \text{if } 0 \leqslant t < \frac{1}{2} \end{cases} \tag{6.10}$$

Remark 6.21. Recall that $H_\infty(p)$ depends on the maximum value of $P_p(x)$, which will either by p^3 or $(1-p)^3$, hence the factor of 3 appearing above.

Cryptographers, in the interest of practicality may wish to narrow the inferred set of min-entropies to a single value, and, moreover, do so by taking the infimum of the set, for the sake of prudence. Instead of thresholds, one may consider confidence level $c = 1 - t$. For each confidence level $c \in [0, 1]$, the inferred $H_\infty(c)$ value is given by

$$H_\infty(c) = \begin{cases} \infty & \text{if } 0 \leqslant c \leqslant \frac{5}{16} \\ -3\log_2(q(1-c)) & \text{if } \frac{5}{16} < c \leqslant 1 \end{cases} \tag{6.11}$$

Remark 6.22. For a given confidence level, in this toy example, using balanced typicality as the grading to be threshold generally gives a higher inference of min-entropy than inclusive typicality. Formally, this is because inclusive typicality is always at least balanced typicality, and therefore its inferred sets contain the inferred sets from balanced typicality. When we infer a value of min-entropy by taking an infimum, we arrive at the inclusive inference being at most the balanced inference.

Remark 6.23. In this example, the infima of the inferred sets do not actually belong to the sets.

Remark 6.24. The infimum of an empty set is, by convention, taken to be infinite. The inference of infinite entropy in three bits is absurd, but this inference is only made at low confidence levels.

6.1.5 Maximal Adjusted Likelihood




Figure 4: Adjusted Likelihood, Toy Example, with $x = (0, 1, 1)$

Adjusted likelihood t_{σ_1}, is a special case of generalized typicality (4.12) where $\sigma_1(x) = x$. In this toy example, adjusted likelihood t_{σ_1}, with $x = (0, 1, 1)$, works out to be:

$$t_{\sigma_1}(x, p) = \frac{1}{2}\left(1 + p^2(1-p) - ((1-p)^2 + p^2)^3\right) = \frac{1}{2}(6p - 17p^2 + 31p^3 - 36p^4 + 24p^5 - 8p^6) \tag{6.12}$$

This adjusted likelihood grading seems to have a maximum at around $\hat{p} \approx 0.559$. The corresponding inference for min-entropy is about 2.52 bits, considerably larger than the inference made with the maximum likelihood inference.

6.1.6 Threshold Adjusted Likelihood

The inferred set of min-entropies for a given threshold level t and sample value $x = (0, 1, 1)$ is given by:

$$H_\infty(i_{t_\iota > t}(x)) = \begin{cases} \infty & \text{if } t \geqslant t_\iota(x, \hat{p}) \approx 0.504 \\ (-3 \log_2 q(t), 3] & \text{if } 0 \leqslant t < t_\iota(x, \hat{p}) \end{cases} \tag{6.13}$$

where $\hat{p}$ is maximal value of adjusted likelihood as described in §6.1.5, and $q(t)$ is now the unique solution in the interval of $[\hat{p}, 1]$ of $t_\iota(x, q(t)) = t$.

At threshold and confidence level $c = t = 1/2$, the taking the infimum of the inferred min-entropy given an estimate of about 2.11 bits of min-entropy.

6.1.7 Frequency Statistic Induced Inferences

The value of the frequency statistic at sample $x = (0, 1, 1)$ is $v = f(x) = (1, 2)$. The induced probability for $v = (1, 2)$ is

$$Q_p(v) = 3p^2(1 - p). \tag{6.14}$$

6.1.7.1 Maximal Induced Likelihood The induced likelihood is $L_v(p) = 3L_x(p)$ where L_x is the direct likelihood. So, L_v is maximized at the same value of L_x, which is $\hat{p} = \frac{2}{3}$. The induced inference for min-entropy is therefore the same as direct inference: about 1.75 bits.

6.1.7.2 Induced Inclusive Typicality The frequency statistic value $v = (1, 2)$ has induced inclusive typicality 1 whenever

$$p \in \left[\tfrac{1}{2}, \tfrac{3}{4}\right]. \tag{6.15}$$

Taking the minimum value of the min-entropy over this range gives an inference of only about 1.24 bits.

Remark 6.25. The directed inference using maximal inclusive typicality was 3 bits of min-entropy, so the use of sample statistic induced inference has, in this case, reduced the entropy estimate, even though the same inference method was used, namely maximal inclusive

Remark 6.26. This example shows that the maximal inclusive typicality is not always too optimistic. Indeed, in this case, it seems to be slightly too pessimistic.

More generally the induced inclusive typicality at $x = (0, 1, 1)$ is the function

$$g_1(x, p) = \begin{cases} 3p^2 - 2p^3 & \text{if } 0 \leqslant p < \frac{1}{1+\sqrt{3}} \approx 0.366 \\ 1 - 3p + 6p^2 - 3p^3 & \text{if } \frac{1}{1+\sqrt{3}} \leqslant p < \frac{1}{2} \\ 1 & \text{if } \frac{1}{2} \leqslant p \leqslant \frac{3}{4} \\ 1 - p^3 & \text{if } \frac{3}{4} < p \leqslant 1 \end{cases} \tag{6.16}$$

6.1.7.3 Induced Balanced Typicality Induced balanced typicality at $x = (0, 1, 1)$ is the function

$$g_{1/2}(x, p) = \begin{cases} \frac{3}{2}p^2 - \frac{1}{2}p^3 & \text{if } 0 \leqslant p < \frac{1}{1+\sqrt{3}} \approx 0.366 \\ \frac{11}{2} - 3\sqrt{3} \approx 0.304 & \text{if } p = \frac{1}{1+\sqrt{3}} \\ 1 - 3p + \frac{9}{2}p^2 - \frac{3}{2}p^3 & \text{if } \frac{1}{1+\sqrt{3}} < p < \frac{1}{2} \\ \frac{5}{8} & \text{if } p = \frac{1}{2} \\ 1 - \frac{3}{2}p^2 + \frac{3}{2}p^3 & \text{if } \frac{1}{2} < p < \frac{3}{4} \\ \frac{37}{64} & \text{if } p = \frac{3}{4} \\ 1 - \frac{3}{2}p^2 + \frac{1}{2}p^3 & \text{if } \frac{3}{4} < p \leqslant 1 \end{cases} \tag{6.17}$$

As p approaches $\frac{1}{2}$ from above, balanced typicality a value of $\frac{13}{16} \approx 0.81$. The supremum of the balanced typicalities is $\frac{13}{16}$, although this value is never attained. One could interpret the maximal frequency-induced balanced typicality to occur at $\hat{p} = \frac{1}{2} + \epsilon$, for arbitrarily small $\epsilon > 0$. The resulting inference for entropy is about $3 - \epsilon$ bits, for arbitrarily small $\epsilon > 0$.

In this case, the maximal frequency-induced balanced typicality still seems to produce a inference about min-entropy that is too optimistic.

Remark 6.27. Balanced typicality actually has a local minimum at $p = \frac{2}{3}$, where it takes value $\frac{7}{9} \approx 0.78$. As p approaches $\frac{3}{4}$ from below, the typicality approaches $\frac{101}{128} \approx 0.79$. At threshold levels, between $\frac{7}{9}$ and $\frac{101}{128}$ the threshold inferred set is not connected.

6.1.7.4 Induced Adjusted Likelihood

The frequency-induced adjusted likelihood at $x = (0, 1, 1)$ is

$$t_{\sigma_1}(x, p) = 3p - \frac{21}{2}p^2 + \frac{53}{2}p^3 - 39p^4 + 30p^5 - 10p^6 \tag{6.18}$$

This function seems to have a maximum at around $p \approx 0.628$, so that the maximal frequency-induced adjusted likelihood estimate for min-entropy is about 2.02 bits.

Remark 6.28. In this case, the effect of inducing on the sample statistic frequency, has reduced the maximum adjusted likelihood estimate.

6.1.8 Partition Statistic Induced Inferences

The value of the partition statistic at sample $x = (0, 1, 1)$ is $\theta = \phi(x) = (2, 1)$. The only other possible value of the partition statistics is $\theta' = (3, 0)$. The induced probability for $\theta = (2, 1)$ is

$$Q_p(\theta) = 3p(1 - p) \tag{6.19}$$

6.1.8.1 Maximal Induced Likelihood

So the induced likelihood is $L_\theta(p) = 3p(1-p) = 3\left(\frac{1}{4} - \left(p - \frac{1}{2}\right)^2\right)$. This form of the likelihood shows to be maximized at $\hat{p} = \frac{1}{2}$. The resulting inference for min-entropy is 3 bits.

Remark 6.29. That the partition statistic essentially ignores the values of the entries may in part explain why the maximal induced likelihood distribution does not favor 1 or 0.

6.1.8.2 Maximal Induced Inclusive Typicality

Because the partition statistic in the $(2, 3)$ independent model can only takes two values, statistic value $\theta = (2, 1)$ has induced inclusive typicality 1 when $3p(1 - p) \geqslant \frac{1}{2}$, which holds whenever

$$p \in \left[\frac{1}{2} - \sqrt{\frac{1}{12}}, \frac{1}{2} + \sqrt{\frac{1}{12}}\right]. \tag{6.20}$$

Taking the minimum value of the min-entropy over this range gives an inference of only about 1.02 bits.

6.1.9 Bayesian Inference

Bayesian inference requires an *a priori* distribution on the probability space Π. For this example, let us assume a uniform distribution on $\Pi = [0, 1]$. For $x = (0, 1, 1)$, recall that the standard likelihood grading was $L_x(p) = p^2(1-p)$, so the Bayesian grading works out to be:

$$B_x(p) = \frac{p^2(1-p)}{\int_0^1 p^2(1-p)dp} = 12p^2(1-p). \tag{6.21}$$

Because this grading is just a constant scaling of the likelihood grading, it gives the same maximally graded inference. It is unclear how to use grading for thresholding.

One can use the Bayesian grading to calculate an average inference over the Π of the inferred min-entropy. Such averaging is probably ill-advised for cryptographic applications, but its computation would be as follows:

$$\int_0^{1/2} -3\log_2(1-p)B_x(p)dp + \int_{1/2}^1 -3\log_2(p)B_x(p)dp \tag{6.22}$$

which seems to be about 1.69 bits.

6.1.10 Working Entropy

The working entropy at 1 and 2 bits of workload are now considered for the $(2,3)$ independent model. Assuming $p \geqslant \frac{1}{2}$, the working entropies are:

$$H_{(w)}(p) = \begin{cases} -2\log_2(p) & \text{if } w = 1 \\ -2\log_2(p) - \log_2(3-2p) & \text{if } w = 2 \end{cases} \tag{6.23}$$

For $p < \frac{1}{2}$, replace p by $1-p$ in the formula above.

6.1.10.1 Maximum Likelihood Estimate

The maximum likelihood inference for p is $\hat{p} = \frac{2}{3}$. Applying (6.23) at workload of 1 bit gives about 1.17 bits of entropy. Applying (6.23) at workload of 2 bits gives about 0.43 bits of entropy.

6.1.10.2 Threshold Inclusive Typicality

At a workload of 1 bit and confidence level c, and using the infimum estimate, the resulting inference is very similar to (6.7), just multiplied by $\frac{2}{3}$, so:

$$H_{(1)}(c) = \begin{cases} \infty & \text{if } c = 0 \\ 2 & \text{if } 0 < c \leqslant \frac{1}{8} \\ -\frac{2}{3}\log_2(c) & \text{if } \frac{1}{8} < c \leqslant 1 \end{cases} \tag{6.24}$$

At a workload of 2 bits, it works out to

$$H_{(2)}(c) = \begin{cases} \infty & \text{if } c = 0 \\ 1 & \text{if } 0 < c \leqslant \frac{1}{8} \\ -\frac{2}{3}\log_2(c) - \log_2(3 - 2\sqrt[3]{c}) & \text{if } \frac{1}{8} < c \leqslant 1 \end{cases} \tag{6.25}$$

6.1.11 Applied Min-Entropy

Suppose that the only information about x that will be applied is $f(x) = x_0 \oplus x_1 \oplus x_2$. The applied model is (Π, Y, Q), with the same probability space $\Pi = [0,1]$ as before, applied sample space $Y = \{0,1\}$, and applied probability function Q, which works out from (3.20) to be

$$Q_p(y) = \begin{cases} (1-p)(1-2p+4p^2) & \text{if } y = 0 \\ p(3-6p+4p^2) & \text{if } y = 1 \end{cases} \tag{6.26}$$

The applied min-entropy, as a function of p is therefore:

$$H_{f(\infty)}(p) = \begin{cases} -\log_2((1-p)(1-2p+4p^2)) & \text{if } 0 \leqslant p \leqslant \frac{1}{2} \\ -\log_2(p(3-6p+4p^2)) & \text{if } \frac{1}{2} \leqslant p \leqslant 1 \end{cases} \tag{6.27}$$

Remark 6.30. The applied min-entropy is strictly less than the min-entropy. At $p = \frac{1}{2}$ the applied min-entropy is one third that of the min-entropy. But as $p \to 1$ or $p \to 0$, the ratio of the min-entropy to the applied min-entropy approaches one.

Remark 6.31. The applied min-entropy of $f(x)$ as function has a plateau around $p = \frac{1}{2}$, whereas the min-entropy x has a sharp peak. So the applied min-entropy, in this example, is is less affected than the min-entropy by slight deviations in p, at least when p is close to a uniform distribution.

6.1.11.1 Maximal Likelihood At the maximal likelihood inference $\hat{p} = \frac{2}{3}$ the applied min-entropy is about $-\log_2(\frac{14}{27}) \approx 0.948$ bits.

6.1.11.2 Inclusive Typicality Taking the infimum of threshold inclusive typicality inference, gives the following:

$$H_{f(\infty)}(c) = \begin{cases} \infty & \text{if } c = 0 \\ 1 & \text{if } 0 < c \leqslant \frac{1}{8} \\ -\log_2(3\sqrt[3]{c} - 6\sqrt[3]{c^2} + 4c) & \text{if } \frac{1}{8} < c \leqslant 1 \end{cases} \tag{6.28}$$

at a confidence of c.

6.1.12 Contingent Min-Entropy

Suppose that the adversary can learn the function $f(x)$ where f is defined as:

$$f(x) = \begin{cases} 0 & \text{if } x \in \{(0,0,0),(1,1,1)\}, \\ 1 & \text{otherwise.} \end{cases} \tag{6.29}$$

Remark 6.32. One reason that an adversary might learn such a function $f(x)$ is that the amount of inferred entropy may depend strongly on the function $f(x)$, and thus the actions that would be taken by cryptographic implementation in an effort to gather enough entropy would need to differ, thereby creating a higher chance of a side channel.

Remark 6.33. Another possible reason that an adversary might learn such a function is that $f(x)$ is the exclusive-or of the bits in the representation of $x_0 + x_1 + x_2$.

For given $p \in \Pi$, the general formula (3.22) for contingent min-entropy works out to be:

$$H_{\infty|f}(p) = \begin{cases} -2\log_2(1-p) & \text{if } 0 \leqslant p \leqslant \frac{1}{2} \\ -2\log_2 p & \text{if } \frac{1}{2} \leqslant p \leqslant 1 \end{cases} \tag{6.30}$$

Remark 6.34. For example, when $p = \frac{1}{2}$, the contingent entropy is 2 bits. Intuitively, this is because an adversary has a strategy to guess x with probability $\frac{1}{4}$.

One such strategy is to guess $x = (0,0,0)$ when $f(x) = 0$ and to guess $x = (0,0,1)$ when $f(x) = 1$.

The first case occurs $\frac{1}{4}$ of the time, and the adversary guess is right $\frac{1}{2}$ of that time, making for a correct guess $\frac{1}{8}$ of the time. The second case occurs $\frac{3}{4}$ of the time and the adversary's guess is right $\frac{1}{6}$ of that time, making for a correct guess $\frac{1}{8}$ of time. These correctly-guessed times are disjoint and total to $\frac{1}{4}$.

The contingent min-entropy in this example works out to always be exactly $\frac{2}{3}$ of the min-entropy. Therefore, all the inferences for min-entropy will scale exactly $\frac{2}{3}$ for inferences of contingent min-entropy.

6.1.13 Filtered Min-Entropy

Suppose that an implementation of a source assumed to be in $(2,3)$ independent model will reject a sample output of $(0,0,0)$ or $(1,1,1)$, perhaps on the ground that these sample could have arisen from a deterministic distribution in the independent model.

Therefore, the adversary wishing to guess x can exclude these values. For cryptographic purposes, one must consider the filtered entropy of x based on the knowledge that the adversary would possess. In other words, the adversary has knowledge that $x \in Y \subsetneq X$ where $Y = \{(0,0,1),(0,1,0),(1,0,0),(0,1,1),(1,0,1),(1,1,0)\}$. Applying (3.26), maximizing, and taking the negative logarithm to the base two, yields a filtered min-entropy value of:

$$\log_2(3) - \log_2(\max(p, 1-p)) \tag{6.31}$$

The inferred set of distribution for p does not depend on the parameter, so to infer the filtered entropy, it will suffice to apply the filtered entropy to the inferred sets, and take infima.

6.1.13.1 Maximum Likelihood Estimate The maximum likelihood inference gives $\hat{p} = \frac{2}{3}$. Applying (6.31) gives an inferred filtered entropy of $2\log_2(3) - 1 \approx 2.17$ bits of filtered entropy.

6.1.13.2 Threshold Inclusive Typicality At confidence level c, the filtered entropy, works out to be (by adapting (6.7)),

$$I_c(x) = \begin{cases} \infty & \text{if } c = 0 \\ \log_2(3) + 1 & \text{if } 0 < c \leqslant \frac{1}{8} \\ \log_2(3) - \frac{1}{3}\log_2(c) & \text{if } \frac{1}{8} < c \leqslant 1 \end{cases} \tag{6.32}$$

Remark 6.35. As confidence levels c approaches 1, the threshold inclusive typicality inferred min-entropy approaches 0, while the threshold inclusive inferred filtered entropy approaches $\log_2 3 \approx 1.58$.

The reason that inferred min-entropy is much lower is that it allows for the possibility of $x = (1, 1, 1)$, which could occur with probability nearly 1 when the confidence level approaches 1. By contrast, the filtered entropy, as defined above, does not allow $x = (1, 1, 1)$, because $(1, 1, 1)$ is filtered.

Remark 6.36. In a cryptographic application, this example is a little artificial. In the case of prospective assessment, as $c \to 1$, the inferred min-entropy approaches 0, because $p \to 1$. A source is likely to result in $(1, 1, 1)$ and therefore by rejected.

So, as noted before, the possibility of rejection is not reflected in the definition of filtered entropy. If the source can be freely sampled until the result is not rejected, then indeed, the higher inferred contingent entropy properly reflects reality.

But in this report, it is generally presumed that the source is expensive to sample. So, a low inferred min-entropy does really reflect something about the rate at which entropy can be drawn from the source.

6.1.14 Sample Entropy

In this section, inferences about the sample-dependent parameter sample entropy (§3.3.1) are made.

Remark 6.37. Recall that sample entropy is mainly useful retrospective inference.

Remark 6.38. Most inference methods given sample $x = (0, 1, 1)$ in the $(2, 3)$ independent model infer a set of distributions whose min-entropy infimum distribution has $p \geqslant \frac{1}{2}$, roughly because 1 appears more often than 0 in the sample x. For such a distribution with $p \geqslant \frac{1}{2}$, the sample value $(1, 1, 1)$ is at least as likely the given sample. Indeed, when $p > \frac{1}{2}$, the sample $(1, 1, 1)$ is more likely, and is the sample which gives rise to the min-entropy of the distribution p.

In these cases, $x = (0, 1, 1)$ has sample entropy at least as high as the min-entropy of p.

Remark 6.39. One usually applies retrospective inference, when one does not wish to waste entropy, so a higher value of sample entropy is not to be discarded.

The inferred set of probability distribution is the same as for min-entropy. The remaining task is to apply the sample entropy parameter, and then take the infimum.

6.1.14.1 Maximal Likelihood Taking the maximum likelihood inference gives $\hat{p} = \frac{2}{3}$ for the solely inferred distribution. The inferred sample entry is $I(\hat{p}, x) = -\log_2(\hat{p}^2(1 - \hat{p})) = 3\log_2(3) - 2 \approx 2.75$.

Remark 6.40. By comparison, the inferred sample entropy 2.75 is exactly 1 bit greater than the inferred min-entropy 1.75, under the same inference method (maximal likelihood estimation).

6.1.14.2 Threshold Inclusive Typicality Each threshold level t determines, under threshold inclusive typicality, a set of inferred distributions $i_{g_1 > t}(x)$, which were calculated in (6.5). Applying the parameter sample entropy

$p \mapsto I(p,x)$ to each of these subsets of Π, seems to gives

$$I(i_{g_1>t}(x),x) = \begin{cases} \varnothing & \text{if } t = 1 \\ \{3\} & \text{if } \frac{7}{8} \leqslant t < 1 \\ \left(-\log_2\left(t-1+(1-t)^{2/3}\right),3\right] & \text{if } \frac{19}{27} \leqslant t < \frac{7}{8} \\ [3\log_2(3)-2,3] & \text{if } \frac{1}{2} \leqslant t < \frac{19}{27} \\ [3\log_2(3)-2,-\log_2(q(t))] & \text{if } 0 \leqslant t < \frac{1}{2} \end{cases} \tag{6.33}$$

where $q(t)$ is the function taking value in interval $[0,\frac{1}{2}]$ such that $3q(t)^2 - 2q(t)^3 = t$.

Remark 6.41. At thresholds below $\frac{1}{2}$, the inferred interval of sample entropies contains values larger than 3, which is an instance of Remark 3.48.

For confidence level $c = 1-t$ the infimum of the inferred sample entropies is:

$$I_c(x) = \begin{cases} 3\log_2(3)-2 & \text{if } c \geqslant \frac{8}{27} \\ -\log_2(-c+c^{2/3}) & \text{if } \frac{1}{8} < c \leqslant \frac{8}{27} \\ 3 & \text{if } 0 < c \leqslant \frac{1}{8} \\ \infty & \text{if } c = 0 \end{cases} \tag{6.34}$$

6.1.15 Eventuated Min-Entropy

Suppose that the probability model is the $(2,5)$ independent model but that, at the time of making inference about p, the first three bits of x have been observed to be $(0,1,1)$. So, in other words, the event E that has occurred is that $(x_0,x_1,x_2) = (0,1,1)$.

The first three bits of the x adhere to the $(2,3)$ independent model, so the inferences to be made about p are those that would be made in $(2,3)$ independent. In particular, the inferences about p of the subsections above apply. In this section, inferences will be made about the sample-dependent parameter eventuated min-entropy (§3.3.2).

The eventuated min-entropy is

$$H_{\infty\|E}(p,x) = -\log_2(1-p)p^2\max(p,1-p)^2 \tag{6.35}$$

6.1.15.1 Maximal Likelihood The maximum likelihood inference for p was $\hat{p} = \frac{2}{3}$ (actually a singleton set), from §6.1.2.

The inferred eventuated min-entropy under maximum likelihood inference is therefore $5\log_2(3)-4 \approx 3.92$ bits.

6.1.15.2 Threshold Inclusive Typicality Applying the infimum value of the eventuated min-entropy parameter $H_{\infty\|E}$ from (6.35) to each of the threshold inclusive typicality inferred sets of distributions from (6.5), and expressing the results as a function of the confidence level $c = 1-t$, gives

$$H_{\infty\|E:g_1>(1-c)}(x) = \begin{cases} \infty & \text{if } c = 0 \\ 5 & \text{if } 0 < c \leqslant \frac{1}{8} \\ -\frac{4}{3}\log_2 c - \log_2(1-\sqrt[3]{c}) & \text{if } \frac{1}{8} < c \leqslant \frac{64}{125} \\ 5\log_2(5)-8 \approx 3.61 & \text{if } \frac{64}{125} \leqslant c \leqslant 1 \end{cases} \tag{6.36}$$

Remark 6.42. The distribution $p = \frac{4}{5}$ actually minimizes the eventuated min-entropy. Consequently at high enough confidence levels, specifically, as shown above, the infimum of the inferred eventuated min-entropies is realized at $p = \frac{4}{5}$.

Recall that the adversary is presumed to know p. When $p > \frac{1}{2}$, the generally optimum single guess at x for adversary is to guess $x = (1,1,1,1,1)$. Because the event E has occurred this optimum strategy will fail, because $x_0 = 0$ in the event E and x_0 in the optimal guess..

Nevertheless, eventuated min-entropy attempts to account for all possible strategies, including a strategy to guess $x = (0,1,1,1,1)$. What eventuated min-entropy measures is the general success rate of such a strategy as if the event E had not occurred.

6.1.16 Applied Eventuated Min-Entropy

Suppose that the probability model is the $(2,5)$ independent model but that, at the time of making inference about p, the first three bits of x have been observed to be $(0,1,1)$. So, in other words, the event E that has occurred is that $(x_0,x_1,x_2)=(0,1,1)$. Explicitly, $E=\{(0,1,1,0,0),(0,1,1,0,1),(0,1,1,1,0),(0,1,1,1,1)\}$. Furthermore, suppose that only the middle three bits $f(x)=(x_1,x_2,x_3)$ are to be used.

The first three bits of the x adhere to the $(2,3)$ independent model, so the inferences to be made about p are those that would be made in $(2,3)$ independent. In particular, the inferences about p of the subsections above apply. In this section, inferences will be made about the sample-dependent parameter applied eventuated min-entropy (§3.3.3).

The applied eventuated min-entropy is

$$H_{f(\infty)\|E}(p,x) = -\log_2 p^2 \max(p,1-p) \tag{6.37}$$

6.1.16.1 Maximal Likelihood The maximum likelihood inference for p was $\hat{p}=\frac{2}{3}$ (actually a singleton set), from §6.1.2. The inferred eventuate min-entropy under maximum likelihood inference is therefore $3\log_2(3)-3\approx 1.75$ bits.

6.1.16.2 Threshold Inclusive Typicality Applying the infimum value of the eventuated min-entropy parameter $H_{f(\infty)\|E}$ from (6.37) to each of the threshold inclusive typicality inferred sets of distributions from (6.5), and expressing the results as a function of the confidence level $c=1-t$, gives the same inferred entropy as the inferred min-entropy from (6.7).

Remark 6.43. For high confidence c, the inferred applied eventuated min-entropy approaches zero in this example, whereas eventuated min-entropy in the previous example did not approach zero. The main difference accounting for this is that here the adversary's ideal strategy (not hinging on event E), knowing $p\geqslant\frac{1}{2}$ is to guess $f(x)=(1,1,1)$, whereas in the previous example, for $p\geqslant\frac{1}{2}$ (not hinging on event E), the adversary's ideal strategy was to guess $(1,1,1,1,1)$.

6.1.17 Contingent Eventuated Min-Entropy

Suppose that the model is the $(2,5)$ independent model. Suppose that, at the time of making an inference, the event E concerning the sample x that $(x_0,x_1,x_2)=(0,1,1)$ is observed. Suppose that the adversary will learn the value of $g(x)$ where g is the function $g:X\to\{0,1,2,3,4,5\}:x\mapsto x_0+x_1+x_2+x_3+x_4$. The contingent eventuated min-entropy from §3.3.4 may be inferred as follows.

A function f supplementary to g that may minimize the applied eventuated min-entropy $H_{f(\infty)\|E)}(p,x)$ is a function $f:X\to\{0,1,2,\ldots,9\}$ such that

$$f:\begin{cases}(0,1,1,0,0) & \mapsto 0\\ (0,1,1,0,1) & \mapsto 0\\ (0,1,1,1,0) & \mapsto 1\\ (0,1,1,1,1) & \mapsto 0\end{cases} \tag{6.38}$$

and such that, for each $j\in\{0,1,2,3,4,5\}$, the function f maps $g^{-1}(j)$ injectively into the set $\{0,1,\ldots,\binom{5}{j}-1\}$. If this function f does indeed minimize the applied min-entropy, then contingent min-entropy is given by

$$H_{\infty|g\|E}(p,x) = -\log_2 p^2(1-p)((1-p)^2+(1-p)p+p^2) \tag{6.39}$$

Contingent eventuated min-entropy seems to have a minimum at

$$p=\frac{8+\sqrt[3]{107+15\sqrt{129}}+\sqrt[3]{107-15\sqrt{129}}}{15}\approx 0.702 \tag{6.40}$$

6.2 Polling Inference

Examples of inference in the $(2, N)$ independent model, with $N \gg 2$ are considered in this section.

As in §6.1, we use the simplified description of the model, in which $\Pi = [0, 1]$, with distribution p mapping to distribution $(1 - p, p)$ in the standard description of the model.

Whereas §6.1, and §6.3 could be considered as low sample size inferences in the independent model, the example in this section could be considered as a large sample size. Intuition suggests the inferences should have higher confidence levels, and that the resulting inference depend less on the inference method.

This example could arise in various ways. Coins could have been flipped, either one coin N times, or N coins once, or something in between. This type of inference also arises in non-cryptographic applications such as in polling: say N people are queried on a yes or no.

It is again emphasized that the independent model is being assumed in this section, not assessed. Again, it is assumed that the N bits are independent and identically distributed. It is under these assumptions that inferences will be made.

Each example will address a distinct inference method. A first part of each example may treat the general case of any N and any sample x. For the sake concreteness, a second part of each example may treat a specific choice of N and x. For consistency of comparison, each example will use the same specific N and x. For ease of computation, the fairly small choice $N = 32$ will be used. For x, we will use:

$$x = (1, 0, 0, 1, 1, 0, 0, 1, 0, 1, 1, 1, 0, 1, 1, 1, 0, 1, 1, 1, 0, 0, 1, 1, 1, 0, 1, 0, 1, 0, 1, 1) \tag{6.41}$$

This value x has 20 entries of value 1, and 12 of value 0. In the independent model, the order of the entries does not matter, so the 20 ones could be have appeared first, following by the zeroes, without affecting the inferences. Of course, if the probability model was a Markov, then the order of the bits is crucially important.

Remark 6.44. The value of x above is derived from the binary expansion of $\sqrt{23}$. Taking into account the specialized manner in which this x was actually selected produces a model much different than the independent model, which would likely yield much lower entropy estimates.

6.2.1 Maximum Likelihood

Recall that likelihood function is $L_x(p) = P_p(x) = (1 - p)^{f(x)_0} p^{f(x)_1}$, where $f(x)$ is the frequency vector §5.4.2 of x. Just as the probability model was given a simplified description, we simplify the frequency vector to a scalar $f(x)$ counting the number of ones in x. (So, the former frequency vector is now $(N - f(x), f(x))$). In this notation,

$$L_x(p) = (1 - p)^{N-f} p^f \tag{6.42}$$

To maximize L_x, calculate its derivative as $L'_x(p) = -(N - f)(1 - p)^{N-f-1} p^f + f(1 - p)^{N-f} p^{f-1} = (f(1 - p) - (N - f)p))(1 - p)^{N-f-1} p^{f-1} = ()$. For $0 < f < N$, the solutions to $L'_x(p) = 0$ are $p = 0$ and $p = \frac{f}{N}$ and $p = 1$. (At $f = 0$ and $f = N$, the solutions are $p = 1$ and $p = 0$ respectively.) Because L_x is differentiable, any local maximum $\hat{p}$ must occur at a critical point with $L_x(\hat{p}) = 0$, or at boundary of $\Pi = [0, 1]$. It is straightforward to confirm, $\hat{p} = \frac{f}{N}$ is the global maximum in $[0, 1]$ for all $0 \leqslant f \leqslant N$. (

Remark 6.45. For the $f \in \{0, N\}$ the global maximum occurs at a boundary point where $L'_x(\hat{p}) \neq 0$. Otherwise the global maximum occurs at a local minimum interior to the domain $[0, 1]$.

With the specific choices of x from (6.41), we get an inferred distribution of $\hat{p} = \frac{20}{23} = \frac{5}{8}$.

The inferred min-entropy is $H_\infty(\hat{p}) = -\log_2 P_{\hat{p}}(\hat{x}) = -\log_2 \hat{p}^{32} \approx 21.7$ bits. This is due the the fact $\hat{p} > \frac{1}{2}$ makes the sample $\hat{x} = (1^{32})$ the most likely sample value.

6.2.2 Inclusive Typicality

Using the notation $f(x)$ from §6.2.1, the inclusive typicality is:

$$\begin{aligned} g_1(x,p) &= \sum_{y:P_p(y)\leqslant P_p(y)} P_p(y) \\ &= \sum_{y:(1-p)^{N-f(y)}p^{f(y)}\leqslant(1-p)^{N-f(x)}p^{f(x)}} (1-p)^{N-f(y)}p^{f(y)} \\ &= \begin{cases} \sum_{e=f(x)}^{N}\binom{N}{e}(1-p)^{N-e}p^e & \text{if } 0\leqslant p<\frac{1}{2} \\ 1 & \text{if } p=\frac{1}{2} \\ \sum_{e=0}^{f(x)}\binom{N}{e}(1-p)^{N-e}p^e & \text{if } \frac{1}{2}<p\leqslant 1 \end{cases} \end{aligned} \tag{6.43}$$

The inclusive typicality at the maximum likelihood estimate $\hat{p}$ is about 0.566. The inclusive typicality as p approaches $\frac{1}{2}$ from above (but not at $\frac{1}{2}$) is about 0.945.

If we want to have confidence level of 0.999, which corresponds to a threshold of 0.001, then the largest value of p meeting this threshold is $p \approx 0.857$. The corresponding infimum inference for the min-entropy of p seems to 7.12 bits of entropy.

Remark 6.46. Generally, as N approaches infinity, if $f(x) > \frac{N}{2}$, then, at all but the most lowest and highest threshold levels, the inferred set of distributions takes the form of an interval $[a,b]$ where $a=\frac{1}{2}$ and $b \approx \frac{f(x)}{N}$. Indeed, in the interval $[a,b]$, the typicality is nearly one, and elsewhere is is nearly zero.

The upper end of the interval corresponds to the maximum likelihood estimate.

The lower bound of the interval reflects the fact that for distributions $p \gtrapprox \frac{1}{2}$, The function P_p of the sample x is sufficiently flat in the sense that with probability near to 1, it holds that $P_p(y) \leqslant P_p(x)$.

Remark 6.47. A more precise description of the approximate shape of the inclusive typicality for large N is given by the Gauss error function.

Remark 6.48. From the perspective of general inference, the inference from inclusive typicality may seem too weak, in that it always infers some distributions close to $\frac{1}{2}$, whereas one might expect that inference should strongly value distributions near to $\frac{f(x)}{N}$. (Sample statistic induced inference may resolve this.)

From the perspective of cryptography, the arguable weakness of the inference makes no difference in this case, because by taking the infimum of the entropies in the interval, we find the infimum is unaffected by the inclusion of distributions near to $\frac{1}{2}$.

6.2.3 Balanced Typicality

As N gets larger, the difference between inclusive and balanced typicality becomes negligible compared to the total typicality.

6.2.4 Adjusted Likelihood

The adjusted likelihood seems to take a maximum value at $\hat{p} \approx 0.5516$. The inferred min-entropy is then about 27.5 bits.

6.2.5 Frequency Statistic Induced Inference

The induced likelihood of frequency is proportional of the standard likelihood in the sense that $L_{f(x)}(p) = \binom{N}{f}L_x(p)$, so taking the induced inference under maximal likelihood is the same, namely $\hat{p} = \frac{f(x)}{N}$.

The frequency induced inclusive typicality takes the form

$$g_1(f,p) = \left(\sum_{e=0}^{a(f,p)} + \sum_{e=b(f,p)}^{N}\right)\binom{N}{e}p^e(1-p)^{N-e} \tag{6.44}$$

where $a(f,p)$ and $b(f,p)$ are integers determined by f and p, because the likelihood function is unimodal, increasing for $f \leqslant pN$ and decreasing for $f \geqslant pN$. So, $a(f,p) = f$ if $f \leqslant pN$ and $b(f,p) = f$ if $f > pN$.

For large N, the probability function takes the shape of a normal curve, due to the Central Limit Theorem. This suggests the approximations $a(f,p) \approx pn - |pn - f|$ and $b(f,p) \approx pn + |pn - f|$.

For the specific $N = 32$ and sample x from (6.41), which has $f(x) = 20$, the function $p \mapsto g_1(f,p)$ from (6.44) was estimated using floating point arithmetic at value $p = \frac{m}{8192}$ for integers m with $0 \leqslant m \leqslant 8192$, and plotted as shown in Figure 5.

Remark 6.49. The non-smooth, stepped appearance of the graph seems to be the actually correct effect of the shifting summation term limits, and is not merely some round effect. As N gets, this curve should probably approach a smoother curve. The shape of Figure 5 might suggest that as N goes to infinity, the curve would approach in shape a normal curve, but actually it should approach in shape the sum of Gauss error function and an reflected Gauss error function. The curve will be smooth except for a sharp peak at the maximum.




Figure 5: Frequency-Induced Inclusive Typicality Plot in the $(2, 32)$ independent model

Remark 6.50. By just casually glancing at Figure 5, at a confidence level of 0.9, the inferred set of distributions seems to be about [0.47,0.75]. Applying the min-entropy parameter gives an inferred set of about [13.28,32]. Taking the infimum of the min-entropies, gives 13.83 bits of entropy at a 90% confidence level.

Remark 6.51. Maximal inclusive typicality should in theory be obtained whenever p gives a peak in the likelihood at frequency value $f = 20$, which should occur when $p \in [\frac{19}{32}, \frac{20}{32}]$. Figure 5 is only slightly off from this.

6.3 Low Sample Sizes in the Independent Model

The hypothetical example from §1.1.2.8 is now addressed under the formal approaches of this report. Recall that the independent probability model was assumed. Specifically, the (m, N) independent with $m = 2^{32}$.

Remark 6.52. Some heuristic justification for the independent model. Muons are elementary particles similar to electrons but much more massive. Because of their large mass, creation of muons requires amounts of localized energy so large that they typically do not arise to nuclear reactions. Thus creation of muons on earth requires accelerators.

Muons passing through the atmosphere arise from the cosmic rays, primarily intergalactic protons that have been accelerated by galactic magnetic fields over very long distances to very high speeds. These protons strike atoms in the atmosphere and create muons. The muons then continue in the nearly the same direction as the original proton, ionizing atoms along the way, until the muons decay into a high-energy electron and neutrinos. Because of the mass, charge, and high speed of cosmic ray muons, they are highly penetrating and can be used to form images of the moon kilometers underground.

Given the above, it seems not unreasonable that each muons passing through a detector may be independent. Especially suggestive of this assumption would the intergalactic source: since perhaps muons from different directions would have sources very far apart within the universe, and ought to have independent speeds.

Of course, hypothesis testing can be applied to this assumption. Possible reasonable causes for lack of independence might be bursts or regularity of muons from a certain directions of the universe.

In our hypothetical example, a third party laboratory is assessing the source, collecting $N = 1024$ muon measures, so the model from the lab's perspective is the $(m, N) = (2^{32}, 2^{10})$ independent model. Because $N \ll m$, the sample size may be deemed as low.

Recall the supposition that the laboratory observes 1023 distinct values among the 1024 muon speed measurements. In other words, one value repeats and all other values are distinct. Because the independent model is assumed, the actual values and the order in which they occurred are irrelevant for inferring entropy. The independent model implies that the assessed entropy is a function of sorted frequencies. So, without loss of generality, it can be assumed that $x = (x_0, x_1, \ldots, x_{N-1}) = (0, 0, 1, \ldots, N-2)$.

Because the independent model is assumed, each muon measurement contributes equally to the entropy. The lab can divide its overall assessment by N to determine the entropy per component. This will determine the amount of entropy per muon measurement.

The lab's observations will not include the sample values used in cryptographic applications. So the entropy assessment will be prospective. In the field where the source is deployed, if the assessed min-entropy per component is h bits, and the goal it is to obtain to k bits of min-entropy, then a value of $N = \lceil k/h \rceil$ can be used.

6.3.1 Maximal Likelihood Estimate

It is verified below that the maximal likelihood estimate inference for the probability distribution p is to take the relative frequencies of from the sample x. More precisely, the maximal likelihood inference for the distribution is the set $\{\hat{p}\}$, where

$$\hat{p}_i = \begin{cases} \frac{2}{N} & \text{if } i = 0 \\ \frac{1}{N} & \text{if } 1 \leqslant i \leqslant N-2 \\ 0 & \text{if } N-1 \leqslant i \leqslant m-1 \end{cases} \tag{6.45}$$

The set-value inference for the min-entropy is then $\{H_\infty(\hat{p})\}$. Narrowing the set-valued inference to a point-valued inference, by taking the minimum, and evaluating the result numerically gives 9216 bits of min-entropy, which is 9 bits of min-entropy per component of x.

Remark 6.53. This estimate is considerably lower than the heuristic argument for about 20 bits in the introduction. On one hand, a lower estimate is more prudent, causing the implementer to seek out more entropy. On the other hand, a low entropy estimate is expensive, because more entropy has to be gathered which can be costly.

The verification mentioned above for given maximum likelihood estimate for independent model is as follows. Apply (A.6) to the objective f

$$f(p) = -L_x(p) = -p_0^2 p_1 p_2 \ldots p_{N-1} \tag{6.46}$$

The gradient of the f is given by

$$\nabla f(p) = (2\Lambda/p_0, \Lambda/p_1, \ldots, \Lambda/p_{N-2}, 0, \ldots, 0), \tag{6.47}$$

where $\Lambda = -L_x(p)$, provided none of $p_0, \ldots, p_{N-2}$ are zero. If any of $p_0, p_1, \ldots, p_{N-2}$ are zero, then the likelihood is zero, and it is easy to find p such that the likelihood is positive. Therefore, we can assume that none of $p_0, \ldots, p_{N-2}$ are zero.

The right hand side of (A.6) can be seen to be simply $N\Lambda$. Therefore the m inequalities of (A.6) becomes, upon multiplication of appropriate denominators and division by $N\Lambda < 0$ become

$$2/N \leqslant p_0 \tag{6.48}$$
$$1/N \leqslant p_1, \ldots, p_{N-2} \tag{6.49}$$
$$0 \geqslant p_{N-1}, \ldots, p_{m-1} \tag{6.50}$$

which, with the usual defining conditions on the probability distribution p, implies the result claimed above.

Remark 6.54. The inclusive typicality of the the sole distribution $\hat{p}$ in the maximum likelihood inferred set has value

$$g_1(x, \hat{p}) = \left(1 - \frac{2}{N}\right)^N \left(5 + \frac{10}{N-2} + \frac{4}{(N-2)^2}\right) \approx 0.68 \tag{6.51}$$

which is less than 1, because other sample values have higher probability than x, in particular, any sample y in which the component 0 appears more than twice, and all other components are at most $N-2$. In particular, the most likely sample is $(0, 0, \ldots, 0)$, and this is the sample value, that an adversary knowing $p = \hat{p}$, should guess. This sample is 2^{N-2} times more probable than the obtained sample x, under the inferred distribution $\hat{p}$.

Remark 6.55. The balanced typicality of the the sole distribution $\hat{p}$ in the maximum likelihood inferred set has value

$$g_1(x, \hat{p}) = \left(1 - \frac{2}{N}\right)^N \left(4 + \frac{4}{N-2} + \frac{2}{(N-2)^2}\right) \approx 0.54 \tag{6.52}$$

which is more than 1/2, indicating that the $\hat{p}$ has higher balanced typicality than any subuniform distribution consistent with x.

6.3.2 Maximal Inclusive Typicality

Inclusive typicality always takes a maximal value of 1. For the given sample x, the inclusive typicality is 1 provided $P_p(x) \geqslant P_p(y)$ for all $y \in X$. We claim that this will be true whenever:

$$p_0 = p_1 = \cdots = p_{N-2} \geqslant p_{N-1}, \ldots, p_m, \tag{6.53}$$

because $P_p(x) = p_0^2 p_1 \ldots p_{N-2}$. To prove this claim, suppose otherwise. This supposition implies $p_i < p_j$ for some i, j with $i \leqslant N-2$. Replace a p_i by a p_j to get a $P_p(y) > P_p(x)$. More precisely, let $y_{i+1} = j$ and let $y_k = x_k$ for $k \neq i+1$.

This set of probability distributions given by (6.53) is more extensive than that given by a maximum likelihood estimate. For example, it includes the (fully) uniform distribution. The directly inferred set of min-entropies is correspondingly extensive. For example, in includes the inference of $\log_2(N)$ bits.

Nevertheless, taking the minimum inferred min-entropy corresponds to the probability distribution in which $p_i = 1/(N-1)$ for $i \in [0, N-2]$ and otherwise $p_i = 0$. For the choice of $N = 2^{10}$, we get an inferred min-entropy $\log_2(2^{10} - 1) \approx 9.9986$ bits of entropy per component of x.

Remark 6.56. This gives an estimate of almost one more bit of entropy than we obtained from maximal likelihood estimate.

Remark 6.57. This estimate is still considerably lower than the heuristic argument for about 20 bits in the introduction.

Remark 6.58. The probability distribution at which the minimum inferred entropy is attained is a subuniform distribution, specifically an $(N-1, m)$-subuniform distribution.

6.3.3 Maximal Balanced Typicality

Consider distributions p with inclusive typicality of 1, that also approach the uniform distribution X. The balanced typicality of these distributions approaches:

$$1 - \frac{1}{2}\left(\frac{N-1}{m}\right)^N \tag{6.54}$$

or about $1 - 2^{22000}$, which is very close to 1. If these distributions have higher balanced typicality than any others, then the maximal balanced typicality is a limit, with no actual distribution hitting the maximum. Nevertheless, the limit of the distributions exists and is the uniform distribution, which gives an estimate of 32 bits of min-entropy per component.

Remark 6.59. At this point, the assessments seem too pessimistic or too optimistic compared to the intuition from the introduction. Indeed, the introduction informally makes use of a sample statistic.

6.3.4 Frequency Statistic Induced Inference

The function $s : X \to Y$ given by the frequency statistic defined in §5.4.2 induces probability model (Π, Y, Q) such that

$$Q_p(y) = M(y)P_p(x) \tag{6.55}$$

where $x \in X$ is such that $s(x) = y$ and $M(y)$ is an integer multiplier counting that the number of x such that $s(x) = y$. This holds from (5.1) because that statistic s has the property that for all $x, x' \in X$ and $p \in \Pi$ if $s(x) = s(x')$ then $P_p(x) = P_p(x')$.

With our specific sample gathered of $x = (0, 0, 1, \ldots, N-2)$, we have $y = s(x) = (2, 1, \ldots, 1, 0, \ldots, 0)$ where there are $N-2$ entries with value 1 and $m - N + 1$ entries with value 0.

The general formula for the integer multiple $M(y)$ is given by $M(y) = \frac{N!}{y_0!y_1!\ldots y_m!}$. The general formula for the probability of y is therefore:

$$N! \prod_{i=0}^{m} \frac{p_i^{y_i}}{y_i!} \tag{6.56}$$

With our specific sample example gathered y, the value of the multiplier is thus $N!/2$ and the probability is $\frac{1}{2}N!p_0^2 p_1 \ldots p_{N-2}$.

6.3.4.1 Induced Inclusive Typicality Consider the distribution $\hat{p}$ maximal likelihood inference from (6.45). The frequency-induced inclusive typicality of $y = s(x)$ at distribution $\hat{p}$ is one. Therefore the maximal induced inclusive typicality consists of all distributions p reaching induced typicality of one. Taking the infimum of min-entropies over this set will be at most $H_\infty(\hat{p})$, which as above, is 9 bits per component.

Similarly, any threshold graded inference with the frequency-induced inclusive typicality will give, once one takes an infimum of min-entropies, will give at most 9 bits of min-entropy per component of the sample.

6.3.4.2 Induced Balanced Typicality Let u be an integer with $N \leqslant u \leqslant m$. Let S be a $(u-2)$-element subset of $\{1, \ldots, m-2\}$, such that $\{1, \ldots, N-2\} \subseteq S$. Let p^S be a distribution defined by

$$p_i^S = \begin{cases} 2/u & \text{if } i = 0 \\ 1/u & \text{if } i \in S \\ 0 & \text{if } i \notin S \end{cases} \tag{6.57}$$

The frequency-induced inclusive typicality of x at p^S is one, and as such, the distributions p^S would seem to good candidates for maximizing balanced typicality. The frequency-induced balanced typicality of x at p^S seems to be:

$$1 - \frac{1}{4}\binom{m-N+1}{u-N}\frac{N!}{u^N} \tag{6.58}$$

This seems to be maximized when $u = N$. When $u = N$, the distribution p^S is the same as distribution in the maximal likelihood inference.

This suggests that threshold frequency-induced balanced typicality inferences would be similar to threshold frequency-induced inclusive typicality inferences.

6.3.4.3 Induced Threshold Adjusted Likelihood Letting v run over the possible frequency vectors, the adjustment term of adjusted likelihood is

$$\begin{aligned}\sum_v Q_p(v)^2 &= \sum_v \binom{N}{v}^2 p^{2v} \\ &= N!^2[u^N] \sum_v \prod_{i=0}^{m-1} \frac{p_i^{2v_i} u^{v_i}}{v_i!^2} \\ &= N!^2[u^N] \prod_{i=0}^{m-1} \sum_{v_i \geqslant 0} \frac{p_i^{2v_i} u^{v_i}}{v_i!^2} \\ &= N!^2[u^N] \prod_{i=0}^{m-1} C_0(p_i^2 u)\end{aligned} \tag{6.59}$$

where $[u^N]F$ means the coefficient of u^N in the power series F, and C_0 is the Clifford-Bessel function of order 0.

To be completed.

6.3.5 Partition Statistic Induced Inference

Recall (5.8) which states that that partition statistic induces a probability function given by

$$Q_p(\theta) = \binom{N}{\theta} m_\theta(p) \tag{6.60}$$

For our sample x, where $\theta = \phi(x) = (2, 1^{1023})$, where the entry 1 is repeated 1023 times. It follows that $\binom{N}{\theta} = \frac{1024!}{2}$.

6.3.5.1 Maximal Induced Likelihood Before considering the general problem of optimizing the likelihood function $L_x(p) = Q_p(\theta)$ over the whole independent model, consider the more restriction model which considers of only subuniform distributions. This restriction is a relaxation of the model from §1.1.2.8 which contains three subuniform distributions, in which the probability vector p has supports of sizes 2^{10}, 2^{20} and 2^{30}.

Let $p^{(u)}$ be a probability vector, in the (m, N) independent model, that has u entries of $\frac{1}{u}$ and all other entries zero. It results in a subuniform distribution on X where the sample with non-zero probability have probability $\frac{1}{u^N}$. With this notation and $(m, N) = (2^{32}, 2^{10})$, the induced likelihood is

$$L_x(p^{(u)}) = \frac{1024!}{2} 1023 \frac{\binom{u}{1023}}{u^{1024}}, \tag{6.61}$$

with a factor of 1023 accounting for the choice of which value is repeated, and the factor of $\binom{u}{1023}$ account for which of the u entries with non-zero probabilities (as individual entries) appear in the sample.

Based on the assumption that this is the a unimodal function of u, some brute force numerical calculations seem to give $u = 2^{19} - 853$ as the value which maximizes the induced likelihood. This is in close agreement with the inference made in §1.1.2.8.

6.4 Toy Examples in the Markov Model

In this section, two toy examples in the Markov model will be considered:

- The first example uses a $(2,3)$-Markov model with sample value $x = (0,1,1)$. The sample space and sample value are the same as in §6.1. The model is a relaxation of the model in §6.1. Relaxation of the model generally the effect of reducing the infimum of inferred entropy. Indeed, this sample value of x is the output of a deterministic distribution in the Markov model.
- The second example uses a $(2,5)$-Markov model with a sample value $x' = (0,1,1,0,1)$.

6.4.1 Maximum Likelihood Estimate

In the first example, the likelihood function, for $x = (0,1,1)$ in the Markov model is

$$L_x(p) = v_0 M_{0,1} M_{1,1} \tag{6.62}$$

where, recall $p = (v, M)$ is a pair of a vector and a matrix. It is fairly easy to see that L_x is optimized at

$$\hat{p} = \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 1 \end{pmatrix} \right) \tag{6.63}$$

The probability of $P_{\hat{p}}$ is maximized at $y = (0,1,1)$, with value 1. So the point-valued inferred min-entropy is $H_\infty(\hat{p}) = 0$.

In the second example, the likelihood function, for $x' = (0,1,1,0,1)$ in the Markov model is

$$L_{x'}(p) = v_0 M_{0,1}^2 M_{1,1} M_{1,0} \tag{6.64}$$

where, recall $p = (v, M)$ is a pair of a vector and a matrix. It is fairly easy to see that $L_{x'}$ is optimized at

$$\hat{p}' = \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ \frac{1}{2} & \frac{1}{2} \end{pmatrix} \right) \tag{6.65}$$

The probability of $P_{\hat{p}'}$ is maximized at $x' = (0,1,1,0,1)$, with value $\frac{1}{4}$. So the point-valued inference for min-entropy is $H_\infty(\hat{p}) = 2$ bits.

Remark 6.60. At distribution $\hat{p}'$: probability $\frac{1}{4}$ is assigned to $(0,1,0,1,0)$, $(0,1,0,1,1)$ and $(0,1,1,0,1)$; probability $\frac{1}{8}$ is assigned to $(0,1,1,1,0)$ and $(0,1,1,1,1)$; and probability 0 is assigned to all other sample values.

Remark 6.61. The balanced typicality of x' at $\hat{p}'$ is $\frac{5}{8}$.

Remark 6.62. The working entropy at a work load two bit of the distribution $\hat{p}'$ is about 0.19 bits.

6.4.2 Inclusive Typicality

In the case of sample $x = (0,1,1)$, the inclusive typicality at the maximum likelihood distribution $\hat{p}$ is $g_1(x, \hat{p}) = 1$. So, the maximally graded or threshold graded inference based on inclusive typicality will include $\hat{p}$ in the set of distributions. Taking, the infimum of min-entropy over the set of inferred distributions, given an inference of zero for the min-entropy.

6.4.2.1 Maximally Graded In the case of the sample $x' = (0,1,1,0,1)$, the inclusive typicality of the maximal likelihood estimate $\hat{p}'$ from the previous section is 1.

So, the inferred set of distributions from taking the maximally graded inference with grading equal to inclusive typicality is all those the distributions with inclusive typicality equal to one. Since this includes the distribution, this is at most 2 bits.

Some numerical exploration suggests that 2 is indeed the minimum value of the min-entropy among the distribution with inclusive typicality one.

Remark 6.63. The set of distributions with inclusive typicality at x' seems, based on numerical computations, largely characterized as follows: $M_{0,1} \geqslant \frac{1}{2}$; and $\frac{1}{2} M_{1,1} \geqslant c(M_{0,1})$ where c is some concave increasing function with $c(1) \approx 0.618$; and $0 \leqslant v_1 \leqslant M_{1,1}$.

6.4.2.2 Threshold Graded At a confidence level $c = 0.99$, meaning a threshold of $t = 0.01$, some numerical calculations give the distribution $p = (v, M)$ with:

$$v = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad M = \begin{pmatrix} 0 & 1 \\ 0.003345 & 0.996655 \end{pmatrix} \tag{6.66}$$

give rise to an inclusive typicality $g_1(x', p) \approx 0.01$. The min-entropy of this distribution is $H_\infty(p) \approx 0.0145$ bits. So, at a confidence level of about 99%, the inferred min-entropy is at least 0.015 bits.

6.5 Dice

This section illustrates statistical inference about dice rolls. Some inference will be done in various models, and some hypothesis testing on the models themselves will be done.

Two separate processes were used to generate two sample vectors of the following dice rolls:

$$x' = (5, 4, 3, 2, 2, 1, 3, 6, 2, 3, 1, 1, 5, 4, 1, 5, 2, 6, 6, 1, 6, 5, 5, 5) \tag{6.67}$$

$$x = (2, 5, 6, 1, 2, 5, 2, 5, 1, 1, 1, 1, 1, 4, 1, 2, 2, 1, 3, 3, 3, 1, 3, 1) \tag{6.68}$$

Both sample vectors were produced by the author dropping a 15mm die (a cube), with embossed numbers $\{1, 2, 3, 4, 5, 6\}$ into a cup. The die was placed so that it touched the inside of the cup, with the top of the die approximately level with the rim of the cup, with the numeral 1 oriented with its top pointing to the center of the cup. The die was held so, and then let go, with an effort to let the die have initial velocity zero, and thereby let gravity create motion. Despite this effort, the motion of the die did seem to have some correlation with the motion of the fingers releasing. The die then fell to the bottom of the cup, bouncing, rotating, and eventually stabilizing. The numeral facing up was recorded as above.

Sample vector x' is the result of 24 consecutive drops into a cup of height 113mm. Sample vector x is the results of 24 drops into a cup of height 45mm.

6.5.1 The Uniform Model

A commonly assumed model for a a single die roll is the uniform model. It is also commonly assumed that multiple rolls are independent and identically distributed. Combining these two assumptions gives results in the uniform model on the sample space $\{1, 2, 3, 4, 5, 6\}^N$ where N is the number of die rolls.

6.5.1.1 Entropy Assessment in the Uniform Model In the uniform model, the single distribution has min-entropy of $24 \log_2(6) \approx 62.0$ bits for each of x and x'. This assessment assumes the uniform model, which as will be shown below, is not very realistic for the sample x.

6.5.1.2 Hypothesis Testing of the Uniform Model A casual inspection of x from (6.68) should suggest that x does adhere well to the uniform model. Incidentally, observations made during the process used to generate x indicated some correlation between the motion of the die and release motion of the fingers.

Formally, we can apply hypothesis testing to the assumption of the uniform model. Although hypothesis testing is not the main topic of this report, a brief foray into hypothesis testing may be illustrative.

One cannot rule out x as being atypical if we limit ourselves to the uniform model, because any sample value is equally likely. Similarly, the inclusive typicality of all x is 1, and the balanced typicality is $\frac{1}{2}$. The tying effect of uniform model is in effect.

Sample statistics can be used as tiebreakers. Generally, this report has somewhat discouraged the use of sample statistics, at least for entropy assessment, and instead encourages the relaxation of the probability model. More precisely, in entropy assessment, the probability model is deemed well-founded, so the sample statistics should be only used for tiebreaking in the case that the sample statistics is very consistent with the assumed model, for example, by being model-neutral.

For hypothesis testing, the probability model is less trusted, but nevertheless, the general idea above for entropy assessment can be used. One could consider a relaxation of the model, do statistical inference in the alternative model.

If the sample has significantly higher typicality than in the hypothesized model, one can reject the hypothesized model, and favor the alternative model.

Alas, in this case, even the approach of an alternative hypothesis above, is ineffective, because the inclusive typicality was 1 in the hypothesized uniform model, so the alternative cannot have higher typicality. One can blame inclusive typicality and use balanced typicality. But even with balanced typicality, the uniform model gives $\frac{1}{2}$, which is very plausible, and not real grounds for rejection. The approach of comparison to inference over the alternative model does not seem to work well.

An intermediate approach is to use a sample statistic appropriate for the alternative model, such as a model-neutral one, and then compute the induced typicality of the sample in the hypothesize model. This intermediate approach seems to have to address the concerns above in the best possible way.

So, in the specific example at hand, the hypothesized model is the uniform model. The alternative model will be independent model. The sample statistic will be the frequency statistic, which is model-neutral in the independent model.

The frequency-induced inclusive typicality for x' is about 0.81 and for x is about 0.02.

These typicality values can be interpreted as follows: x' can perhaps be considered as highly consistent with the uniform model. Of course, it is probably always more conservative to consider a more relaxed model. So, given x', our confidence in the assumption of the uniform model is not decreased. That is, whatever confidence we had in the assumption of uniform model is the confidence that we could have, as cryptographers, in x having arisen from a uniform distribution.

The other sample x has lower typicality, only 0.02. This alone may not be grounds for rejection of the uniform model, because if the uniform model was correct, one would still get such a result have 2% of the time. In an entropy assessment context, rejection is somewhat wasteful. So, the low typicality 0.02 should be taken as strong incentive for relaxing to the alternative model.

6.5.2 The Independent Model

6.5.2.1 Entropy Assessment in the Independent Model The maximum likelihood inference of min-entropy in the independent model is exactly 48 bits for x' and about 30.3 bits for x.

Remark 6.64. The values are lower than the inference in the uniform model, as expected because the probability model has been relaxed.

Remark 6.65. In the case of x, the maximum likelihood estimate means that inferred distribution takes its maximum probability at $\hat{x} = (1, 1, \ldots, 1)$, the all ones sample vector, and that this probability is about $2^{-30.3}$. The sample $\hat{x}$ is the best guess an adversary can make given the distribution.

Remark 6.66. The inferred sample entropy, under the maximum likelihood distribution, of x is about 52.4 bits.

Other types of inference methods as applied to the independent model have been illustrated in other parts of this report, so will not be illustrated again for this example.

Remark 6.67. If one applies a uniformity extractor to x, assuming the independent model, one can derive an integer y uniformly distributed between 1 and $\frac{24!}{10!5!4!1!3!1!} \approx 2^{43}$. Note that this should not be compared to the inferred min-entropy but rather to the inferred sample entropy of x.

To make prospective inference about some uniformity extractor as the applied function, all that is needed is a precise description of the uniformity extractor function.

6.5.2.2 Hypothesis Testing of the Independent Model Intuition may suggest that the long subsequence of 1 entries in x means the independent model is not an accurate assumption for x. In particular, the entry 1 is is more frequently followed by another 1 than by something else, so order seems to matter, whereas in the independent model order does not matter. In this section, we attempt to formally quantify this intuition, adhering to the general principles of this report.

So the approach from §6.5.1.2 will be followed again. The hypothesized model is the independent model. The alternative model is the Markov model, which is chosen as a simple relaxation of the independent model in which the

order plays a role. So, the idea is to apply a compute the maximal typicality of x as induced by a sample statistic that is model-neutral in the Markov model. The maximality is taken over all distributions in the independent model. If the maximal typicality is low, then it is formally justified to reject the independent model.

Before, embarking on this task, we can see what happens when we compute the of maximal typicality x using no sample statistic, and with a the frequency statistic which is model-neutral in the independent model. Let us use inclusive typicality because it is larger than balanced typicality, so a low value of inclusive typicality is a stronger reason for rejection. Because the uniform distribution u is included the independent model, and inclusive typicality of x at u is 1, the maximal inclusive typicality is 1.

The frequency statistic for x is $y = f(x) = (10, 5, 4, 1, 3, 1)$. The induced probability for any frequency statistic $z = (z_1, \ldots, z_6)$ under distribution $p = (p_1, \ldots, p_6)$ is

$$N! \prod_{j=1}^{6} \frac{p_j^{z_j}}{z_j!} \tag{6.69}$$

Numerical computation of the induced probability at the maximal likelihood distribution $p = \frac{y}{24}$ for each possible frequency vector z, show that the induced probability is uniquely maximized at $z = y$. Therefore, the inclusive typicality at p is exactly one. Therefore, the maximal frequency-induced inclusive typicality is one.

For the sample statistic, use Markov frequency statistic in the alternative (Markov) model from §5.5.1, which is model-neutral in the Markov model. So, we should compute the statistic-induced inclusive typicality for both observed samples x' and x. More precisely, we should compute the maximum value of the inclusive typicality, taking the maximum over all distributions in the independent model. If it is low, then we should prefer the Markov model over the independent model.

First, we note that we will be indexing vector matrix entries from 1 to 6, rather than from 0 to 5, as in earlier sections of this report. The resulting Markov frequency statistic value at our observed sample vectors are:

$$F(x') = \left(\begin{pmatrix} 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 1 & 0 & 2 & 1 \\ 1 & 1 & 1 & 0 & 0 & 1 \\ 1 & 1 & 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 2 & 2 & 0 \\ 1 & 1 & 0 & 0 & 1 & 1 \end{pmatrix} \right) \tag{6.70}$$

and

$$F(x) = \left(\begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 4 & 2 & 2 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 & 3 & 0 \\ 2 & 0 & 2 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \right) \tag{6.71}$$

The induced probabilities can be computed using (5.12) and (5.17). For example, the induced probability of of $F(x')$ at the maximum likelihood distribution as given in (6.72) is about 2.495×10^{-6}. The induced probability of $F(x')$ under the maximum likelihood distribution in the independent model is about 2.086×10^{-12}. The induced probability of $F(x')$ under the uniform distribution is about 5.689×10^{-13}. (Low values of induced probabilities are particular distributions (or maximized over all distributions) are not grounds for model rejection.)

Similarly, the induced probability of $F(x)$ at the maximum likelihood distribution in the Markov model, as given by (6.73), is about 1.723×10^{-4}. The induced probability of $F(x)$ under the maximum likelihood distribution in the independent model is about 1.322×10^{-10}. The induced probability of $F(x)$ under the uniform distribution is 1.723×10^{-13}.

Induced typicalities are computed by summing the induced probabilities over the set of values of the sample statistic. Maximal induced typicality are then computed by determining the maximum over all distribution. For the hypothesis testing task at hand, the space of distributions over which maximum typicality is calculated is the probability space of the hypothesized independent model.

There at most $6\binom{36+23-1}{23} \approx 5.3 \times 10^{16} \approx 2^{55.6}$ values for the frequency statistic (e, U), because the entries of matrix U are non-negative integers summing to 23. This number is smaller than the number of values for x, which is $6^{24} \approx 4.7 \times 10^{18} \approx 2^{62}$, but it is still too large for any currently practical calculation. The b may condition from (5.17) may help somewhat to reduce this number, but perhaps it may not reduce the number to a practical value over which sums can be computed.

A general method to probabilistically estimate the inclusive typicality at a given distribution can be given based on the fact the induced inclusive typicality at $y = F(x)$ is the expected value of the random variable $\gamma(Q_p(F(x)) - Q_p(F(x')))$ where Q_p is the induced probability function, which we can compute, and x' is drawn randomly according to the distribution p, x is fixed, the γ evaluates to one if its input is non-negative, and to zero otherwise. So, based on this expectation, one can compute the random variable for a large number of x' drawn from p, and take the average.

Recall that, generally, we wish to avoid probabilistic algorithms, because the underlying problem of entropy assessment involves inferring probabilities, and thus probabilistic assessment presents a logical circularity. In this case though, a direct calculation was deemed infeasible. One way overcome the circularity is to use a a second source of to assess a given source. This may be useful in the context of unconstrained, system-wide, pre-deployment assessment of sources, but may be much more difficult in retrospective, mid-deployment assessment of sources. Another way to overcome the circularity is to use deterministic pseudorandom generators, such as those based on cryptographic hash function (which are likely to already be available in a cryptographic implementation).

Another potential disadvantage of probabilistic methods is the difficult of maximization of functions that can only be computed probabilistically.

Using this method with 8192 random samples, and just an ordinary pseudorandom number generator, gives an estimate of around 0.62 for the inclusive typicality of $F(x')$ at the distribution which is the maximum likelihood in the independent model. Therefore, the maximal induced inclusive typicality is at least around 0.62. The independent model cannot be rejected from the case because this typicality is too high.

Similarly, for $F(x)$, the estimated induced inclusive is typicality is around 0.38, so the independent model would be not be rejected by this test. In words, the intuition that x has too many successive ones for the independent model is not quantifiably justifiable according to the Markov frequency sample statistic.

Just for comparison, the suppose that x'' consisted of twelve ones followed by twelve twos. The Markov-frequency induced inclusive typicality was estimate for four distributions: the maximum likelihood estimate in the independent model $p = (\frac{1}{2}, \frac{1}{2}, 0, 0, 0, 0)$, at the uniform distribution, and also at the maximum likelihood distribution in the independent model for the sample x' and x. The first three estimated typicalities were zero (within the precision of the numerical computations), but the last distribution resulted in an estimate of around 0.027. The maximal typicality could be much larger, but if it is not, then the independent model could be rejected upon observing the sample x''.

6.5.3 The Markov model

Maximum likelihood estimate in the Markov model gives inferences:

$$\hat{p}(x') = \left(\begin{pmatrix} 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 1/5 & 0 & 1/5 & 0 & 2/5 & 1/5 \\ 1/4 & 1/4 & 1/4 & 0 & 0 & 1/4 \\ 1/3 & 1/3 & 0 & 0 & 0 & 1/3 \\ 1/2 & 0 & 1/2 & 0 & 0 & 0 \\ 0 & 1/5 & 0 & 2/5 & 2/5 & 0 \\ 1/4 & 1/4 & 0 & 0 & 1/4 & 1/4 \end{pmatrix} \right) \tag{6.72}$$

and

$$\hat{p}(x) = \left(\begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 4/9 & 2/9 & 2/9 & 1/9 & 0 & 0 \\ 1/5 & 1/5 & 0 & 0 & 3/5 & 0 \\ 1/2 & 0 & 1/2 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 1/3 & 1/3 & 0 & 0 & 0 & 1/3 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \right) \tag{6.73}$$

The resulting inferred min-entropies are:

$$H_\infty(\hat{p}(x')) \approx 27.8, \tag{6.74}$$
$$H_\infty(\hat{p}(x)) \approx 23.5. \tag{6.75}$$

6.6 Toy Model for a Ring Oscillator

In this section, we consider a toy model for a ring oscillator. No assertion is being made on the accuracy or appropriateness of this model for actual ring oscillators. The purpose of this section is to illustrate of the principles of this report on other types of model.

We begin with a continuous probability model (Π, X_T, P_T). The probability space is

$$\Pi = \{(a,b) : 0 \leqslant a < b\} \tag{6.76}$$

where a and b are defined a interval in the real line. The sample space is

$$X_T = \mathbb{R}^+ = \{t : 0 \leqslant t\} \tag{6.77}$$

where t is real number. (The value t which will be represent the time period of the oscillator). The probability function is now replaced by a probability density function, which is

$$P_{(a,b)}(x) = \begin{cases} \frac{1}{b-a} & \text{if } a \leqslant x \leqslant b \\ 0 & \text{otherwise.} \end{cases} \tag{6.78}$$

An applied model will be used in the cryptographic application. Let the function

$$f : \mathbb{R}^+ \to X = \{0,1\}^N \tag{6.79}$$

be defined by

$$f : t \mapsto (x_0, x_1, x_2, \ldots, x_{N-1}) \tag{6.80}$$

where

$$x_i = \left\lfloor \frac{i}{t} \right\rfloor \bmod 2 \tag{6.81}$$

The idea is that the ring oscillator alternates values between 0 and 1 every t units of time. Every single unit of time, the state of the ring oscillator is sampled and an entry of x is recorded.

Remark 6.68. The model described above is actually a hull model, as in §2.4.4, obtained from tow models. The first model has probability space $\Pi' = [0, \infty)$, with distributions t, with each distribution on the discrete sample space X being deterministic as given by (6.81). The second model has probability space Π from (6.76), and continuous sample space Π'.

Remark 6.69. For $t \in \mathbb{R}^+$ and $x = f(t)$, it is true that $x_0 = 0$. Therefore, any $x \in \{0,1\}$ with $x_0 = 1$ is non-occurring in the applied probability model.

In the toy model above, each distribution gives a uniform continuous distribution on the value of t within some interval $[a,b]$. After applying f, the distribution gives rise to some distribution on the sequences. Because we are starting from a continuous distribution, we cannot directly calculate the applied probabilities from (3.20), which assumed discrete initial distribution. Rather, we need a continuous variant of (3.20). To this end, for $x \in \{0,1\}^N$, let function $\tau_x : \mathbb{R}^+ \to \{0,1\}$ be

$$\tau_x(t) = \begin{cases} 1 & \text{if } f(t) = x, \\ 0 & \text{if } f(t) \neq x. \end{cases} \tag{6.82}$$

Then the applied model is (Π, X, P) where probability function is given by

$$P_{(a,b)}(x) = \frac{\int_a^b \tau_x(t)dt}{b-a} \tag{6.83}$$

The equation above implies that

$$P_{(a,b)}(x) = \frac{1}{b-a}\left|[a,b] \cap \left(\bigcap_{i=1}^{N-1}\bigcup_{m=0}^{\infty}\left[\frac{2m+x_i}{i}, \frac{2m+x_i+1}{i}\right]\right)^{-1}\right| \tag{6.84}$$

where for a subset $S \subset \mathbb{R}^+$, the S^{-1} indicates the set $\{s^{-1} : s \in S\}$ and the $|S|$ indicates the total length of a set. For example if $x = (x_0, x_1, x_2) = (0, 1, 1)$, then

$$P_{(a,b)}(x) = \frac{1}{b-a}\left|[a,b] \cap \left(\left[\frac{1}{2}, \frac{2}{3}\right] \cup \left[\frac{1}{4}, \frac{2}{7}\right] \cup \left[\frac{1}{6}, \frac{2}{11}\right] \cup \left[\frac{1}{8}, \frac{2}{15}\right] \cup \dots\right)\right|. \tag{6.85}$$

By taking $p = [a, b]$ as a sub-interval of one of the connected components of the set $T_x = \{t : f(t) = x\}$, the finds that $P_p(x) = 1$. So, this model is pseudodeterministic.

From a cryptographic standpoint, any reasonable inference should include all the deterministic distributions consistent with the observed sample. Taking prudent principle of using the infimum of the inferred set of entropies should generally give a result of zero, because deterministic distributions will all have entropy of zero. So, any prudent inference in this model gives an entropy assessment of zero.

Although the original model (without the standard deviation restriction above) does not provide any hope for prudent entropy assessment, it, like any model, can be subject to hypothesis testing. Given a sample x, one can calculate its typicalities in the model. Remark 6.69 has shown the model has non-occurring sample values, such as x with $x_0 = 1$. If the sample is a non-occurring value then its typicality, under any distribution, is zero. This should lead to rejection of the model. But for any occurring sample value, a deterministic distribution exists, and therefore the model cannot be rejected on this basis.

Remark 6.70. In the cryptographic context, one has incentive to reject this model, so the cryptographer would seek an alternative model, that some support in terms of other evidence, such as further testing, such as good typicality under extensive sampling, and so.

Under hypothesis testing, the non-occurring x are those for which $\{t : f(t) = x\} = \varnothing$.

There are at most $N^2 - N + 1$ values of x for which x is occurring. To see that, consider $u = t^{-1} \bmod 2$, that is, $t^{-1} - 2\lfloor\frac{1}{2t}\rfloor$. Then $f(t) = f(u^{-1})$, so only the value of u affects the value of f. The value of each x_i changes as a function of u at most $2i$ times as u ranges from 0 to 2. Therefore, there are at most $2(1 + \dots + (N-1)$ transitions in the value of $f(u^{-1})$ as u ranges from 0 to 2.

For large N, the proportion of the space X that is occurring is small. If the hypothesized model is false, then perhaps that some under the true model, the probability that of obtaining an x that is occurring in the hypothesized model becomes small, at least for large N. Once a non-occurring sample x is observed, our toy hypothesized model can be rejected.

If our toy model is rejected, one could move try to move a relaxation of the model, and hopefully one that is not pseudodeterministic.

If the model cannot be rejected, then one's only hope is find a restriction of the model, such as the one described above. Again, to support the restriction, one would have to do some hypothesis testing on the restricted model. But even if the restricted model is supportable, the resulting entropy will always be low, because any restricted model can take at most $N^2 - N + 1$ values, which bounds the entropy to about $-2\log_2(N)$ bits, which is can considerably smaller than the N bits in the representation sample value, and more important than the N units of time needed to generate the sample value. For small enough N, then it might still be worthwhile.

Perhaps the model can salvaged by restricting it, such as by supposing that the random variable t associated with each distribution has some minimum standard deviation. In a real world example, there would have to some justification for adding such a restriction. This would force in each $p = (a, b)$ to be such that $b \geqslant (1 + \epsilon)a$ for some fixed $\epsilon > 0$.

For example, taking $\epsilon = \frac{1}{2}$, then the distribution $p = (\frac{4}{9}, \frac{2}{3})$ is still allowed in the restricted model and we have:

$$P_p(0, 1, 1) = \frac{3}{4}. \tag{6.86}$$

If this is the unique most likely distribution for $x = (0, 1, 1)$, the maximum likelihood estimate would give an infimum inference of min-entropy of about 0.415 bits.

Another approach to salvaging this toy model would to be strengthen by taking its common product with itself (a common power). Such a product model may have a some justification. If ring oscillators are manufactured according to some strict process, then each oscillators should have a rate independent of the others, and the rates should be identically distributed. (Perhaps a natural model for the common distribution of the rates would be a bell curve such as a positive valued version of an normal curve, but our toy model distributions of interval may serve as a decent approximation to such a bell curve.)

Assuming this common model, one might take multiple readings over several of the rings oscillators, and try to infer the probability distribution from the sample values, and then deduce the various entropy parameters.

6.7 Models Based on Poisson Processes

The Poisson and Poisson process models were defined in §2.5.3.1. In a Poisson distribution p, a value of x that maximizes $P_p(x)$ is $x = \lceil p - 1 \rceil$. The min-entropy of the the Poisson distribution p is therefore

$$H_\infty(p) = -\log_2 \left(\frac{e^{-p} p^{\lceil p-1 \rceil}}{\lceil p - 1 \rceil !} \right), \tag{6.87}$$

which can, for large enough p, be approximated using Stirling's formula as

$$H_\infty(p) \approx \tfrac{1}{2} \log_2(p) + 0.92 \tag{6.88}$$

When a Poisson process model is assumed for a source, then one likely has a time-interval $[a, b]$ in which one can access the source. If the time source is sufficiently long compared to the distribution q, then $p = q(b - a)$ is large enough to use the approximation (6.88) to estimate the min-entropy of the cardinality of $x \cap [a, b]$.

Instead one could divide the interval into two pieces, say $\left[a, \frac{a+b}{2}\right]$ and $\left[\frac{a+b}{2}, b\right]$, and consider the min-entropy of the counts for each interval. If the approximation above still applies, then the resulting estimate for the min-entropy is $\log_2(p) + 0.84$, which is about twice as much entropy. One can divide the intervals again, perhaps about doubling the min-entropy, but eventually the approximation (6.88) will no longer apply as p gets too small.

Even though the doubling approximation cannot be applied indefinitely, infinite min-entropy may seem theoretically available if p is sufficiently large, because if at least one real number r is expected in the set $x \cap [a, b]$ it contains an infinite amount of precision, and therefore contains an infinite amount of information. However, min-entropy is not formally defined for continuous distributions. In practice, the real numbers in $x \cap [a, b]$ can only be determined to a finite precision, which limits the min-entropy to a finite amount. As shown below, there remains an upper limit on the min-entropy even if arbitrary precision is available.

Suppose that a source adheres to a Poisson process distribution q, and that a cryptographic implementation can measure down to shortest size interval τ, and that $q\tau < 1$, and that N such intervals can be measured. The resulting sample space is $\mathbb{Z}^N_{\geqslant 0}$, and the min-entropy of the resulting distribution is:

$$H_\infty(q, \tau, N) = Nq\tau \log_2(e) \tag{6.89}$$

because in each τ-interval the count maximizing the probability is $x = 0$, and the intervals are independent counts. For starting interval of length t, we can choose $\tau t/N$. So (6.89), which holds whenever $\tau < 1/q$, bounds the min-entropy to $qt \log_2(e)$, and decreasing τ further below $1/q$ does not boost the min-entropy. The greater precision does not increase the min-entropy, because even though most of the time a large amount of information from the occurrence of real number in the interval, the probability that $x \cap [a, b]$ is empty remains fixed and does not depend on the precision, and this determines the min-entropy.

In other words, a Poisson process gives a realistic example of a distribution that has a relatively large spike. As the precision goes to the zero, the Shannon entropy can go to infinity, yet the min-entropy remains bounded.

Acknowledgments

Aleksandra Zivkovic, Alfred Menezes, Greg Zaverucha and Adrian Antipa provided helpful comments.

References

[FIPS 140-1] National Institute of Standards and Technology. *Security Requirements for Cryptographic Modules*, Federal Information Processing Standard 140-1, 1994. csrc.nist.gov/groups/ST/toolkit/random_number.html.

[NIST 800-90] E. Barker and J. Kelsey. *Recommendation for Random Number Generation Using Deterministic Bit Generators*, Special Publication 800-90. National Institute of Standards and Technology, Mar. 2007. csrc.nist.gov/groups/ST/toolkit/random_number.html.

[X9.82] Accredited Standards Committee X9F1. *Draft American National Standard X9.82 (Random Number Generation), Part 2, Entropy Sources*, Jun. 2005.

[BG07] D. R. L. Brown and K. Gjøsteen. *A security analysis of the NIST SP 800-90 elliptic curve random number generator*. In A. J. Menezes (ed.), *Advances in Cryptology — CRYPTO 2007*, Lecture Notes in Computer Science 4622, pp. 466–481. International Association for Cryptologic Research, Springer, Aug. 2007. http://eprint.iacr.org/2007/048.

[BH05] B. Barak and S. Halevi. *A model and architecture for pseudo-random generation with applications to `/dev/random`*. In C. Meadows (ed.), *Proceedings of the 12th ACM conference on Computer and communications security*, pp. 203–212. ACM, Nov. 2005.

[BLMT11] M. Baudet, D. Lubicz, J. Micolod and A. Tassiaux. *On the security of oscillator-based random number generators*. J. of Cryptology, **24**(2):398–425, Apr. 2011.

[BPSW70] L. E. Baum, T. Petrie, G. Soules and N. Weiss. *A maximization technique occurring in the statistical analysis of probabilistic functions of markov chains*. The Annals of Mathematical Statistics, **41**(1):164–171, 1970.

[Cac97] C. Cachin. *Smooth entropy and Rényi entropy*. In W. Fumy (ed.), *Advances in Cryptology – EUROCRYPT '97*, Lecture Notes in Computer Science 1233, pp. 193–208. International Association for Cryptologic Research, Springer, May 1997.

[GJ83] I. P. Goulden and D. M. Jackson. *Combinatorial Enumeration*. Dover, 1983.

[JAW+00] T. Jennewein, U. Achleitner, G. Weihs, H. Weinfurter and A. Zeilinger. *A fast and compact quantum random number generator*. Review of Scientific Instruments, **71**(4):1675–1680, 2000.

[JJSH98] A. Juels, M. Jakobsson, E. Shrver and B. K. Hillyer. *How to turn loaded dice into fair coins*. IEEE Transactions on Information Theory, 1998.

[Lub96] M. Luby. *Pseudorandomness and Cryptographic Applications*. Princeton University Press, 1996.

[Mac95] I. G. Macdonald. *Symmetric Functions and Hall Polynomials*. Oxford University Press, second edn., 1995.

[Mau90] U. Maurer. *A universal statistical test for random bit generators*. In A. J. Menezes and S. A. Vanstone (eds.), *Advances in Cryptology — CRYPTO '90*, Lecture Notes in Computer Science 537, pp. 409–420. International Association for Cryptologic Research, Springer, Aug. 1990.

[MvOV97] A. J. Menezes, P. C. van Oorschot and S. A. Vanstone. *Handbook of Applied Cryptography.* CRC Press, 1997.

[Rén60] A. Rényi. *On measures of entropy and information.* In *Proceedings of the 4th Symposium on Mathematics, Statistics and Probability*, pp. 547–561. 1960.

[Sch01] C. P. Schnorr. *Small generic hardcore subsets for the discrete logarithm: Short secret Dl-keys.* Information Processing Letters, **79**:93–98, 2001.

[SMS07] B. Sunar, W. J. Martin and D. R. Stinson. *A provably secure true random number generator with built-in tolerance to active attacks.* IEEE Transactions on Computers, **56**(1):109–119, Jan. 2007.

A Optimization Methods

This section briefly reviews a few results from the theory of optimization. Optimization problems arise often in statistical inference, such as in maximally graded inference methods. Optimization also arises in entropy assessment when narrowing by minimization a set-valued inference about min-entropy into a single-valued inference.

A.1 Karush-Kuhn-Tucker Condition

Here we state the well-known Karush-Kuhn-Tucker theorem and conditions. Suppose that we want to minimize a function f subject to the constraints $g_i(x) \leqslant 0$ and $h_j(x) = 0$, for various indices i and j. Suppose that $\hat{x}$ is a local minimum of f within the constrained space of f. Suppose that f, g_i, h_j satisfy certain regularity conditions at $\hat{x}$. Then there exists μ_i and λ_j such that the following holds:

$$\nabla f(\hat{x}) + \sum_i \mu_i \nabla g_i(\hat{x}) + \sum_j \lambda_j \nabla h_j(\hat{x}) = 0, \tag{A.1}$$

$$\mu_i \geqslant 0, \tag{A.2}$$

$$\sum_i \mu_i g_i(\hat{x}) = 0. \tag{A.3}$$

These three conditions are called *stationarity*, *dual feasibility* and *complementary slackness*, respectively. By definition, we also have

$$g_i(\hat{x}) \leqslant 0, \tag{A.4}$$

$$h_i(\hat{x}) = 0, \tag{A.5}$$

which together are called the condition *primal feasibility*. These four conditions together are called the *Karush-Kuhn-Tucker* (KKT) conditions.

Several different regularity conditions exist such that a local minimum $\hat{x}$ ensures the KKT conditions.

Unless otherwise noted, the regularity conditions that we will use are as follows. The first part of regularity is that all f and g_i and h_j must be continuously differentiable. We say g_i is *active* at x when $g_i(x) = 0$. A second part of regularity is that the set consisting of all the gradients of the active g_i at $\hat{x}$ together the gradients of all the h_j is a linearly independent set of vectors.

A.2 Optimizing Non-Smooth and Non-Continuous Functions

Min-entropy is generally not a smooth function of p. More precisely, it is only piece-wise smooth, and does not have a well-defined gradient at some points. Typical optimization method, such as those the KKT conditions, use gradients. One approach to deal with the non-smoothness is to note that min-entropy may be viewed as the minimum of a number of smooth functions. Then one can optimize each such smooth function separately.

Generally, one is minimizing min-entropy over some set of inferred of distributions (either a maximally graded or threshold graded). In some case, this inferred set of distributions is symmetric with respect to the entries of the probability distribution vectors, so that it suffices to optimize just a single of the many smooth functions mentioned above.

Typicality, such as inclusive or balanced typicality can not only be non-smooth, but can also be non-continuous. It may be possible to handle such non-continuous functions by breaking the optimization problem into multiple variations, based on the case corresponding to each piece. As an alternative, this report has suggested forms of generalized typicality which can be chosen to be continuous and smooth, such as adjusted likelihood.

A.3 Model Constraints

A possible approach to handle the optimization problems arising from statistical inference, such as maximizing a grading or minimizing the entropy over a grading-thresholded set, is to parametrize the probability space Π using one coordinate for sample value x, with the coordinate taking the value $P_p(x)$.

In this approach, the probability model may be viewed as a set of equality constraints on these coordinates. This approach uses a lot of variables, but has the potential of simplifying the various functions involved.

A.4 Optimizations for the Independent Model

The KKT conditions simplify in the case of the independent model. First we revert the general KKT notation so that instead of optimizing over a vector x, we optimize over a vector $p = (p_0, \ldots, p_{m-1})$.

There m inequality constraints defined by $g_i(p) = -p_i \leqslant 0$ and one equality constraint $h(p) = p_0 + p_1 + \cdots + p_m - 1 = 0$. If f is continuously differentiable, then we note that the regularity conditions described above are met. This is because at most $m-1$ inequality constraints can be active and the gradients are thus obviously independent. Therefore, at a local minimum, the KKT conditions will hold.

Upon elimination the μ_i and λ, the KKT conditions are equivalent to the following conditions. For $0 \leqslant i \leqslant m-1$,

$$(\nabla f(\hat{p}))_i \geqslant (\nabla f(\hat{p})) \cdot \hat{p} \tag{A.6}$$

where $(\nabla f(\hat{p}))_i$ is the i^{th} entry in the vector $\nabla f(\hat{p})$ and $(\nabla f(\hat{p})) \cdot p$ is the usual dot product of vectors.

Remark A.1. To see how to explicitly eliminate the intermediate KKT variables μ_i and λ, do as follows. The gradients of the constraints are given by $\nabla g_i = -e_i$ where e_i is the elementary vector with value 1 in position i and value 0 elsewhere; and $\nabla h = \sum_{i=0}^{m-1} e_i$.

Apply the dot product of the stationarity condition (A.1) with $\hat{p}$. The complementary slackness (A.3) eliminates each contributions of $\mu_i \nabla g_i \cdot \hat{p} = -\mu_i e_i \cdot p = -\mu_i p_i = \mu_i g_i(\hat{p})$. The contribution from the equality constraint is $\lambda \nabla h \cdot \hat{p} = \lambda \sum_{i=0}^{m-1} p_i = \lambda$. It follows that $\lambda = -\nabla f(\hat{p})$.

Apply the dot product of the stationarity condition (A.1) with e_i, to get $(\nabla f(\hat{p}))_i + \mu_i + \lambda$, which gives (A.6).

To see (A.6) directly, note that (A.6) is equivalent to the condition that the objective f is nondecreasing along each line ray emanating from $\hat{p}$ and heading towards a vertex of the simplex Π, that is a distribution $p^{(i)} \in \Pi$ with $p^{(i)}_j$ equal to 1 if $i = j$ and equal to 0 otherwise. If the objective function is continuously differentiable and $\hat{p}$ is a local minimum, then clearly f will be non-decreasing along each such ray. The derivative along the ray is $(\nabla f(\hat{p})) \cdot (p^{(i)} - \hat{p})$. Taking the

Remark A.2. The converse, however, may not hold: conditions (A.6) do not suffice to ensure a local minimum at $\hat{p}$. If $\hat{p}$ is not a local minimum, but f is continuously differentiable at $\hat{p}$, then f has a saddle at $\hat{p}$: in some directions f increases and in other directions f decreases.

Remark A.3. As an example to consider, suppose that we are in the $(2, 3)$ independent model, and we want to maximize the likelihood function $L_{011}(p) = p_0 p_1^2$. Put $f = -L_{011}$, and (A.6) works to the following two inequalities:

$$-p_1^2 \geqslant -3p_0 p_1^2, \tag{A.7}$$

$$-2p_0 p_1 \geqslant -3p_0 p_1^2 \tag{A.8}$$

which becomes respectively

$$p_1^2(3p_0 - 1) \geqslant 0, \tag{A.9}$$

$$p_0 p_1(3p_1 - 2) \geqslant 0. \tag{A.10}$$

Since $p_0 \geqslant 0$, the first inequality implies $p_1 = 0$ or $p_0 \geqslant 1/3$. Since $p_0, p_1 \geqslant 0$, the second implies that $0 \in \{p_0, p_1\}$ or $p_1 \geqslant 2/3$. The only (p_0, p_1) that meet these conditions are $(1, 0)$, $(1/3, 2/3)$ and $(0, 1)$.

Remark A.4. When the objective is more complicated, the gradient of the objective is usually correspondingly more complicated. In some cases, it may be better to add some additional variables to the optimization problem: variables that represent some intermediate values in the objective function. Being intermediate values, these additional variables are governed by some additional equality constraints. An example may help.

In the independent model, the given variables are $p_0, \ldots, p_{m-1}$. The given constraints are m inequalities $p_i \geqslant 0$ and one equality $\sum_i p_i = 1$. First, we introduce m^N supplemental variables p_x for each $x \in X$. (Similar to what was suggested in §A.3.)

Second, we introduce m^N supplemental equality constraints $p_x = P_p(x)$, which may be explicitly given by the polynomial equations:

$$p_{x_0} p_{x_1} \cdots p_{x_{N-1}} - p_x = 0 \tag{A.11}$$

Although this introduces many variables and constraints, a benefit may be to simplify the objective function. To maximize likelihood L_x, the objective function is simply $-p_x$. To maximize adjusted likelihood, it suffices to minimize the objective function is $-p_x + \sum_y p_y^2$, a quadratic function in the variables.

B Modeling

This report concerns the task of formally assessing cryptographic entropy of a source. A pre-requisite to this task is that a probability model appropriate for the source has been selected. Selection of the model is not the main focus of this report. This section briefly describes approaches to selecting a probability model for a source. Two types of approaches are outline below.

B.1 Relaxation Approach to Modeling

In the *relaxation* approach, one selects as an initial model a very restricted model. Either the restricted model could be selected as the ideal for the intended use of the source, such as being the uniform model, or as the model that hypothetically describes the source in most detail, such as a deterministic model. Next, one does hypothesis testing on the initial model, as outlined in §C. If the model is rejected, one must select another model. In the relaxation approach, a relaxation of model initial model is selected. The choice of relaxation requires some inspiration or intuition. If one chooses the relaxation of the model before hypothesis testing, then one can do comparative hypothesis testing, which is the method preferred by this report. The relxation approach can be iterated. Examples of the relaxation approach to modeling are given in §6.5 and §6.6.

Section 6.5 deals with dice, and starts with an initial model which is the uniform model. It uses comparative hypothesis testing with the alternative model being the independent model. For one sample, it rejects the uniform model. In accordance with the relaxation approach to modeling, the model is relaxed from the uniform model to the independent model, and then the independent model is tested. Again, comparative testing is used, with the alterative model being the Markov model. In this case, the independent model is accepted.

Section 6.6 deals with ring oscillators. It starts with an initial model in which the bit sequence produced by a ring oscillator has a simple description determined by its frequency. In this case, the model has very low entropy, so the starting point is not the most optimistic one from the perspective of use the source for cryptographic entropy. (It is optimistic from the perpsective of an adversary.)

Two disadvantages of the relaxation approach are:

- The relaxation approach risks being over-optimistic, which could occur if, firstly, hypothesis testing yields a false acceptance, meaning that source has a distribution p which is not contained in the tested model, and if, secondly, the distribution p has significantly lower entropy than what would be inferred using the falsely accepted model.
- In the event that a tested model is rejected, the relaxation approach provides no formally quantified guidance on how to relax the probably model. The choice of relaxation is outside the scope of the formal techniques in this report.

B.2 Restrictive Approach to Modeling

In the *restriction* approach to modeling, one starts from some a very relaxed model for the source. The selection of this model must based on intuition or inspiration, but the goal is to think only about the source, and to capture all the possible ways in which the source could be descibed, making as minimal assumptions as possible. Next, one makes infer the distributions in the relaxed model, using the formal methods described in this reaport. These inferences will depend on the observed sample.

The restriction approach tries to derive from the formal inference a new model. For example, a set-valued inference on the distribution may itself be thought as a new model, restricting the original model. The restriction approach is not just to use this inference as a restricted model, because then all one has done really is inference. Instead, the restriction approach uses the formal inferences as inspiration to hypothesize a different restricted model.

It may be noted that a simple model or distribution, such as the uniform distribution, can yield sample values that with high complexity. A formal inference made from a complex is liekly to have high complexity. The restriction approach is to choose take complex inference about the distribution and to find a simpler description of the inference and express this as another probability. This simplification step may require some inspiration, and is outside the main scope of this report.

Any such restriction of the model must be also subjected to hyptothesis testing. Comparative hypothesis testing using the original model as the alternative may be applied. If the restricted model is accepted, then the process can be iterated.

The relaxation and restrication approaches can be mixed. Indeed, they are not entirely different, since they both involve steps of selecting models that are relaxed and restricted.

Remark B.1. In the restriction approach, one should not start from a model is relaxed but not too relaxed, otherwise the inferences may be too weak.

Remark B.2. For example, to model ring oscillators, one can gather multiple ring oscillators manufactured by the same process. For an initial relaxed model, one might model their outputs as indpendent from each other, and furthermore assume that each has the same distribution. So, one is formulating the common power of the models. If one does not assume anything about each individual oscillator, then one essentially has the independent model. So, if one takes 32 ring oscillators, and looks at their first 2^{10} output bits, then one could take as one is starting with a relaxed model in the restriction approach consisting the of $(2^{2^{10}}, 32)$ independent model. The large is due there being $2^{2^{10}}$ possible outputs of the first 2^{10}. The length 32 is much shorter than the width, so the sample size is very low. Formal inferences that can can say very little therefore.

Essentially, in this case, the formal inferences may so modest, the sample values themselves should be used directly for inspiration of how to hypothesis a restriction of the model.

Remark B.3. Remark B.2 uses initial model that may be more relaxed than necessary. For example, it completely ignores that the 2^{10} bits produced obtained from each ring oscillator sample are produced chronologically. It seems a mild assumption that the initial bits could not depend on the latter bits. So, a hidden Markov model could be hypothesized as an initial model for each individual oscillator. In this case, the formal initial model would the common power of 32 copies of the hidden Markov model. The larger the size of the hidden state for the Markov model, the more relaxed the model would be.

C Hypothesis Testing

This report focuses on the task of formally assessing cryptographic entropy of a source. As noted in §B, A prerequisite to this task relies is that a probability model appropriate for the rousec has been selected. An entropy assesment of a source can only be as appropriate for the source as the model is appropriate.

In this report, *hypothesis testing* is to determine whether a given probability model is appropriate for a given source. For example, extensive hyptothesis testing may provide the confidence in the probability model for a given source, thereby providing confidence in entropy assessments for the source.

Hypothesis testing of a model on a source requires one to gather a sample value x from the source. Two types of hypothesis testing are conisdred.

Hypothesis testing has the risk that the model can be falsely accepted. For example, if the model contains the uniform distribution, but the source is deterministic but pseudorandom, then it is unlikely to be rejected. In other words, a badly seeded cryptographic number generator would pass any general hypothesis test. The only way to truly overcome this risk is to ensure that both hypothesis testing and cryptoraphic entropy asessment look at sample values drawn from the actual source of entropy, before any cryprographic processing.

C.1 Non-Comparative Hypothesis Testing

In *non-comparative* hypothesis testing does not rely on any other models. In the formalism of this report, one evaluates the maximal typicality of x, perhaps as induced by a sample statistic. If the maximal typicality of x is too low, then one rejects the model. Some difficulties with non-comparative hypothesis testing are:

- The model may be subject to the tying effect may be make rejection impossible for any x. Choosing an arbitrary tiebreaker sample statistic can overcome this, but only at the cost making an arbitrary choice of sample statistic, and the risk of making an artificial choice of statistic.
- The model has a probability of being falsely rejected, depending the threshold for the maximal typicality value used for rejection. In practice, this means setting the threshold very low.
- Setting a very low threshold for rejection risks that the may exists a distribution outside the model which is much better fit for the source, but would not be detected.
- If the model is rejected, then one has to start from scratch.

C.2 Comparative Hypothesis Testing

In *comparative* hypothesis testing, the hypothesized model is tested against an alternative model which is relaxation of the hypothesized model. Comparative hypothesis testing is an attempt to address some of the difficulties with non-compartive hypothesi testing.

In *comparative* hypothesis testing, one computes the maximal typicality of the x using a sample statistic that is model-neutral with respect to the alternative model. One rejects the hypothesis if the maximal typicality obtained is below some threshold. In this case, rejection means to change the hypothesis into the alternative model instead of the hypothesized model. In comparative hyptothesis one can set the threshold much higher than in non-comparative testing, because the cost of false rejection is only to relax the model to the alternative model.

D Game-Theoretic Analysis

In this section, we consider a situation in which the adversary can choose the distribution $p \in \Pi$, and then tries to guess the sample value x drawn from p. But the entropy assessor has an opportunity to make statistical inference about p. The assessor must bound the chance the adversary wins.

For example, suppose that the system is using prospective inference, with the first sample used for assessment, and the second for deployment. (In other words, the overall source model is a common product (Π, X^2, R) of the model (Π, X, P) with itself. The game works as follows:

1. The adversary chooses $p \in \Pi$.
2. A sample (x_1, x_2) is drawn in the model (Π, X^2, R).
3. The assessor is given x_1.
4. The assessor outputs an entropy estimate $H \in \mathbb{R}$.
5. The adversary guesses a value y.
6. If $y = x_2$, the score of the game is $s_1(H)$.
7. If $y \neq x_2$, the score of the game is $s_0(H)$.

The assessor tries to maximize the score, while the adversary tries to minimize the score. The strategies of the assessor and the adversary in this game depend on the scoring function $s_0, s_1 : \mathbb{R} \to \mathbb{R}$.

More formally, the game can be viewed as a two-player game in which the two players make choices simultaneously. The adversary choice consists of the pair (p, y), a distribution and a sample value. The assessor's choice consists of a entropy assessment function $H : X \to \mathbb{R}$. The score of the game is then a random variable, taking values in $\mathbb{R}$. Although the range of the score is a continuum, the random variable is discrete if the set X is finite. The probability that a score results in s is

$$P_p(y) \left(\sum_{x : s = s_1(H(x))} P_p(x) \right) + (1 - P_p(y)) \left(\sum_{x : s = s_0(H(x))} P_p(x) \right) \tag{D.1}$$

As mentioned earlier, the adversary wishes to minimize the score, while the assessor wishes to maximize the score. But since, since the score is a random variable, maximization and minimization are not clearly defined.

For the sake of simplicity, suppose that both the adversary and the assessor attempt to optimize (in different directions) the expected value of the resultant score. The expected value of the score is

$$\sum_{x \in X} P_p(x) \left(P_p(y) s_1(H(x)) + (1 - P_p(y)) s_0(H(x)) \right). \tag{D.2}$$

Note that the summation index x corresponds to the observed value x_1 in the step-by-step game above.

For the sake of concreteness, now suppose that (Π, X, P) is the $(2, 3)$ independent model, and that the scoring functions are $s_0(h) = h$ and $s_1(h) = -h$. The objective function for both players, the expected score, then simplifies to:

$$1 - 2P_p(y) \sum_{x \in X} P_p(x) H(x) \tag{D.3}$$

In cryptographic applications, the assessor's only source of randomness is the source. So, effectively, we can assume that the assessor must fix the choice of assessment function H. In game theory terminology, the assessor is forced to use a pure strategy. The adversary may use a mixed strategy. For example, the choice of (p, y) in the game may not be fixed, but actually drawn from a distribution.

E Estimation Theory

Estimation theory is an approach to statistical inference that takes a given inference method, and produces an evaluation of its quality.

For example suppose that (Π, X, P) and $r : \Pi \to R$ is a parameter, and $i : X \to R$ is a inference function. Furthermore, suppose that the space R is a convex space, in the sense that convex combinations are defined in R. Then the inference function i is said to be *unbiased estimator* for r if, for all $p \in \Pi$, it is true that

$$E(i(x)) = r(p), \tag{E.1}$$

where E is the expected value of $i(x)$ according to the probability distribution p, which means that:

$$E(i(x)) = \sum_{x \in X} P_p(x) i(x). \tag{E.2}$$

Remark E.1. The parameter min-entropy parameter H_∞ is a non-polynomial function of $p \in \Pi$. The left hand side of (E.1) is a polynomial function in p, as seen by the definition (E.2). Therefore, for almost all choices of Π, no inference function is an unbiased estimator for H_∞.

Remark E.2. The maximum likelihood estimate (inference) is an unbiased estimator for the probability distribution p itself in the unrestricted model.

Remark E.3. Suppose that we have two models (Π, X, P) and (Π, Y, Q), with a shared probability space. Suppose we have probability parameters $r : \Pi \to R$ and $s : \Pi \to S$ and inference functions $i : X \to R$ and $j : Y \to S$. Recall that we defined there product model $(\Pi, X \times Y, P \times Q)$ such that $(P \times Q)_p(x, y) = P_p(x) Q_p(y)$. Similarly, we may define the product of the parameters $r \times s : \Pi \to R \times S : p \mapsto (r(p), s(p))$ and the product of the inference functions $i \times j : X \times Y \to R \times S : (x, y) \mapsto (i(x), j(y))$. Then $r \times s$ is a parameter for the product model and $i \times j$ is an inference function for the product model with the same range as the product parameter, namely $R \times S$. If i and j are unbiased estimator for r and s respectively, then $i \times j$ is an unbiased estimator for $r \times s$

Remark E.4. The maximum likelihood inference function is an unbiased estimator of the probability distribution itself in the (m, N) independent model. It is easy to see that the maximum likelihood inference function i is defined such that $i(x) = f(x)$ where f is the frequency vector of the sample vector x, meaning that, $f(x)_i = j/N$ if the number of k such that $x_k = i$ is j. To show that this is unbiased, by symmetry, it suffices to show that the expected value of $f(x)_0$ is p_0. The expected value of $f(x)_0$ is

$$\begin{aligned}
f(x)_0 &= \sum_{x_0, x_1, \ldots, x_{N-1}} p_{x_0} p_{x_1} \cdots p_{x_{N-1}} \left| \{k : x_k = 0\} \right| / N \\
&= \tfrac{1}{N} \sum_{x_0, x_1, \ldots, x_{N-1}} p_0 \frac{\partial}{\partial p_0} p_{x_0} p_{x_1} \cdots p_{x_{N-1}} \\
&= \tfrac{1}{N} p_0 \frac{\partial}{\partial p_0} \sum_{x_0, x_1, \ldots, x_{N-1}} p_{x_0} p_{x_1} \cdots p_{x_{N-1}} \\
&= \tfrac{1}{N} p_0 \frac{\partial}{\partial p_0} (p_0 + \cdots + p_{m-1})^N \\
&= \tfrac{1}{N} p_0 N (p_0 + \cdots + p_{m-1})^{N-1} \frac{\partial p_0}{\partial p_0} \\
&= p_0,
\end{aligned} \tag{E.3}$$

where the p_i are treated as indeterminates when calculating partial derivatives, then treated again as probabilities in the final steps. The sums are over $x \in X$, meaning $0 \leqslant x_0, \ldots, x_{N-1} \leqslant m - 1$, with the $x_i \in \mathbb{Z}$.

Other measures of the quality of the estimator can be defined given a metric d on the space R. For each p, we can define the error of i as an estimator of r:

$$\epsilon_e(p, i, r) = E(d(i(x) - r(p))^e) \tag{E.4}$$

where e would typically be one or two. However, in statistical inference, p is unknown, so therefore so is the error above. What one can do is take the maximum error over all p as the total error of the inference i of r. Or, what one can do is, if Π is equipped with a measure, take the average error over Π.

These notions suggest that one can define an inference function in terms of achieving best quality. For example, perhaps choose an unbiased estimator, if possible. Generally, among the remaining choices, choose an inference function that produces the least total error, or average error.

Remark E.5. The suitability of estimation theory for cryptology is unclear, primarily because of Remark E.1.

Furthermore, all estimation methods above use expectation, and thus averages over the sample space X. It is a theme of cryptology that danger lurks in using averages, because an adversary, unlike a natural process, will not confine itself to random behavior, and thus to average behavior. An adversary will search over the sample space X, so averages over sample space X may not be a good measure of anything.